

Linear Algebra Review

CS221: Introduction to Artificial Intelligence

Carlos Fernandez-Granda

10/11/2011

Index

- 1 Vectors
- 2 Matrices
- 3 Matrix Decompositions
- 4 Application : Image Compression

- 1 Vectors
- 2 Matrices
- 3 Matrix Decompositions
- 4 Application : Image Compression

What is a vector?

What is a vector?

- An ordered tuple of numbers :


$$\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ u_n \end{pmatrix}$$

What is a vector?

- An ordered tuple of numbers :

$$\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ u_n \end{pmatrix}$$

- A quantity that has a magnitude and a direction.



Vector spaces

- More formally a vector is an element of a **vector space**.

Vector spaces

- More formally a vector is an element of a **vector space**.
- A vector space \mathcal{V} is a set that contains **all linear combinations** of its elements :

Vector spaces

- More formally a vector is an element of a **vector space**.
- A vector space \mathcal{V} is a set that contains **all linear combinations** of its elements :
 - If vectors \mathbf{u} and $\mathbf{v} \in \mathcal{V}$, then $\mathbf{u} + \mathbf{v} \in \mathcal{V}$.

Vector spaces

- More formally a vector is an element of a **vector space**.
- A vector space \mathcal{V} is a set that contains **all linear combinations** of its elements :
 - If vectors \mathbf{u} and $\mathbf{v} \in \mathcal{V}$, then $\mathbf{u} + \mathbf{v} \in \mathcal{V}$.
 - If vector $\mathbf{u} \in \mathcal{V}$, then $\alpha \mathbf{u} \in \mathcal{V}$ for any scalar α .

Vector spaces


- More formally a vector is an element of a **vector space**.
- A vector space \mathcal{V} is a set that contains **all linear combinations** of its elements :
 - If vectors \mathbf{u} and $\mathbf{v} \in \mathcal{V}$, then $\mathbf{u} + \mathbf{v} \in \mathcal{V}$.
 - If vector $\mathbf{u} \in \mathcal{V}$, then $\alpha \mathbf{u} \in \mathcal{V}$ for any scalar α .
 - There exists $\mathbf{0} \in \mathcal{V}$ such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$ for any $u \in \mathcal{V}$.

Vector spaces

- More formally a vector is an element of a **vector space**.
- A vector space \mathcal{V} is a set that contains **all linear combinations** of its elements :
 - If vectors \mathbf{u} and $\mathbf{v} \in \mathcal{V}$, then $\mathbf{u} + \mathbf{v} \in \mathcal{V}$.
 - If vector $\mathbf{u} \in \mathcal{V}$, then $\alpha \mathbf{u} \in \mathcal{V}$ for any scalar α .
 - There exists $\mathbf{0} \in \mathcal{V}$ such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$ for any $u \in \mathcal{V}$.
- A **subspace** is a subset of a vector space that is **also** a vector space.


Examples

- Euclidean space \mathbb{R}^n .



Examples

- Euclidean space \mathbb{R}^n .




- The **span** of any set of vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$, defined as :

$$\text{span}(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n) = \{\alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \dots + \alpha_n \mathbf{u}_n \mid \alpha_i \in \mathbb{R}\}$$


Subspaces

- A **line** through the origin in \mathbb{R}^n (span of a vector).




Subspaces

- A **line** through the origin in \mathbb{R}^n (span of a vector).



- A **plane** in \mathbb{R}^n (span of two vectors).



Linear Independence and Basis of Vector Spaces

- A vector \mathbf{u} is linearly independent of a set of vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ if it does **not** lie in their **span**.

Linear Independence and Basis of Vector Spaces

- A vector \mathbf{u} is linearly independent of a set of vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ if it does **not** lie in their **span**.
- A set of vectors is linearly independent if every vector is linearly independent of the rest.

Linear Independence and Basis of Vector Spaces

- A vector \mathbf{u} is linearly independent of a set of vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ if it does **not** lie in their **span**.
- A set of vectors is linearly independent if every vector is linearly independent of the rest.
- A **basis** of a vector space \mathcal{V} is a linearly independent set of vectors whose **span is equal to \mathcal{V}** .

Linear Independence and Basis of Vector Spaces

- A vector \mathbf{u} is linearly independent of a set of vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ if it does **not** lie in their **span**.
- A set of vectors is linearly independent if every vector is linearly independent of the rest.
- A **basis** of a vector space \mathcal{V} is a linearly independent set of vectors whose **span is equal to** \mathcal{V} .
- If a vector space has a basis with d vectors, its **dimension** is d .

Linear Independence and Basis of Vector Spaces

- A vector \mathbf{u} is linearly independent of a set of vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ if it does **not** lie in their **span**.
- A set of vectors is linearly independent if every vector is linearly independent of the rest.
- A **basis** of a vector space \mathcal{V} is a linearly independent set of vectors whose **span is equal to \mathcal{V}** .
- If a vector space has a basis with d vectors, its **dimension** is d .
- Any other basis of the same space will consist of **exactly d** vectors.

Linear Independence and Basis of Vector Spaces

- A vector \mathbf{u} is linearly independent of a set of vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ if it does **not** lie in their **span**.
- A set of vectors is linearly independent if every vector is linearly independent of the rest.
- A **basis** of a vector space \mathcal{V} is a linearly independent set of vectors whose **span is equal to** \mathcal{V} .
- If a vector space has a basis with d vectors, its **dimension** is d .
- Any other basis of the same space will consist of **exactly d** vectors.
- The dimension of a vector space can be infinite (function spaces).

Norm of a Vector

- A norm $\|\mathbf{u}\|$ measures the **magnitude** of a vector.

Norm of a Vector

- A norm $\|\mathbf{u}\|$ measures the **magnitude** of a vector.
- Mathematically, any function from the vector space to \mathbb{R} that satisfies :

Norm of a Vector

- A norm $\|\mathbf{u}\|$ measures the **magnitude** of a vector.
- Mathematically, any function from the vector space to \mathbb{R} that satisfies :
 - Homogeneity $\|\alpha \mathbf{u}\| = |\alpha| \|\mathbf{u}\|$

Norm of a Vector


- A norm $\|\mathbf{u}\|$ measures the **magnitude** of a vector.
- Mathematically, any function from the vector space to \mathbb{R} that satisfies :
 - Homogeneity $\|\alpha \mathbf{u}\| = |\alpha| \|\mathbf{u}\|$
 - Triangle inequality $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$

Norm of a Vector


- A norm $\|\mathbf{u}\|$ measures the **magnitude** of a vector.
- Mathematically, any function from the vector space to \mathbb{R} that satisfies :
 - Homogeneity $\|\alpha \mathbf{u}\| = |\alpha| \|\mathbf{u}\|$
 - Triangle inequality $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$
 - Point separation $\|\mathbf{u}\| = 0$ if and only if $\mathbf{u} = \mathbf{0}$.

Norm of a Vector

- A norm $\|\mathbf{u}\|$ measures the **magnitude** of a vector.
- Mathematically, any function from the vector space to \mathbb{R} that satisfies :
 - Homogeneity $\|\alpha \mathbf{u}\| = |\alpha| \|\mathbf{u}\|$
 - Triangle inequality $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$
 - Point separation $\|\mathbf{u}\| = 0$ if and only if $\mathbf{u} = 0$.
- Examples :
 - Manhattan or ℓ_1 norm : $\|\mathbf{u}\|_1 = \sum_i |u_i|$.
 - Euclidean or ℓ_2 norm : $\|\mathbf{u}\|_2 = \sqrt{\sum_i u_i^2}$.



Manhattan




Euclidean

Inner Product

- Inner product between \mathbf{u} and \mathbf{v} : $\langle \mathbf{u}, \mathbf{v} \rangle = \sum_{i=1}^n u_i v_i$.


Inner Product

- Inner product between \mathbf{u} and \mathbf{v} : $\langle \mathbf{u}, \mathbf{v} \rangle = \sum_{i=1}^n u_i v_i$.
- It is the **projection** of one vector onto the other.



Inner Product




- Inner product between \mathbf{u} and \mathbf{v} : $\langle \mathbf{u}, \mathbf{v} \rangle = \sum_{i=1}^n u_i v_i$.
- It is the **projection** of one vector onto the other.



- Related to the Euclidean norm : $\langle \mathbf{u}, \mathbf{u} \rangle = \|\mathbf{u}\|_2^2$.




Inner Product

The inner product is a measure of **correlation** between two vectors, scaled by the norms of the vectors :



Inner Product




The inner product is a measure of **correlation** between two vectors, scaled by the norms of the vectors :



- If $\langle \mathbf{u}, \mathbf{v} \rangle > 0$, \mathbf{u} and \mathbf{v} are **aligned**.

Inner Product




The inner product is a measure of **correlation** between two vectors, scaled by the norms of the vectors :



- If $\langle \vec{u}, \vec{v} \rangle > 0$, \vec{u} and \vec{v} are **aligned**.
- If $\langle \vec{u}, \vec{v} \rangle < 0$, \vec{u} and \vec{v} are **opposed**.

Inner Product

The inner product is a measure of **correlation** between two vectors, scaled by the norms of the vectors :



- If $\langle \mathbf{u}, \mathbf{v} \rangle > 0$, \mathbf{u} and \mathbf{v} are **aligned**.
- If $\langle \mathbf{u}, \mathbf{v} \rangle < 0$, \mathbf{u} and \mathbf{v} are **opposed**.
- If $\langle \mathbf{u}, \mathbf{v} \rangle = 0$, \mathbf{u} and \mathbf{v} are **orthogonal**.

Orthonormal Basis

- The vectors in an **orthonormal** basis have unit Euclidean norm and are orthogonal to each other.

Orthonormal Basis

- The vectors in an **orthonormal** basis have unit Euclidean norm and are orthogonal to each other.
- To express a vector x in an orthonormal basis, we can just take inner products.

Orthonormal Basis

- The vectors in an **orthonormal** basis have unit Euclidean norm and are orthogonal to each other.
- To express a vector \mathbf{x} in an orthonormal basis, we can just take inner products.
- Example : $\mathbf{x} = \alpha_1 \mathbf{b}_1 + \alpha_2 \mathbf{b}_2$.

Orthonormal Basis

- The vectors in an **orthonormal** basis have unit Euclidean norm and are orthogonal to each other.
- To express a vector \mathbf{x} in an orthonormal basis, we can just take inner products.
- Example : $\mathbf{x} = \alpha_1 \mathbf{b}_1 + \alpha_2 \mathbf{b}_2$.
- We compute $\langle \mathbf{x}, \mathbf{b}_1 \rangle$:

$$\begin{aligned}\langle \mathbf{x}, \mathbf{b}_1 \rangle &= \langle \alpha_1 \mathbf{b}_1 + \alpha_2 \mathbf{b}_2, \mathbf{b}_1 \rangle \\ &= \alpha_1 \langle \mathbf{b}_1, \mathbf{b}_1 \rangle + \alpha_2 \langle \mathbf{b}_2, \mathbf{b}_1 \rangle \quad \text{By linearity.} \\ &= \alpha_1 + 0 \quad \text{By orthonormality.}\end{aligned}$$

Orthonormal Basis

- The vectors in an **orthonormal** basis have unit Euclidean norm and are orthogonal to each other.
- To express a vector \mathbf{x} in an orthonormal basis, we can just take inner products.
- Example : $\mathbf{x} = \alpha_1 \mathbf{b}_1 + \alpha_2 \mathbf{b}_2$.
- We compute $\langle \mathbf{x}, \mathbf{b}_1 \rangle$:

$$\begin{aligned}\langle \mathbf{x}, \mathbf{b}_1 \rangle &= \langle \alpha_1 \mathbf{b}_1 + \alpha_2 \mathbf{b}_2, \mathbf{b}_1 \rangle \\ &= \alpha_1 \langle \mathbf{b}_1, \mathbf{b}_1 \rangle + \alpha_2 \langle \mathbf{b}_2, \mathbf{b}_1 \rangle \quad \text{By linearity.} \\ &= \alpha_1 + 0 \quad \text{By orthonormality.}\end{aligned}$$

- Likewise, $\alpha_2 = \langle \mathbf{x}, \mathbf{b}_2 \rangle$.

- 1 Vectors
- 2 Matrices**
- 3 Matrix Decompositions
- 4 Application : Image Compression

Linear Operator

- A linear operator $\mathcal{L} : \mathcal{U} \rightarrow \mathcal{V}$ is a map from a vector space \mathcal{U} to another vector space \mathcal{V} that satisfies :

Linear Operator

- A linear operator $\mathcal{L} : \mathcal{U} \rightarrow \mathcal{V}$ is a map from a vector space \mathcal{U} to another vector space \mathcal{V} that satisfies :
 - $\mathcal{L}(\mathbf{u}_1 + \mathbf{u}_2) = \mathcal{L}(\mathbf{u}_1) + \mathcal{L}(\mathbf{u}_2)$.

Linear Operator

- A linear operator $\mathcal{L} : \mathcal{U} \rightarrow \mathcal{V}$ is a map from a vector space \mathcal{U} to another vector space \mathcal{V} that satisfies :
 - $\mathcal{L}(\mathbf{u}_1 + \mathbf{u}_2) = \mathcal{L}(\mathbf{u}_1) + \mathcal{L}(\mathbf{u}_2)$.
 - $\mathcal{L}(\alpha \mathbf{u}) = \alpha \mathcal{L}(\mathbf{u})$ for any scalar α .

Linear Operator

- A linear operator $\mathcal{L} : \mathcal{U} \rightarrow \mathcal{V}$ is a map from a vector space \mathcal{U} to another vector space \mathcal{V} that satisfies :
 - $\mathcal{L}(\mathbf{u}_1 + \mathbf{u}_2) = \mathcal{L}(\mathbf{u}_1) + \mathcal{L}(\mathbf{u}_2)$.
 - $\mathcal{L}(\alpha \mathbf{u}) = \alpha \mathcal{L}(\mathbf{u})$ for any scalar α .
- If the dimension n of \mathcal{U} and m of \mathcal{V} are finite, \mathcal{L} can be represented by an $m \times n$ matrix \mathbf{A} .

$$\mathbf{A} = \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix}$$

Matrix Vector Multiplication

$$\mathbf{A}\mathbf{u} = \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ u_n \end{pmatrix} = \begin{pmatrix} v_1 \\ v_2 \\ \dots \\ v_m \end{pmatrix}$$

Matrix Vector Multiplication

$$\mathbf{A}\mathbf{u} = \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{pmatrix} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{pmatrix}$$

Matrix Vector Multiplication

$$\mathbf{A}\mathbf{u} = \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ \color{red}{A_{21}} & \color{red}{A_{22}} & \dots & \color{red}{A_{2n}} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} \color{red}{u_1} \\ \color{red}{u_2} \\ \dots \\ \color{red}{u_n} \end{pmatrix} = \begin{pmatrix} v_1 \\ \color{red}{v_2} \\ \dots \\ v_m \end{pmatrix}$$

Matrix Vector Multiplication

$$\mathbf{A}\mathbf{u} = \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ u_n \end{pmatrix} = \begin{pmatrix} v_1 \\ v_2 \\ \dots \\ v_m \end{pmatrix}$$

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}\mathbf{AB} &= \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} B_{11} & B_{12} & \dots & B_{1p} \\ B_{21} & B_{22} & \dots & B_{2p} \\ & & \dots & \\ B_{n1} & B_{n2} & \dots & B_{np} \end{pmatrix} \\ &= \begin{pmatrix} AB_{11} & AB_{12} & \dots & AB_{1p} \\ AB_{21} & AB_{22} & \dots & AB_{2p} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}\end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}
 \mathbf{AB} &= \begin{pmatrix} \textcolor{red}{A}_{11} & \textcolor{red}{A}_{12} & \dots & \textcolor{red}{A}_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} \textcolor{red}{B}_{11} & B_{12} & \dots & B_{1p} \\ \textcolor{red}{B}_{21} & B_{22} & \dots & B_{2p} \\ & & \dots & \\ \textcolor{red}{B}_{n1} & B_{n2} & \dots & B_{np} \end{pmatrix} \\
 &= \begin{pmatrix} \textcolor{red}{AB}_{11} & AB_{12} & \dots & AB_{1p} \\ AB_{21} & AB_{22} & \dots & AB_{2p} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}
 \end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}
 \mathbf{AB} &= \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} & \dots & \mathbf{A}_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} B_{11} & \mathbf{B}_{12} & \dots & B_{1p} \\ B_{21} & \mathbf{B}_{22} & \dots & B_{2p} \\ & & \dots & \\ B_{n1} & \mathbf{B}_{n2} & \dots & B_{np} \end{pmatrix} \\
 &= \begin{pmatrix} AB_{11} & \mathbf{AB}_{12} & \dots & AB_{1p} \\ AB_{21} & AB_{22} & \dots & AB_{2p} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}
 \end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}
 \mathbf{AB} &= \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} B_{11} & B_{12} & \dots & B_{1p} \\ B_{21} & B_{22} & \dots & B_{2p} \\ & & \dots & \\ B_{n1} & B_{n2} & \dots & B_{np} \end{pmatrix} \\
 &= \begin{pmatrix} AB_{11} & AB_{12} & \dots & AB_{1p} \\ AB_{21} & AB_{22} & \dots & AB_{2p} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}
 \end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}
 \mathbf{AB} &= \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ \color{red}{A_{21}} & \color{red}{A_{22}} & \dots & \color{red}{A_{2n}} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} \color{red}{B_{11}} & B_{12} & \dots & B_{1p} \\ \color{red}{B_{21}} & B_{22} & \dots & B_{2p} \\ & & \dots & \\ \color{red}{B_{n1}} & B_{n2} & \dots & B_{np} \end{pmatrix} \\
 &= \begin{pmatrix} AB_{11} & AB_{12} & \dots & AB_{1p} \\ \color{red}{AB_{21}} & AB_{22} & \dots & AB_{2p} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}
 \end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}
 \mathbf{AB} &= \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ \color{red}{A_{21}} & \color{red}{A_{22}} & \dots & \color{red}{A_{2n}} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} B_{11} & \color{red}{B_{12}} & \dots & B_{1p} \\ B_{21} & \color{red}{B_{22}} & \dots & B_{2p} \\ & & \dots & \\ B_{n1} & \color{red}{B_{n2}} & \dots & B_{np} \end{pmatrix} \\
 &= \begin{pmatrix} AB_{11} & AB_{12} & \dots & AB_{1p} \\ AB_{21} & \color{red}{AB_{22}} & \dots & AB_{2p} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}
 \end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}
 \mathbf{AB} &= \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ \color{red}{A_{21}} & \color{red}{A_{22}} & \dots & \color{red}{A_{2n}} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} B_{11} & B_{12} & \dots & \color{red}{B_{1p}} \\ B_{21} & B_{22} & \dots & \color{red}{B_{2p}} \\ & & \dots & \\ B_{n1} & B_{n2} & \dots & \color{red}{B_{np}} \end{pmatrix} \\
 &= \begin{pmatrix} AB_{11} & AB_{12} & \dots & AB_{1p} \\ AB_{21} & AB_{22} & \dots & \color{red}{AB_{2p}} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}
 \end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}
 \mathbf{AB} &= \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} B_{11} & B_{12} & \dots & B_{1p} \\ B_{21} & B_{22} & \dots & B_{2p} \\ & & \dots & \\ B_{n1} & B_{n2} & \dots & B_{np} \end{pmatrix} \\
 &= \begin{pmatrix} AB_{11} & AB_{12} & \dots & AB_{1p} \\ AB_{21} & AB_{22} & \dots & AB_{2p} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}
 \end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}
 \mathbf{AB} &= \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} B_{11} & B_{12} & \dots & B_{1p} \\ B_{21} & B_{22} & \dots & B_{2p} \\ & & \dots & \\ B_{n1} & B_{n2} & \dots & B_{np} \end{pmatrix} \\
 &= \begin{pmatrix} AB_{11} & AB_{12} & \dots & AB_{1p} \\ AB_{21} & AB_{22} & \dots & AB_{2p} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}
 \end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Matrix Product

Applying two linear operators **A** and **B** defines another linear operator, which can be represented by another matrix **AB**.

$$\begin{aligned}
 \mathbf{AB} &= \begin{pmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ & & \dots & \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{pmatrix} \begin{pmatrix} B_{11} & B_{12} & \dots & B_{1p} \\ B_{21} & B_{22} & \dots & B_{2p} \\ & & \dots & \\ B_{n1} & B_{n2} & \dots & B_{np} \end{pmatrix} \\
 &= \begin{pmatrix} AB_{11} & AB_{12} & \dots & AB_{1p} \\ AB_{21} & AB_{22} & \dots & AB_{2p} \\ & & \dots & \\ AB_{m1} & AB_{m2} & \dots & AB_{mp} \end{pmatrix}
 \end{aligned}$$

Note that if **A** is $m \times n$ and **B** is $n \times p$, then **AB** is $m \times p$.

Transpose of a Matrix

- The transpose of a matrix is obtained by flipping the rows and columns.

$$\mathbf{A}^T = \begin{pmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ & & \dots & \\ A_{1m} & A_{2m} & \dots & A_{nm} \end{pmatrix}$$

Transpose of a Matrix

- The transpose of a matrix is obtained by flipping the rows and columns.

$$\mathbf{A}^T = \begin{pmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ & & \dots & \\ A_{1m} & A_{2m} & \dots & A_{nm} \end{pmatrix}$$

- Some simple properties :

Transpose of a Matrix

- The transpose of a matrix is obtained by flipping the rows and columns.

$$\mathbf{A}^T = \begin{pmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ & & \dots & \\ A_{1m} & A_{2m} & \dots & A_{nm} \end{pmatrix}$$

- Some simple properties :
 - $(\mathbf{A}^T)^T = \mathbf{A}$

Transpose of a Matrix

- The transpose of a matrix is obtained by flipping the rows and columns.

$$\mathbf{A}^T = \begin{pmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ & & \dots & \\ A_{1m} & A_{2m} & \dots & A_{nm} \end{pmatrix}$$

- Some simple properties :
 - $(\mathbf{A}^T)^T = \mathbf{A}$
 - $(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$

Transpose of a Matrix

- The transpose of a matrix is obtained by flipping the rows and columns.

$$\mathbf{A}^T = \begin{pmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ \dots & \dots & \dots & \dots \\ A_{1m} & A_{2m} & \dots & A_{nm} \end{pmatrix}$$

- Some simple properties :
 - $(\mathbf{A}^T)^T = \mathbf{A}$
 - $(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$
 - $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$

Transpose of a Matrix

- The transpose of a matrix is obtained by flipping the rows and columns.

$$\mathbf{A}^T = \begin{pmatrix} A_{11} & A_{21} & \dots & A_{n1} \\ A_{12} & A_{22} & \dots & A_{n2} \\ & & \dots & \\ A_{1m} & A_{2m} & \dots & A_{nm} \end{pmatrix}$$

- Some simple properties :
 - $(\mathbf{A}^T)^T = \mathbf{A}$
 - $(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$
 - $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$
- The inner product for vectors can be represented as $\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u}^T \mathbf{v}$.

Identity Operator

$$I = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ & & \dots & \\ 0 & 0 & \dots & 1 \end{pmatrix}$$

Identity Operator

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ & & \dots & \\ 0 & 0 & \dots & 1 \end{pmatrix}$$

- For any matrix \mathbf{A} , $\mathbf{AI} = \mathbf{A}$.

Identity Operator

$$I = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ & & \dots & \\ 0 & 0 & \dots & 1 \end{pmatrix}$$

- For any matrix \mathbf{A} , $\mathbf{AI} = \mathbf{A}$.
- I is the **identity operator** for the matrix product.

Column Space, Row Space and Rank

- Let \mathbf{A} be an $m \times n$ matrix.

Column Space, Row Space and Rank

- Let \mathbf{A} be an $m \times n$ matrix.
- **Column space :**

Column Space, Row Space and Rank

- Let \mathbf{A} be an $m \times n$ matrix.
- **Column space :**
 - Span of the columns of \mathbf{A} .

Column Space, Row Space and Rank

- Let \mathbf{A} be an $m \times n$ matrix.
- **Column space :**
 - Span of the columns of \mathbf{A} .
 - Linear subspace of \mathbb{R}^m .

Column Space, Row Space and Rank

- Let \mathbf{A} be an $m \times n$ matrix.
- **Column space :**
 - Span of the columns of \mathbf{A} .
 - Linear subspace of \mathbb{R}^m .
- **Row space :**

Column Space, Row Space and Rank

- Let \mathbf{A} be an $m \times n$ matrix.
- **Column space :**
 - Span of the columns of \mathbf{A} .
 - Linear subspace of \mathbb{R}^m .
- **Row space :**
 - Span of the rows of \mathbf{A} .

Column Space, Row Space and Rank

- Let \mathbf{A} be an $m \times n$ matrix.
- **Column space :**
 - Span of the columns of \mathbf{A} .
 - Linear subspace of \mathbb{R}^m .
- **Row space :**
 - Span of the rows of \mathbf{A} .
 - Linear subspace of \mathbb{R}^n .

Column Space, Row Space and Rank

- Let \mathbf{A} be an $m \times n$ matrix.
- **Column space :**
 - Span of the columns of \mathbf{A} .
 - Linear subspace of \mathbb{R}^m .
- **Row space :**
 - Span of the rows of \mathbf{A} .
 - Linear subspace of \mathbb{R}^n .
- **Important fact :**

The column and row spaces of **any** matrix have **the same dimension**.

Column Space, Row Space and Rank

- Let \mathbf{A} be an $m \times n$ matrix.
- **Column space :**
 - Span of the columns of \mathbf{A} .
 - Linear subspace of \mathbb{R}^m .
- **Row space :**
 - Span of the rows of \mathbf{A} .
 - Linear subspace of \mathbb{R}^n .
- **Important fact :**

The column and row spaces of **any** matrix have **the same dimension**.
- This dimension is the **rank** of the matrix.

Range and Null space

- Let \mathbf{A} be an $m \times n$ matrix.

Range and Null space

- Let \mathbf{A} be an $m \times n$ matrix.
- **Range :**

Range and Null space

- Let \mathbf{A} be an $m \times n$ matrix.
- **Range :**
 - Set of vectors equal to $\mathbf{A}\mathbf{u}$ for some $\mathbf{u} \in \mathbb{R}^n$.

Range and Null space

- Let \mathbf{A} be an $m \times n$ matrix.
- **Range :**
 - Set of vectors equal to $\mathbf{A}\mathbf{u}$ for some $\mathbf{u} \in \mathbb{R}^n$.
 - Linear subspace of \mathbb{R}^m .

Range and Null space

- Let \mathbf{A} be an $m \times n$ matrix.
- **Range :**
 - Set of vectors equal to $\mathbf{A}\mathbf{u}$ for some $\mathbf{u} \in \mathbb{R}^n$.
 - Linear subspace of \mathbb{R}^m .
- **Null space :**

Range and Null space

- Let \mathbf{A} be an $m \times n$ matrix.
- **Range :**
 - Set of vectors equal to $\mathbf{A}\mathbf{u}$ for some $\mathbf{u} \in \mathbb{R}^n$.
 - Linear subspace of \mathbb{R}^m .
- **Null space :**
 - \mathbf{u} belongs to the null space if $\mathbf{A}\mathbf{u} = \mathbf{0}$.

Range and Null space

- Let \mathbf{A} be an $m \times n$ matrix.
- **Range :**
 - Set of vectors equal to $\mathbf{A}\mathbf{u}$ for some $\mathbf{u} \in \mathbb{R}^n$.
 - Linear subspace of \mathbb{R}^m .
- **Null space :**
 - \mathbf{u} belongs to the null space if $\mathbf{A}\mathbf{u} = \mathbf{0}$.
 - Subspace of \mathbb{R}^n .

Range and Null space

- Let \mathbf{A} be an $m \times n$ matrix.
- **Range :**
 - Set of vectors equal to $\mathbf{A}\mathbf{u}$ for some $\mathbf{u} \in \mathbb{R}^n$.
 - Linear subspace of \mathbb{R}^m .
- **Null space :**
 - \mathbf{u} belongs to the null space if $\mathbf{A}\mathbf{u} = \mathbf{0}$.
 - Subspace of \mathbb{R}^n .
 - Every vector in the null space is orthogonal to the rows of \mathbf{A} .
The **null space** and **row space** of a matrix are **orthogonal**.

Range and Column Space

Another interpretation of the matrix vector product (\mathbf{A}_i is column i of \mathbf{A}) :

$$\begin{aligned}\mathbf{A}\mathbf{u} &= \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_2 & \cdots & \mathbf{A}_n \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{pmatrix} \\ &= u_1\mathbf{A}_1 + u_2\mathbf{A}_2 + \cdots + u_n\mathbf{A}_n\end{aligned}$$

Range and Column Space

Another interpretation of the matrix vector product (\mathbf{A}_i is column i of \mathbf{A}) :

$$\begin{aligned}\mathbf{A}\mathbf{u} &= \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_2 & \dots & \mathbf{A}_n \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ u_n \end{pmatrix} \\ &= u_1\mathbf{A}_1 + u_2\mathbf{A}_2 + \dots + u_n\mathbf{A}_n\end{aligned}$$

- The result is a **linear combination** of the columns of \mathbf{A} .

Range and Column Space

Another interpretation of the matrix vector product (\mathbf{A}_i is column i of \mathbf{A}) :

$$\begin{aligned}\mathbf{A}\mathbf{u} &= \begin{pmatrix} \mathbf{A}_1 & \mathbf{A}_2 & \dots & \mathbf{A}_n \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ u_n \end{pmatrix} \\ &= u_1\mathbf{A}_1 + u_2\mathbf{A}_2 + \dots + u_n\mathbf{A}_n\end{aligned}$$

- The result is a **linear combination** of the columns of \mathbf{A} .
- For any matrix, the **range** is **equal** to the **column space**.

Matrix Inverse

- For an $n \times n$ matrix \mathbf{A} : **rank + dim(null space) = n.**

Matrix Inverse

- For an $n \times n$ matrix \mathbf{A} : **rank** + **dim(null space)** = **n**.
- If $\dim(\text{null space})=0$ then \mathbf{A} is **full rank**.

Matrix Inverse

- For an $n \times n$ matrix \mathbf{A} : **rank** + **dim(null space)** = **n**.
- If $\dim(\text{null space})=0$ then \mathbf{A} is **full rank**.
- In this case, the action of the matrix is invertible.

Matrix Inverse

- For an $n \times n$ matrix \mathbf{A} : **rank + dim(null space) = n**.
- If $\dim(\text{null space})=0$ then \mathbf{A} is **full rank**.
- In this case, the action of the matrix is invertible.
- The inversion is also linear and consequently can be represented by another matrix \mathbf{A}^{-1} .

Matrix Inverse

- For an $n \times n$ matrix \mathbf{A} : **rank** + **dim(null space)** = **n**.
- If $\dim(\text{null space})=0$ then \mathbf{A} is **full rank**.
- In this case, the action of the matrix is invertible.
- The inversion is also linear and consequently can be represented by another matrix \mathbf{A}^{-1} .
- \mathbf{A}^{-1} is the **only** matrix such that $\mathbf{A}^{-1}\mathbf{A} = \mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$.

Orthogonal Matrices

- An **orthogonal** matrix \mathbf{U} satisfies $\mathbf{U}^T \mathbf{U} = \mathbf{I}$.

Orthogonal Matrices

- An **orthogonal** matrix \mathbf{U} satisfies $\mathbf{U}^T \mathbf{U} = \mathbf{I}$.
- Equivalently, \mathbf{U} has **orthonormal columns**.

Orthogonal Matrices

- An **orthogonal** matrix \mathbf{U} satisfies $\mathbf{U}^T \mathbf{U} = \mathbf{I}$.
- Equivalently, \mathbf{U} has **orthonormal columns**.
- Applying an orthogonal matrix to two vectors **does not change** their inner product :

$$\begin{aligned}\langle \mathbf{U}\mathbf{u}, \mathbf{U}\mathbf{v} \rangle &= (\mathbf{U}\mathbf{u})^T \mathbf{U}\mathbf{v} \\ &= \mathbf{u}^T \mathbf{U}^T \mathbf{U}\mathbf{v} \\ &= \mathbf{u}^T \mathbf{v} \\ &= \langle \mathbf{u}, \mathbf{v} \rangle\end{aligned}$$

Orthogonal Matrices

- An **orthogonal** matrix \mathbf{U} satisfies $\mathbf{U}^T \mathbf{U} = \mathbf{I}$.
- Equivalently, \mathbf{U} has **orthonormal columns**.
- Applying an orthogonal matrix to two vectors **does not change** their inner product :

$$\begin{aligned}\langle \mathbf{U}\mathbf{u}, \mathbf{U}\mathbf{v} \rangle &= (\mathbf{U}\mathbf{u})^T \mathbf{U}\mathbf{v} \\ &= \mathbf{u}^T \mathbf{U}^T \mathbf{U}\mathbf{v} \\ &= \mathbf{u}^T \mathbf{v} \\ &= \langle \mathbf{u}, \mathbf{v} \rangle\end{aligned}$$

- Example : Matrices that represent **rotations** are orthogonal.

Trace and Determinant

- The **trace** is the sum of the diagonal elements of a square matrix.

Trace and Determinant

- The **trace** is the sum of the diagonal elements of a square matrix.
- The **determinant** of a square matrix \mathbf{A} is denoted by $|\mathbf{A}|$.

Trace and Determinant


- The **trace** is the sum of the diagonal elements of a square matrix.
- The **determinant** of a square matrix \mathbf{A} is denoted by $|\mathbf{A}|$.
- For a 2×2 matrix $\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$:

Trace and Determinant

- The **trace** is the sum of the diagonal elements of a square matrix.
- The **determinant** of a square matrix \mathbf{A} is denoted by $|\mathbf{A}|$.
- For a 2×2 matrix $\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$:
 - $|\mathbf{A}| = ad - bc$

Trace and Determinant

- The **trace** is the sum of the diagonal elements of a square matrix.
- The **determinant** of a square matrix \mathbf{A} is denoted by $|\mathbf{A}|$.
- For a 2×2 matrix $\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$:
 - $|\mathbf{A}| = ad - bc$
 - The absolute value of $|\mathbf{A}|$ is the area of the parallelogram given by the rows of \mathbf{A} .



Properties of the Determinant

- The definition can be generalized to larger matrices.

Properties of the Determinant

- The definition can be generalized to larger matrices.
- $|\mathbf{A}| = |\mathbf{A}^T|$.

Properties of the Determinant

- The definition can be generalized to larger matrices.
- $|\mathbf{A}| = |\mathbf{A}^T|$.
- $|\mathbf{AB}| = |\mathbf{A}| |\mathbf{B}|$

Properties of the Determinant

- The definition can be generalized to larger matrices.
- $|\mathbf{A}| = |\mathbf{A}^T|$.
- $|\mathbf{AB}| = |\mathbf{A}| |\mathbf{B}|$
- $|\mathbf{A}| = 0$ if and only if \mathbf{A} is **not invertible**.

Properties of the Determinant

- The definition can be generalized to larger matrices.
- $|\mathbf{A}| = |\mathbf{A}^T|$.
- $|\mathbf{AB}| = |\mathbf{A}| |\mathbf{B}|$
- $|\mathbf{A}| = 0$ if and only if \mathbf{A} is **not invertible**.
- If \mathbf{A} is invertible, then $|\mathbf{A}^{-1}| = \frac{1}{|\mathbf{A}|}$.

- 1 Vectors
- 2 Matrices
- 3 Matrix Decompositions**
- 4 Application : Image Compression

Eigenvalues and Eigenvectors

- An **eigenvalue** λ of a square matrix **A** satisfies :

$$\mathbf{A}\mathbf{u} = \lambda\mathbf{u}$$

for some vector \mathbf{u} , which we call an **eigenvector**.

Eigenvalues and Eigenvectors

- An **eigenvalue** λ of a square matrix **A** satisfies :

$$\mathbf{A}\mathbf{u} = \lambda\mathbf{u}$$

for some vector **u**, which we call an **eigenvector**.

- Geometrically, the operator **A** **expands** ($\lambda > 1$) or **contracts** ($\lambda < 1$) eigenvectors but **does not rotate** them.

Eigenvalues and Eigenvectors

- An **eigenvalue** λ of a square matrix **A** satisfies :

$$\mathbf{A}\mathbf{u} = \lambda\mathbf{u}$$

for some vector **u**, which we call an **eigenvector**.

- Geometrically, the operator **A** **expands** ($\lambda > 1$) or **contracts** ($\lambda < 1$) eigenvectors but **does not rotate** them.
- If **u** is an eigenvector of **A**, it is in the null space of $\mathbf{A} - \lambda\mathbf{I}$, which is consequently **not invertible**.

Eigenvalues and Eigenvectors

- An **eigenvalue** λ of a square matrix **A** satisfies :

$$\mathbf{A}\mathbf{u} = \lambda\mathbf{u}$$

for some vector \mathbf{u} , which we call an **eigenvector**.

- Geometrically, the operator **A** **expands** ($\lambda > 1$) or **contracts** ($\lambda < 1$) eigenvectors but **does not rotate** them.
- If \mathbf{u} is an eigenvector of **A**, it is in the null space of $\mathbf{A} - \lambda\mathbf{I}$, which is consequently **not invertible**.
- The eigenvalues of **A** are the roots of the equation $|\mathbf{A} - \lambda\mathbf{I}| = 0$.

Eigenvalues and Eigenvectors

- An **eigenvalue** λ of a square matrix **A** satisfies :

$$\mathbf{A}\mathbf{u} = \lambda\mathbf{u}$$

for some vector **u**, which we call an **eigenvector**.

- Geometrically, the operator **A** **expands** ($\lambda > 1$) or **contracts** ($\lambda < 1$) eigenvectors but **does not rotate** them.
- If **u** is an eigenvector of **A**, it is in the null space of $\mathbf{A} - \lambda\mathbf{I}$, which is consequently **not invertible**.
- The eigenvalues of **A** are the roots of the equation $|\mathbf{A} - \lambda\mathbf{I}| = 0$.
- **Not** the way eigenvalues are calculated in practice.

Eigenvalues and Eigenvectors

- An **eigenvalue** λ of a square matrix \mathbf{A} satisfies :

$$\mathbf{A}\mathbf{u} = \lambda\mathbf{u}$$

for some vector \mathbf{u} , which we call an **eigenvector**.

- Geometrically, the operator \mathbf{A} **expands** ($\lambda > 1$) or **contracts** ($\lambda < 1$) eigenvectors but **does not rotate** them.
- If \mathbf{u} is an eigenvector of \mathbf{A} , it is in the null space of $\mathbf{A} - \lambda\mathbf{I}$, which is consequently **not invertible**.
- The eigenvalues of \mathbf{A} are the roots of the equation $|\mathbf{A} - \lambda\mathbf{I}| = 0$.
- **Not** the way eigenvalues are calculated in practice.
- Eigenvalues and eigenvectors can be **complex valued**, even if all the entries of \mathbf{A} are real.

Eigendecomposition of a Matrix

- Let \mathbf{A} be an $n \times n$ square matrix with n linearly independent eigenvectors $\mathbf{p}_1 \dots \mathbf{p}_n$ and eigenvalues $\lambda_1 \dots \lambda_n$.

Eigendecomposition of a Matrix

- Let \mathbf{A} be an $n \times n$ square matrix with n linearly independent eigenvectors $\mathbf{p}_1 \dots \mathbf{p}_n$ and eigenvalues $\lambda_1 \dots \lambda_n$.
- We define the matrices :

$$\mathbf{P} = \begin{pmatrix} \mathbf{p}_1 & \dots & \mathbf{p}_n \end{pmatrix}$$
$$\mathbf{\Lambda} = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ & & \dots & \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$$

Eigendecomposition of a Matrix

- Let \mathbf{A} be an $n \times n$ square matrix with n linearly independent eigenvectors $\mathbf{p}_1 \dots \mathbf{p}_n$ and eigenvalues $\lambda_1 \dots \lambda_n$.
- We define the matrices :

$$\mathbf{P} = \begin{pmatrix} \mathbf{p}_1 & \dots & \mathbf{p}_n \end{pmatrix}$$
$$\mathbf{\Lambda} = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ & & \dots & \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$$

- \mathbf{A} satisfies $\mathbf{AP} = \mathbf{P}\mathbf{\Lambda}$.

Eigendecomposition of a Matrix

- Let \mathbf{A} be an $n \times n$ square matrix with n linearly independent eigenvectors $\mathbf{p}_1 \dots \mathbf{p}_n$ and eigenvalues $\lambda_1 \dots \lambda_n$.
- We define the matrices :

$$\mathbf{P} = \begin{pmatrix} \mathbf{p}_1 & \dots & \mathbf{p}_n \end{pmatrix}$$
$$\mathbf{\Lambda} = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ & & \dots & \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$$

- \mathbf{A} satisfies $\mathbf{AP} = \mathbf{P}\mathbf{\Lambda}$.
- \mathbf{P} is full rank. Inverting it yields the **eigendecomposition** of \mathbf{A} :

$$\mathbf{A} = \mathbf{P}\mathbf{\Lambda}\mathbf{P}^{-1}$$

Properties of the Eigendecomposition

- **Not all** matrices are **diagonalizable** (have an eigendecomposition).
Example : $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Properties of the Eigendecomposition

- **Not all** matrices are **diagonalizable** (have an eigendecomposition).
Example : $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.
- $\text{Trace}(\mathbf{A}) = \text{Trace}(\mathbf{\Lambda}) = \sum_{i=1}^n \lambda_i$.

Properties of the Eigendecomposition

- **Not all** matrices are **diagonalizable** (have an eigendecomposition).
Example : $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.
- $\text{Trace}(\mathbf{A}) = \text{Trace}(\mathbf{\Lambda}) = \sum_{i=1}^n \lambda_i$.
- $|\mathbf{A}| = |\mathbf{\Lambda}| = \prod_{i=1}^n \lambda_i$.

Properties of the Eigendecomposition

- **Not all** matrices are **diagonalizable** (have an eigendecomposition).
Example : $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.
- $\text{Trace}(\mathbf{A}) = \text{Trace}(\mathbf{\Lambda}) = \sum_{i=1}^n \lambda_i$.
- $|\mathbf{A}| = |\mathbf{\Lambda}| = \prod_{i=1}^n \lambda_i$.
- The **rank** of \mathbf{A} is equal to the **number of nonzero eigenvalues**.


Properties of the Eigendecomposition

- **Not all** matrices are **diagonalizable** (have an eigendecomposition).
Example : $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.
- $\text{Trace}(\mathbf{A}) = \text{Trace}(\mathbf{\Lambda}) = \sum_{i=1}^n \lambda_i$.
- $|\mathbf{A}| = |\mathbf{\Lambda}| = \prod_{i=1}^n \lambda_i$.
- The **rank** of \mathbf{A} is equal to the **number of nonzero eigenvalues**.
- If λ is a nonzero eigenvalue of \mathbf{A} , $\frac{1}{\lambda}$ is an eigenvalue of \mathbf{A}^{-1} with the **same eigenvector**.

Properties of the Eigendecomposition

- **Not all** matrices are **diagonalizable** (have an eigendecomposition).
Example : $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.
- $\text{Trace}(\mathbf{A}) = \text{Trace}(\mathbf{\Lambda}) = \sum_{i=1}^n \lambda_i$.
- $|\mathbf{A}| = |\mathbf{\Lambda}| = \prod_{i=1}^n \lambda_i$.
- The **rank** of \mathbf{A} is equal to the **number of nonzero eigenvalues**.
- If λ is a nonzero eigenvalue of \mathbf{A} , $\frac{1}{\lambda}$ is an eigenvalue of \mathbf{A}^{-1} with the **same eigenvector**.
- The eigendecomposition makes it possible to compute **matrix powers** efficiently :
$$\mathbf{A}^m = (\mathbf{P}\mathbf{\Lambda}\mathbf{P}^{-1})^m = \mathbf{P}\mathbf{\Lambda}\mathbf{P}^{-1} \mathbf{P}\mathbf{\Lambda}\mathbf{P}^{-1} \dots \mathbf{P}\mathbf{\Lambda}\mathbf{P}^{-1} = \mathbf{P}\mathbf{\Lambda}^m\mathbf{P}^{-1}.$$

Matlab Example



Eigendecomposition of a Symmetric Matrix

- If $\mathbf{A} = \mathbf{A}^T$ then \mathbf{A} is **symmetric**.

Eigendecomposition of a Symmetric Matrix

- If $\mathbf{A} = \mathbf{A}^T$ then \mathbf{A} is **symmetric**.
- The eigenvalues of symmetric matrices are **real**.

Eigendecomposition of a Symmetric Matrix

- If $\mathbf{A} = \mathbf{A}^T$ then \mathbf{A} is **symmetric**.
- The eigenvalues of symmetric matrices are **real**.
- The eigenvectors of symmetric matrices are **orthonormal**.

Eigendecomposition of a Symmetric Matrix

- If $\mathbf{A} = \mathbf{A}^T$ then \mathbf{A} is **symmetric**.
- The eigenvalues of symmetric matrices are **real**.
- The eigenvectors of symmetric matrices are **orthonormal**.
- Consequently, the eigendecomposition becomes :
 $\mathbf{A} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$ for $\mathbf{\Lambda}$ real and \mathbf{U} orthogonal.

Eigendecomposition of a Symmetric Matrix

- If $\mathbf{A} = \mathbf{A}^T$ then \mathbf{A} is **symmetric**.
- The eigenvalues of symmetric matrices are **real**.
- The eigenvectors of symmetric matrices are **orthonormal**.
- Consequently, the eigendecomposition becomes :
 $\mathbf{A} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$ for $\mathbf{\Lambda}$ real and \mathbf{U} orthogonal.
- The eigenvectors of \mathbf{A} are an orthonormal **basis** for the **column space** (and the row space, since they are equal).

Eigendecomposition of a Symmetric Matrix

- The action of a symmetric matrix on a vector $\mathbf{A}\mathbf{u} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{u}$ can be decomposed into :

Eigendecomposition of a Symmetric Matrix

- The action of a symmetric matrix on a vector $\mathbf{A}\mathbf{u} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{u}$ can be decomposed into :
 - 1 Projection of \mathbf{u} onto the column space of \mathbf{A} (multiplication by \mathbf{U}^T).

Eigendecomposition of a Symmetric Matrix

- The action of a symmetric matrix on a vector $\mathbf{A}\mathbf{u} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{u}$ can be decomposed into :
 - 1 Projection of \mathbf{u} onto the column space of \mathbf{A} (multiplication by \mathbf{U}^T).
 - 2 Scaling of each coefficient $\langle \mathbf{U}_i, \mathbf{u} \rangle$ by the corresponding eigenvalue (multiplication by $\mathbf{\Lambda}$).

Eigendecomposition of a Symmetric Matrix

- The action of a symmetric matrix on a vector $\mathbf{A}\mathbf{u} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{u}$ can be decomposed into :
 - 1 Projection of \mathbf{u} onto the column space of \mathbf{A} (multiplication by \mathbf{U}^T).
 - 2 Scaling of each coefficient $\langle \mathbf{U}_i, \mathbf{u} \rangle$ by the corresponding eigenvalue (multiplication by $\mathbf{\Lambda}$).
 - 3 Linear combination of the eigenvectors scaled by the resulting coefficient (multiplication by \mathbf{U}).

Eigendecomposition of a Symmetric Matrix

- The action of a symmetric matrix on a vector $\mathbf{A}\mathbf{u} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{u}$ can be decomposed into :
 - 1 Projection of \mathbf{u} onto the column space of \mathbf{A} (multiplication by \mathbf{U}^T).
 - 2 Scaling of each coefficient $\langle \mathbf{U}_i, \mathbf{u} \rangle$ by the corresponding eigenvalue (multiplication by $\mathbf{\Lambda}$).
 - 3 Linear combination of the eigenvectors scaled by the resulting coefficient (multiplication by \mathbf{U}).
- Summarizing :

$$\mathbf{A}\mathbf{u} = \sum_{i=1}^n \lambda_i \langle \mathbf{U}_i, \mathbf{u} \rangle \mathbf{U}_i$$

Eigendecomposition of a Symmetric Matrix

- The action of a symmetric matrix on a vector $\mathbf{A}\mathbf{u} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T\mathbf{u}$ can be decomposed into :
 - 1 Projection of \mathbf{u} onto the column space of \mathbf{A} (multiplication by \mathbf{U}^T).
 - 2 Scaling of each coefficient $\langle \mathbf{U}_i, \mathbf{u} \rangle$ by the corresponding eigenvalue (multiplication by $\mathbf{\Lambda}$).
 - 3 Linear combination of the eigenvectors scaled by the resulting coefficient (multiplication by \mathbf{U}).
- Summarizing :

$$\mathbf{A}\mathbf{u} = \sum_{i=1}^n \lambda_i \langle \mathbf{U}_i, \mathbf{u} \rangle \mathbf{U}_i$$

- It would be great to generalize this to all matrices...

Singular Value Decomposition

- **Every** matrix has a **singular value decomposition** :

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$$

Singular Value Decomposition

- **Every** matrix has a **singular value decomposition** :

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$$

- The columns of **U** are an orthonormal basis for the **column space** of **A**.

Singular Value Decomposition

- **Every** matrix has a **singular value decomposition** :

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$$

- The columns of **U** are an orthonormal basis for the **column space** of **A**.
- The columns of **V** are an orthonormal basis for the **row space** of **A**.

Singular Value Decomposition

- **Every** matrix has a **singular value decomposition** :

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$$

- The columns of \mathbf{U} are an orthonormal basis for the **column space** of \mathbf{A} .
- The columns of \mathbf{V} are an orthonormal basis for the **row space** of \mathbf{A} .
- $\mathbf{\Sigma}$ is diagonal. The entries on its diagonal $\sigma_i = \Sigma_{ii}$ are positive real numbers, called the **singular values** of \mathbf{A} .

Singular Value Decomposition

- **Every** matrix has a **singular value decomposition** :

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$$

- The columns of \mathbf{U} are an orthonormal basis for the **column space** of \mathbf{A} .
- The columns of \mathbf{V} are an orthonormal basis for the **row space** of \mathbf{A} .
- $\mathbf{\Sigma}$ is diagonal. The entries on its diagonal $\sigma_i = \Sigma_{ii}$ are positive real numbers, called the **singular values** of \mathbf{A} .
- The action of \mathbf{A} on a vector \mathbf{u} can be decomposed into :

$$\mathbf{A}\mathbf{u} = \sum_{i=1}^n \sigma_i \langle \mathbf{V}_i, \mathbf{u} \rangle \mathbf{U}_i$$

Properties of the Singular Value Decomposition

- The eigenvalues of the symmetric matrix $\mathbf{A}^T \mathbf{A}$ are equal to the **square** of the singular values of \mathbf{A} :

$$\mathbf{A}^T \mathbf{A} = \mathbf{V} \Sigma \mathbf{U}^T \mathbf{U} \Sigma \mathbf{V}^T = \mathbf{V} \Sigma^2 \mathbf{V}^T$$

Properties of the Singular Value Decomposition

- The eigenvalues of the symmetric matrix $\mathbf{A}^T \mathbf{A}$ are equal to the **square** of the singular values of \mathbf{A} :

$$\mathbf{A}^T \mathbf{A} = \mathbf{V} \Sigma \mathbf{U}^T \mathbf{U} \Sigma \mathbf{V}^T = \mathbf{V} \Sigma^2 \mathbf{V}^T$$

- The **rank** of a matrix is equal to the number of nonzero singular values.

Properties of the Singular Value Decomposition

- The eigenvalues of the symmetric matrix $\mathbf{A}^T \mathbf{A}$ are equal to the **square** of the singular values of \mathbf{A} :

$$\mathbf{A}^T \mathbf{A} = \mathbf{V} \Sigma \mathbf{U}^T \mathbf{U} \Sigma \mathbf{V}^T = \mathbf{V} \Sigma^2 \mathbf{V}^T$$

- The **rank** of a matrix is equal to the number of nonzero singular values.
- The largest singular value σ_1 is the solution to the optimization problem :

$$\sigma_1 = \max_{\mathbf{x} \neq 0} \frac{\|\mathbf{A}\mathbf{x}\|_2}{\|\mathbf{x}\|_2}$$

Properties of the Singular Value Decomposition

- The eigenvalues of the symmetric matrix $\mathbf{A}^T \mathbf{A}$ are equal to the **square** of the singular values of \mathbf{A} :

$$\mathbf{A}^T \mathbf{A} = \mathbf{V} \Sigma \mathbf{U}^T \mathbf{U} \Sigma \mathbf{V}^T = \mathbf{V} \Sigma^2 \mathbf{V}^T$$

- The **rank** of a matrix is equal to the number of nonzero singular values.
- The largest singular value σ_1 is the solution to the optimization problem :

$$\sigma_1 = \max_{\mathbf{x} \neq 0} \frac{\|\mathbf{Ax}\|_2}{\|\mathbf{x}\|_2}$$

- It can be verified that the largest singular value satisfies the properties of a norm, it is called the **spectral norm** of the matrix.

Properties of the Singular Value Decomposition

- The eigenvalues of the symmetric matrix $\mathbf{A}^T \mathbf{A}$ are equal to the **square** of the singular values of \mathbf{A} :

$$\mathbf{A}^T \mathbf{A} = \mathbf{V} \Sigma \mathbf{U}^T \mathbf{U} \Sigma \mathbf{V}^T = \mathbf{V} \Sigma^2 \mathbf{V}^T$$

- The **rank** of a matrix is equal to the number of nonzero singular values.
- The largest singular value σ_1 is the solution to the optimization problem :

$$\sigma_1 = \max_{\mathbf{x} \neq 0} \frac{\|\mathbf{A}\mathbf{x}\|_2}{\|\mathbf{x}\|_2}$$

- It can be verified that the largest singular value satisfies the properties of a norm, it is called the **spectral norm** of the matrix.
- In statistics analyzing data with the singular value decomposition is called **Principal Component Analysis**.

- 1 Vectors
- 2 Matrices
- 3 Matrix Decompositions
- 4 Application : Image Compression**

Compression using the Singular Value Decomposition

- The singular value decomposition can be viewed as a sum of rank 1 matrices :

$$\mathbf{A} = \begin{array}{c} \sigma_1 \\ \boxed{} \\ \mathbf{u}_1 \end{array} \overline{} \mathbf{v}_1^T + \sigma_2 \begin{array}{c} \boxed{} \\ \mathbf{u}_2 \end{array} \overline{} \mathbf{v}_2^T \dots + \sigma_R \begin{array}{c} \boxed{} \\ \mathbf{u}_R \end{array} \overline{} \mathbf{v}_R^T$$

Compression using the Singular Value Decomposition

- The singular value decomposition can be viewed as a sum of rank 1 matrices :

$$\mathbf{A} = \begin{array}{c} \sigma_1 \\ \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right. \end{array} \begin{array}{c} \mathbf{v}_1^T \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \sigma_2 \begin{array}{c} \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right. \end{array} \begin{array}{c} \mathbf{v}_2^T \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \dots + \sigma_R \begin{array}{c} \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right. \end{array} \begin{array}{c} \mathbf{v}_R^T \\ \text{---} \\ \text{---} \\ \text{---} \end{array}$$

\mathbf{u}_1 \mathbf{u}_2 \mathbf{u}_R

- If some of the singular values are very small, we can **discard** them.

Compression using the Singular Value Decomposition

- The singular value decomposition can be viewed as a sum of rank 1 matrices :

$$\mathbf{A} = \begin{array}{c} \sigma_1 \\ \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right. \end{array} \begin{array}{c} \mathbf{v}_1^T \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \sigma_2 \begin{array}{c} \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right. \end{array} \begin{array}{c} \mathbf{v}_2^T \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \dots + \sigma_R \begin{array}{c} \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right. \end{array} \begin{array}{c} \mathbf{v}_R^T \\ \text{---} \\ \text{---} \\ \text{---} \end{array}$$

\mathbf{u}_1 \mathbf{u}_2 \mathbf{u}_R

- If some of the singular values are very small, we can **discard** them.
- This is a form of **lossy compression**.

Compression using the Singular Value Decomposition

- Truncating the singular value decomposition allows us to represent the matrix with **less parameters**.

$$\mathbf{A}_{\text{TRUNC}} = \mathbf{U}_{\text{TRUNC}} \mathbf{\Sigma}_{\text{TRUNC}} \mathbf{V}_{\text{TRUNC}}^T$$

Compression using the Singular Value Decomposition

- Truncating the singular value decomposition allows us to represent the matrix with **less parameters**.

$$\boxed{\mathbf{A}_{\text{TRUNC}}} = \boxed{\mathbf{U}_{\text{TRUNC}}} \boxed{\mathbf{\Sigma}_{\text{TRUNC}}} \boxed{\mathbf{V}_{\text{TRUNC}}^T}$$

- For a 512×512 matrix :

Compression using the Singular Value Decomposition

- Truncating the singular value decomposition allows us to represent the matrix with **less parameters**.

$$\boxed{\mathbf{A}_{\text{TRUNC}}} = \boxed{\mathbf{U}_{\text{TRUNC}}} \boxed{\mathbf{\Sigma}_{\text{TRUNC}}} \boxed{\mathbf{V}_{\text{TRUNC}}^T}$$

- For a 512×512 matrix :
 - Full representation : $512 \times 512 = \mathbf{262\ 144}$

Compression using the Singular Value Decomposition

- Truncating the singular value decomposition allows us to represent the matrix with **less parameters**.

$$\boxed{\mathbf{A}_{\text{TRUNC}}} = \begin{bmatrix} \mathbf{U}_{\text{TRUNC}} \end{bmatrix} \begin{bmatrix} \mathbf{\Sigma}_{\text{TRUNC}} \end{bmatrix} \begin{bmatrix} \mathbf{V}_{\text{TRUNC}}^T \end{bmatrix}$$

- For a 512×512 matrix :
 - Full representation : $512 \times 512 = \mathbf{262\ 144}$
 - Rank 10 approximation : $512 \times 10 + 10 + 512 \times 10 = \mathbf{10\ 250}$

Compression using the Singular Value Decomposition

- Truncating the singular value decomposition allows us to represent the matrix with **less parameters**.

$$\mathbf{A}_{\text{TRUNC}} = \mathbf{U}_{\text{TRUNC}} \mathbf{\Sigma}_{\text{TRUNC}} \mathbf{V}_{\text{TRUNC}}^T$$

- For a 512×512 matrix :
 - Full representation : $512 \times 512 = \mathbf{262\ 144}$
 - Rank 10 approximation : $512 \times 10 + 10 + 512 \times 10 = \mathbf{10\ 250}$
 - Rank 40 approximation : $512 \times 40 + 40 + 512 \times 40 = \mathbf{41\ 000}$

Compression using the Singular Value Decomposition

- Truncating the singular value decomposition allows us to represent the matrix with **less parameters**.

$$\mathbf{A}_{\text{TRUNC}} = \mathbf{U}_{\text{TRUNC}} \mathbf{\Sigma}_{\text{TRUNC}} \mathbf{V}_{\text{TRUNC}}^T$$

- For a 512×512 matrix :
 - Full representation : $512 \times 512 = \mathbf{262\ 144}$
 - Rank 10 approximation : $512 \times 10 + 10 + 512 \times 10 = \mathbf{10\ 250}$
 - Rank 40 approximation : $512 \times 40 + 40 + 512 \times 40 = \mathbf{41\ 000}$
 - Rank 80 approximation : $512 \times 80 + 80 + 512 \times 80 = \mathbf{82\ 000}$