

High-Throughput Sequencing Technologies

Jason A. Reuter,¹ Damek V. Spacek,¹ and Michael P. Snyder^{1,*}

¹Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305, USA

*Correspondence: mpsnyder@stanford.edu

<http://dx.doi.org/10.1016/j.molcel.2015.05.004>

The human genome sequence has profoundly altered our understanding of biology, human diversity, and disease. The path from the first draft sequence to our nascent era of personal genomes and genomic medicine has been made possible only because of the extraordinary advancements in DNA sequencing technologies over the past 10 years. Here, we discuss commonly used high-throughput sequencing platforms, the growing array of sequencing assays developed around them, as well as the challenges facing current sequencing platforms and their clinical application.

Introduction

The human genome sequence was completed in draft form in 2001 (Lander et al., 2001; Venter et al., 2001). Shortly thereafter, the genome sequences of several model organisms were determined (Chinwalla et al., 2002; Gibbs et al., 2004; Chimpanzee Sequencing and Analysis Consortium, 2005). These feats were accomplished with Sanger DNA sequencing, which was limited in throughput and high cost; indeed, the first human genome sequence was estimated to cost 0.5–1 billion dollars. These limitations reduced the potential of DNA sequencing for other applications, such as personal genome sequencing. Following the release of the “finished” human genome (International Human Genome Sequencing Consortium, 2004), the National Human Genome Research Institute (NHGRI) created a 70 million dollar DNA sequencing technology initiative aimed at achieving a \$1,000 human genome in 10 years (Schloss, 2008), and a flurry of high-throughput sequencing (HTS) technologies emerged.

To put this initiative in perspective, improvements to traditional Sanger sequencing had decreased the per base cost by around 100-fold by the completion of the Human Genome Project (Schloss, 2008). To reach the \$1,000 dollar genome threshold, however, an additional leap of five orders of magnitude was necessary. Much of this divide has been traversed—the cost of a genome sequence (without interpretation) is presently less than \$2,000. The road to this milestone involved many commercial HTS platforms, which differ in their details but typically follow a similar general paradigm: template preparation, clonal amplification, followed by cyclical rounds of massively parallel sequencing. The specific strategy employed by each platform determines the quality, quantity, and biases of the resulting sequence data and the platform’s usefulness for particular applications.

Several excellent reviews have covered HTS platform strategies in great depth (Metzker, 2010; Morey et al., 2013). Many important platforms are not covered here, including Roche/454’s pyrophosphate Genome Sequencer (Margulies et al., 2005) and Helicos’ single-molecule Heliscope sequencer (Harris et al., 2008) as well as the Polonator (Shendure et al., 2005), ABI’s SOLiD (Valouev et al., 2008), and Complete Genomics’ DNA nano-array sequencer (Drmanac et al., 2010). Instead, we focus on the most commonly used platforms today as well as more recent developments. We also provide an overview of

the growing array of HTS applications and highlight their use by the genomics community to illuminate previously intractable topics in biology. Finally, we discuss the limitations of current platforms and challenges to clinical sequencing.

Overview of Selected Commercially Available HTS Platforms

Illumina

Illumina/Solexa released the Genome Analyzer II in 2006, and advances in Illumina’s technology over the intervening years have largely set the pace for the tremendous gains in output and reductions in cost (Figure 1). As a consequence, Illumina machines currently dominate the HTS market. The sequencing process involves clonal amplification of adaptor-ligated DNA fragments on the surface of a glass slide (Bentley et al., 2008) (Figure 2A). Bases are read using a cyclic reversible termination strategy, which sequences the template strand one nucleotide at a time through progressive rounds of base incorporation, washing, imaging, and cleavage. In this strategy, fluorescently labeled 3′-O-azidomethyl-dNTPs are used to pause the polymerization reaction, enabling removal of unincorporated bases and fluorescent imaging to determine the added nucleotide (Guo et al., 2008). Following scanning of the flow cell with a coupled-charge device (CCD) camera, the fluorescent moiety and the 3′ block are removed, and the process is repeated. Across all Illumina models the overall error rates are below 1%, and the most common type of error is substitution (Dohm et al., 2008).

Illumina currently produces a suite of sequencers (MiSeq, NextSeq 500, and the HiSeq series) optimized for a variety of throughputs and turnaround times. The MiSeq and HiSeqs are the most established platforms. The MiSeq is designed as a fast, personal benchtop sequencer, with run times as low as 4 hr and outputs intended for targeted sequencing and sequencing of small genomes. The HiSeq 2500, on the other hand, is engineered for high-throughput applications, yielding current outputs of 1 Tb in 6 days. Unlike previous HiSeq models, the HiSeq 2500 can also be run in rapid mode, which is less cost effective but can produce a 30× human genome in 27 hr.

In early 2014, Illumina introduced the NextSeq 500 as well as the HiSeq X Ten. Similar to the MiSeq, the NextSeq 500 is designed as a fast benchtop sequencer for individual labs.




Figure 1. Timeline and Comparison of Commercial HTS Instruments

Plot of commercial release dates versus machine outputs per run are shown. For the MinION, outputs from an 18 hr run were used (Ashton et al., 2015). Numbers inside data points denote current read lengths. Sequencing platforms are color coded.

(Figure 2B). Unlike pyrosequencing, which couples base incorporation with luciferase-based light production, Ion Torrent's semiconductor sequencing measures pH changes induced by the release of hydrogen ions during DNA extension (Rothberg et al., 2011). These pH changes are detected by a sensor positioned at the bottom of the microwell and converted into a voltage signal. The voltage signal is proportional to the number of bases incorporated, and the sequential addition of individual nucleotides during each sequencing cycle allows base discrimination.

Moreover, Ion Torrent avoids optical scanning to distinguish nucleotides during cycles of sequencing, a difference that dramatically speeds sequencing runs and reduces costs.

Ion Torrent released a second machine in 2012, the Ion Proton, which increases output over the PGM by an order of magnitude (1 Gb versus 10 Gb). However, the Proton currently features a maximum of 200 bp read lengths as opposed to 400 bp for the PGM. Multiple chips are also available to tailor outputs for different applications. The PGM is most useful for targeted resequencing projects and small genome analysis, whereas the Proton is capable of exome sequencing and whole-transcriptome analysis. The speed of sequencing, 2–8 hr depending on the machine and chip used, make these sequencers particularly useful for clinical applications (Mellmann et al., 2011). Insertions and deletions (indels) are the most common error types (Liu et al., 2012). Because the correlation between the number of bases incorporated and the subsequent voltage change does not perfectly scale, homopolymer repeats longer than 6 bp lead to increased error rates (Rothberg et al., 2011).

Pacific Biosciences

Single-molecule real-time (SMRT) sequencing was pioneered by Nanofluidics, Inc. and commercialized by Pacific Biosciences. Template preparation involves ligation of single-stranded, hairpin adapters onto the ends of digested DNA or cDNA molecules, generating a capped template (SMRT-bell). By using a strand displacing polymerase, the original DNA molecule can be sequenced multiple times, thereby increasing accuracy (Travers et al., 2010). Importantly, clonal amplification is avoided, allowing direct sequencing of native, and potentially modified, DNA. DNA synthesis occurs in zeptoliter-sized chambers, called zero-mode waveguides (ZMWs), in which a single polymerase is immobilized at the bottom of the chamber (Levene et al., 2003) (Figure 3A). The physics of these chambers reduces background

However, the NextSeq is capable producing 120 Gb, or a single 30× genome, in less than 30 hr. The NextSeq 500 system also employs a novel two-channel sequencing strategy. In this approach, cytosine is labeled red, thymine is labeled green, adenine is effectively yellow (labeled with a mixture of red and green), and guanine is unlabeled. In contrast to the four-channel strategy used in the MiSeq and HiSeq platforms, two-channel sequencing requires only two images for nucleotide detection, reducing data processing times and increasing throughput. Despite the reduced complexity, the overall error rates (<1%) are similar to the more established HiSeq machines.

The HiSeq X Ten is a population-scale whole-genome sequencing (WGS) system that was also released in 2014. It is capable of outputting 1.8 Tb in 3 days or 18,000 genomes at 30× coverage per year. Currently, Illumina only supports WGS of human samples on HiSeq X Ten systems. In addition to enhanced optics and computing capacity, the HiSeq X Ten dramatically increases throughput by incorporating a new patterned flow cell technology that improves cluster generation chemistry. Patterned flow cells contain billions of nanowells that standardize cluster spacing and size, allowing higher cluster densities. Patterned flow cell technology is also used in the recently released HiSeq 3000/4000 machines, which provide outputs and run times in between the HiSeq X Ten and the HiSeq 2500.

Life Technologies/ThermoFisher/Ion Torrent

Life Technologies commercialized Ion Torrent's semiconductor sequencing technology in 2010 in the form of the benchtop Ion PGM sequencer. The template preparation and sequencing steps are conceptually similar to the Roche/454 pyrosequencing platform (Margulies et al., 2005). Namely, emulsion-PCR is used to clonally amplify adaptor-ligated DNA fragments on the surface of beads. The beads are subsequently distributed into microwells where a sequencing-by-synthesis reaction occurs




Figure 2. Clonal Amplification-Based Sequencing Platforms

(A) Illumina's four-color reversible termination sequencing method. DNA templates are first clonally amplified on the surface of a glass flow cell. Sequencing occurs via successive rounds of base incorporation, washing, and imaging. A cleavage step after image acquisition removes the fluorescent dye and regenerates the 3' OH for the next cycle. Analysis of four-color images is used to determine base composition.

(B) Ion Torrent's semiconductor sequencing method. Emulsion-PCR is used to clonally amplify DNA templates on the surface of beads, which are subsequently placed into microwells. pH changes induced by the release of hydrogen ions during DNA extension are detected by a sensor positioned at the bottom of the microwell. These pH changes are converted into a voltage signal, which is proportional to the number of nucleotides added by the polymerase.

In addition to providing long, unbiased reads, another distinguishing characteristic of SMRT sequencing is that the polymerization reaction is monitored in real-time, allowing data pertaining to both base composition and enzymatic kinetics to be collected. Distinct kinetic profiles

are produced as the polymerase encounters various types of DNA methylation (Flusberg et al., 2010). These kinetic signatures have been utilized to map sites of potential 6-methyladenine and 5-methylcytosine genome wide in bacteria (Fang et al., 2012). It is possible that these approaches will be extended to map other types of DNA modifications, including DNA damage induced in cancer cells. Moreover, SMRT sequencing instruments are not limited to studying DNA alone, as other molecules, such as ribosomes, can be tethered to the bottom of the ZMW and monitored at single-molecule resolution (Uemura et al., 2010).

Oxford Nanopore Technologies

Nanopore-based sequencing is an emerging single-molecule strategy that has made significant progress in recent years, with Oxford Nanopore Technologies leading the development and commercialization of this method. Nanopore sequencing can take a variety of forms, but it principally relies on the transition of DNA or individual nucleotides through a small channel (Wang et al., 2015). In Oxford Nanopore's current technology, a sequencing flow cell comprises hundreds of independent micro-wells, each containing a synthetic bilayer perforated by biologic nanopores. Sequencing is accomplished by measuring characteristic changes in current that are induced as the bases are threaded through the pore by a molecular motor protein (Figure 3B). Library preparation is minimal, involving fragmentation of DNA and ligation of adapters. Much like SMRT sequencing, this library preparation methodology can be done with or without PCR amplification. The first adaptor is bound with a propriety motor enzyme as well as a molecular tether, whereas the second adaptor is a hairpin oligonucleotide that is

noise such that phosphate-labeled versions of all four nucleotides can be present simultaneously. Thus, polymerization occurs continuously, and the DNA sequence can be read in real-time from the fluorescent signals recorded in a video (Eid et al., 2009).

Released in 2010, the RS II remains Pacific Biosciences only commercially available machine. However, altering the chemistry and doubling the number of ZMWs to 150 k per SMRT cell have greatly enhanced performance. Using the latest chemistry, each SMRT cell produces ~50 k reads and up to 1 Gb of data in 4 hr. The average read lengths are >14 kb, but individual reads can be as long 60 kb. As with most single-molecule sequencing platforms, high error rates (~11%) are evident for single pass reads, and these errors are dominated by indels. Sequencing errors, however, are distributed randomly, allowing accurate consensus calls with increasing coverage or multiple passes around the same template, so-called circular consensus sequences (Carneiro et al., 2012; Koren et al., 2012). By avoiding clonal amplification, SMRT sequencing is also much less sensitive to GC sequence content than other platforms (Loomis et al., 2013). This suite of characteristics makes SMRT sequencing particularly useful for projects involving de novo assembly of small bacterial and viral genomes as well as large genome finishing (English et al., 2012). Reconstructing structural variation (SV) in the genome (Chaisson et al., 2015) and isoform usage in the transcriptome (Sharon et al., 2013) are also key areas where SMRT sequencing has clear advantages over short read technologies. However, lower throughput and higher per base sequencing costs currently limit the scope of most genome-wide studies.




Figure 3. Single Molecule Sequencing Platforms

(A) Pacific Bioscience's SMRT sequencing. A single polymerase is positioned at the bottom of a ZMW. Phosphate-labeled versions of all four nucleotides are present, allowing continuous polymerization of a DNA template. Base incorporation increases the residence time of the nucleotide in the ZMW, resulting in a detectable fluorescent signal that is captured in a video.

(B) Oxford Nanopore's sequencing strategy. DNA templates are ligated with two adapters. The first adaptor is bound with a motor enzyme as well as a tether, whereas the second adaptor is a hairpin oligo that is bound by the HP motor protein. Changes in current that are induced as the nucleotides pass through the pore are used to discriminate bases. The library design allows sequencing of both strands of DNA from a single molecule (two-direction reads).

bound by a second so-called HP motor protein (Quick et al., 2014). This library design allows sequencing of both strands of DNA from a single molecule, which increases accuracy (Ashton et al., 2015; Quick et al., 2014).

The first commercially available device for nanopore sequencing is the MinION, a USB-powered portable sequencer, which Oxford Nanopore Technologies released in early 2014 as part of an early access program. A single 18 hr run can produce >90 Mb of data from around 16,000 total reads, with median and maximum read lengths of ~6 kb and >60 kb, respectively (Ash-ton et al., 2015). As with all single-molecule sequencing methodologies, error rates are high. Jain and colleagues most recently reported insertion, deletion, and substitution rates of 4.9%, 7.8%, and 5.1%, respectively (Jain et al., 2015). Presently, it also has a very high run failure rate. Despite the high error rates, MinION reads have been successfully used to determine the position and structure of a bacterial resistance island in combination with Illumina-derived reads (Ashton et al., 2015) and resolve an assembly gap on human Xq24 (Jain et al., 2015). Given the relatively high error rates and low throughput, nanopore sequencing is unlikely to overtake current sequencing platforms in the near future; however, the combination of size, speed, read lengths, and machine cost hold promise for the future.

The Development and Use of HTS Applications

As sequencing costs have fallen, HTS machines have become widely present in university core facilities and even individual labs. Decreasing costs and increased accessibility have enabled researchers to develop a rich catalog of HTS applications (Figure 4; Table 1). Some of these technologies were initially developed using DNA microarrays, but many are enabled only by using sequencing. HTS offers many advantages over DNA microarrays. In particular, it is more precise and not subject to cross-hybridization, thereby providing higher accuracy and a larger dynamic range (>10⁵ for DNA sequencing versus 10² for DNA microarrays) (Wang et al., 2009). Similar to microarrays, however, HTS-based applications can be biased by a number of variables, such as sequencing platform and library preparation method. The Sequencing Quality Control Consortium and similar initiatives are designed to study these biases and develop approaches to control for them, as has been recently demonstrated for RNA-seq (Su et al., 2014).

As HTS-based applications have become more robust, they have not only enabled individual researchers but also a variety of consortia-based projects. These large-scale projects have both provided valuable resources to the community and also have addressed questions that would be difficult for individual




Figure 4. Overview of Selected HTS Applications

Publication date of a representative article describing a method versus the number of citations that the article received. Methods are colored by category, and the size of the data point is proportional to publication rate (citations/months). The inset indicates the color key as well the proportion of methods in each group. For clarity, seq has been omitted from the labels.

labs to approach. Some of these projects are listed in Table 2 and include efforts to characterize the human genome (ENCODE, Roadmap Epigenomics Project, 2015), study human genetic variation (The 1000 Genomes Project Consortium, 2010), analyze gene expression (GTEx), and discover the molecular underpinnings of human disease (many; see Table 2). These coordinated efforts produce foundational resources that are of high utility to the scientific community by depositing the data into easily accessible public databases. Moreover, consortia often implement robust experimental and computational standards (e.g., Landt et al., 2012), ensuring high-quality data. Use of HTS applications by both individual laboratories and the large consortia have enabled researchers to illuminate previously intractable topics in biology, some of which are discussed below.

Genome Sequencing and Variation

The utility of HTS technologies for determining genome sequences de novo was first demonstrated by sequencing the genome of *Acinetobacter baumannii* (Smith et al., 2007). As the technologies and throughput improved, they were applied to “resequencing” human genomes and exomes, which was accomplished by first mapping reads to a reference genome and then identifying variants that differ between the sample genome and the reference (Wheeler et al., 2008). The different genome sequencing projects have since revealed that individuals typically harbor 3.5–4 million single nucleotide variants (SNVs) in total and several hundred thousand short indels relative

to the reference genome. Importantly, these variants include hundreds of loss of function alterations in genes (The 1000 Genomes Project Consortium, 2010).

HTS has also been used to globally characterize SV in the human genome. SVs include large (>1 kb) segments of the genome that have been duplicated, deleted, or rearranged. The short read lengths of most HTS platforms make determining SVs and indels more challenging than SNVs (Snyder et al., 2010). Typically, at least four independent approaches are utilized to identify SVs in a genome. These approaches include depth of read coverage (Abyzov et al., 2011), mapping of paired end reads that are discordant from the reference genome (Korbel et al., 2007), identifying split reads (Zhang et al., 2011), and mapping of breakpoint junctions (Kidd et al., 2010). Although each method has shortcomings, the improvement in resolution over array-based approaches has greatly enhanced our understanding of the prevalence of SVs throughout the genome and their contribution to disease. However, because no method or combination of them is comprehensive, SVs are never characterized in their entirety, if at all, in most sequencing projects.

In addition to identifying variants, it is also useful to assign them to paternal and maternal alleles, or “phase” them. Similar to SVs, current read lengths hinder our ability to phase genomes. This limitation can be circumvented by several methods, including sequencing parents, sequencing proximity ligated fragments (Selvaraj et al., 2013), or dilution and barcoding strategies during template preparation to allow long read assembly (Kuleshov

Table 1. Selected HTS Methods

Method	Purpose	Reference
RNA-seq	Transcript analysis	Nagalakshmi et al., 2008
Global run-on sequencing (GRO-seq)	Transcription	Core et al., 2008
Nascent-seq	Transcription	Khodor et al., 2011
Native elongating transcript sequencing (NET-seq)	Transcription	Churchman and Weissman, 2011
Ribo-seq	Translation	Ingolia et al., 2009
Replication sequencing (Repli-seq)	Replication	Hansen et al., 2010
Hi-C	Chromatin conformation	Lieberman-Aiden et al., 2009
Chromatin interaction analysis by paired-end tag sequencing (ChIA-PET)	Chromatin conformation	Fullwood et al., 2009
Chromosome conformation capture carbon copy (5-C)	Chromatin conformation	Dostie et al., 2006
Chromatin isolation by RNA purification sequencing (ChIRP-seq)	Genome localization	Chu et al., 2011
Reduced representation bisulphite sequencing (RRBS-seq)	Genome methylation	Meissner et al., 2008
Bisulfite sequencing (BS-seq)	Genome methylation	Cokus et al., 2008
DNAse-seq	Open chromatin	Crawford et al., 2006
Assay for transposase-accessible chromatin using sequencing (ATAC-seq)	Open chromatin	Buenrostro et al., 2013
Parallel Analysis of RNA structure (PARS)	RNA structure	Kertesz et al., 2010
Structure-seq	RNA structure	Ding et al., 2014
RNA on a massively parallel array (RNA-MaP)	RNA-protein interactions	Buenrostro et al., 2014
RNA immunoprecipitation sequencing (RIP-seq)	RNA-protein interactions	Sephton et al., 2011
Parallel analysis of RNA ends sequencing (PARE-seq)	microRNA target discovery	German et al., 2008
Massively parallel functional dissection sequencing (MPFD)	Enhancer assay	Patwardhan et al., 2012

[et al., 2014](#); [Voskoboynik et al., 2013](#)). With approximately 30 Gb of additional sequence data, ~99% of the SNVs identified in a 50× genome can be phased into blocks that are 0.2–1 Mb in length ([Kuleshov et al., 2014](#)). Understanding the phase of variants can have important clinical implications when determining if multiple damaging variants affect both copies of a gene or only one copy. To date, HTS has been applied to many thousands of genomes and many tens of thousands of exomes, yielding tremendous insight into human diversity and disease.

Mapping Regulatory Information of the Genome

HTS has applications beyond simply sequencing genomes. Perhaps one of the highest impact areas is the genome-wide mapping of DNA regulatory elements at high resolution. The first of these technologies was ChIP-seq in which DNA associated with a transcription factor (TF) or chromatin modification is immuno-selected and then sequenced using HTS ([Johnson et al., 2007](#)). Mapping the sequences back to the genome reveals the location of bound regions or chromatin modifications. A more general method for discovering many putative regulatory regions is to map “open” regions of the genome using DNase I digestion, followed by DNA sequencing of the ends of the fragments ([Crawford et al., 2006](#)). This method identifies approximately 50% of regions that are TF-bound as measured by ChIP-seq ([Cheng et al., 2014](#)). DNase-seq, however, is quickly being replaced by Assay for Transposon Accessible Chromatin-seq (ATAC-seq) in which transposon-based insertion is used to map open chromatin regions with approximately 50 million mapped reads ([Buenrostro et al., 2013](#)). The ATAC-seq protocol is also simpler and can be applied to small numbers of cells, even single cells.

Regulatory information is especially revealing when compared across many individual genomes or within a single genome

across many cell or tissue types. Large-scale application of these methods by the ENCODE project has provided a wealth of invaluable information regarding TF binding networks ([ENCODE Project Consortium, 2012](#)), epigenetic maps ([Thurman et al., 2012](#)), and transcript annotations ([Djebali et al., 2012](#)). Moreover, recent studies have found more than 3.5 million regulatory elements located throughout the genome in different cell types. One of the most striking findings from these studies as a whole, however, was the higher than expected portion of the genome that appears to be functional. The exact percentage is a source of significant debate ([Doolittle, 2013](#)), highlighting the importance of further experimental evidence to assign function to genomic elements. Genome targeting techniques, such as CRISPR-Cas9 ([Gilbert et al., 2014](#)) as well as high-throughput enhancer assays ([Kheradpour et al., 2013](#)), provide researchers with new tools to interrogate putative regulatory elements. Nonetheless, a variety of lines of evidence (GWAS and ENCODE) suggest that the total amount of regulatory regions is likely greater than that of protein coding regions ([Kellis et al., 2014](#)).

Mapping the 3D Organization of the Genome

Our understanding of the global organization and compartmentalization of chromosomes has been profoundly advanced by HTS technologies. 3D chromatin interactions can be studied using a variety of HTS assays, such as chromatin interaction analysis by paired-end tag sequencing (ChIA-PET) and Hi-C ([Fullwood et al., 2009](#); [Lieberman-Aiden et al., 2009](#)). Each of these assays relies upon proximity-based ligation of cross-linked, sheared chromatin followed by sequencing to derive contact maps. Hi-C was the first technique to allow unbiased, genome-wide interrogation of chromatin organization and revealed that the genome broadly partitions into open and closed chromatin

Table 2. Examples of Consortia-Based Projects

Initiative	Purpose	Website
1000 Genomes Project	Cataloging normal variation in diverse human populations.	www.1000genomes.org
The Encyclopedia of DNA Elements	Identifying functional genomic elements in the human genome.	www.encodeproject.org
Roadmap Epigenomics Project	Catalogue human epigenomic data with the goal of advancing basic biology and disease-oriented research.	www.roadmapepigenomics.org
Human Microbiome Project	Comprehensive characterization of the human microbiome and analysis of its role in human health and disease.	www.hmpdacc.org
Genotype-Tissue Expression Program	Characterizing gene expression and regulation in many human tissues and correlating with genetic variation and disease.	www.commonfund.nih.gov/GTEX/index
Human Immunology Project Consortium	Characterizing the diverse states of the human immune system following infection, vaccination or treatment.	http://www.immuneprofiling.org
Grand Opportunity Exome Sequencing Project	Discovery of novel genes and mechanisms contributing to heart, lung and blood disorders.	https://esp.gs.washington.edu/drupal
The Cancer Genome Atlas	Understanding the molecular basis of cancer.	www.cancergenome.nih.gov
International Cancer Genome Consortium	Describing the genomic, transcriptomic and epigenomic changes in 50 different tumor types.	www.icgc.org
Clinical Sequencing Exploratory Research Program	Develop methods as well as the legal and ethical frameworks necessary to integrate sequencing into the clinic.	www.genome.gov/27546194
Centers for Mendelian Genomics	Discovering the genes and genetic variants underlying human Mendelian disorders.	www.mendelian.org
Undiagnosed Diseases Network	Promoting the use of genomic data to elucidate the mechanisms underlying the diseases of unknown etiology.	www.commonfund.nih.gov/Diseases/index
Newborn Sequencing in Genomic Medicine and Public Health	Exploring the challenges and opportunities associated with using genomic sequence information in the newborn period.	www.genome.gov/27558493
The Pediatric Cardiac Genomics Consortium	Determining the genes responsible for congenital heart disease.	www.benchtopassinet.com
Alzheimer's Disease Sequencing Project	Identifying genes contributing to risk of developing Alzheimer's disease in multiethnic populations.	www.niagads.org/adsp

states (Lieberman-Aiden et al., 2009). Hi-C also demonstrated that the genome is organized into topological associating domains (TADs), which show high amounts of intra-domain interactions but exhibit infrequent interactions across domain boundaries (Dixon et al., 2012). Interestingly, TAD organization is stable across cell types and evolutionarily conserved across species. The boundaries between TADs were also enriched for house-keeping genes and binding sites for the insulator protein CTCF, raising the possibility that the distribution of TADs is chromosomally encoded (Dixon et al., 2012).

Recent advancements to the Hi-C technique combined with extremely deep sequencing (billions of reads per sample) have produced much higher resolution contact maps (~1 kb), which refine TAD domain size from 1 Mb to less than 200 kb (Rao et al., 2014). These new Hi-C maps demonstrated intra-chromosomal looping events, often containing promoter-to-enhancer contacts that were associated with gene activation. Most loops were anchored with directionally oriented CTCF binding sites, suggesting a mechanistic role for CTCF in establishing stable loops. Strikingly, fewer than 10,000 looping events were observed genome wide, which is far smaller than previous estimates (Jin et al., 2013). Modeling of Hi-C data has also suggested a fractal globule chromatin state, a conformation that both maximizes packing while preserving the flexibility to access any genomic locus (Lieberman-Aiden et al., 2009).

Characterizing the Transcriptome

Our appreciation for the diverse cellular roles of RNA has been greatly enhanced by the advent of HTS. Much of this evolution in thought has been a direct result of the many HTS applications designed to systematically identify various classes of RNA as well as to characterize RNA structure, RNA-protein interactions, and genomic localization. Cap analysis of gene expression (CAGE) and RNA-seq have been utilized to great effect to deeply characterize transcriptomes, providing precise, comprehensive measurements for message abundance, isoform usage, RNA-editing, and allele-specific expression. Deep sequencing of RNA has suggested that roughly three-quarters of the human genome is transcribed (Djebali et al., 2012). Most of this transcription covers introns or is very low, non-coding and of unclear biologic significance. However, many interesting species of non-coding RNA, including long non-coding RNAs (lncRNAs), small nucleolar RNAs (snoRNAs), and micro-RNAs, have been systematically described with RNA-seq and derivative techniques. A subset of lncRNAs, for example, have been revealed by overlaying RNA-seq data with ChIP-seq profiles characteristic of expressed genes (Guttman et al., 2009). Building upon earlier cDNA sequencing and tiling array experiments, these HTS approaches expanded the list to include more than a thousand mammalian lncRNAs. Analogous expansions have occurred for many aspects of RNA biology,

such as the number of sites undergoing RNA editing (Li et al., 2009).

Understanding the structure and biology of these newly discovered transcripts has led to the development of additional HTS applications. For instance, microRNA-target discovery has been facilitated by sequencing signatures of miRNA-mediated mRNA decay, using parallel analysis of RNA ends (PAREs) (German et al., 2008). Furthermore, RNA immunoprecipitation chip (RIP-chip) and subsequently RIP-seq were utilized to show that approximately 20% of the lncRNAs associate with polycomb repressor complex 2 (PRC2), a chromatin-modifying complex (Khalil et al., 2009; Zhao et al., 2010). Given these links to chromatin, methods analogous to ChIP-seq were developed, such as chromatin isolation by RNA purification (ChIRP-seq), to determine the genomic localization of lncRNAs (Chu et al., 2011). HTS applications have also made it possible to determine transcript structure both in vitro (parallel analysis of RNA sequencing; PARS) and in vivo (Structure-seq), providing insight into the effects of various structural features on translation efficiency, splicing, and polyadenylation (Ding et al., 2014; Kertesz et al., 2010). More recently, systematic interrogation of sequence-function relationships for RNA-protein interactions has been made possible using a high-throughput biochemical assay called RNA on a massively parallel array (RNA-MaP) (Buenrostro et al., 2014). The use of these assays, and many others, have enabled researchers to study RNA biology both comprehensively and with great detail, thereby enhancing our appreciation for the varied roles RNA plays in normal cellular homeostasis as well as human disease.

Microbiome Sequencing

Advances in HTS have enabled extensive cataloging of metagenomic samples, providing insight into the diversity of microbial species from a wide variety of sources, including the ocean, soil, and human body. These studies use both 16S rRNA gene sequencing to determine phylogenetic relationships as well as more comprehensive shotgun sequencing to predict detailed species and gene composition. In particular, much attention has been paid to characterizing the diverse microbes resident to healthy human populations (Human Microbiome Project Consortium, 2012). These studies found extensive variation in both body site habitat and among different individuals, giving rise to the concept of a “personal microbiome.” Microbial diversity, or the number and abundance distribution of microorganisms in a given niche, also correlates with several human diseases. For instance, an increase in diversity is associated with bacterial vaginosis (Fredricks et al., 2005), whereas obesity and inflammatory bowel disease exhibit a decrease in the diversity of gut microbes (Qin et al., 2010; Turnbaugh et al., 2009). Although transplant studies in mice have demonstrated a direct link between the gut microbiome, energy metabolism, and obesity (Turnbaugh et al., 2006), causal relationships for the majority of human diseases are not well established. A deeper understanding will require more detailed characterizations of the dynamics of microbiomes across health states as well as more integrative studies to investigate the functional interplay between the microbiota, the host, and the environment.

Genome Sequencing of Rare Diseases

The capacity to sequence genomes, exomes, and transcriptomes has profoundly influenced our understanding of the ge-

netics of human disease, especially for rare Mendelian disorders and cancer. According to the Online Mendelian Inheritance in Man database, there are more than 7800 Mendelian disorders, but the causative gene for less than one half of these are known. By sequencing unrelated patients or affected and unaffected family members, early exome studies demonstrated the ability to identify causal alleles for a variety of inherited diseases (Bilgüvar et al., 2010; Ng et al., 2010). In rare cases, sequencing of patient samples has suggested specific clinical interventions that have dramatically altered patient outlook. In one early example, exome sequencing of a child with severe inflammatory bowel disease uncovered a mutation in an important regulator of inflammation, X-linked inhibitor of apoptosis (XIAP). Based on the severity of the child's symptoms as well as the molecular diagnosis, a bone marrow transplant was given to the patient, which subsequently alleviated his symptoms (Worthey et al., 2011). Despite the power of HTS for disease gene discovery, however, exome sequencing currently identifies the genetic defect in only 25% of cases (Yang et al., 2013).

Cancer Genome Sequencing

Cancer is another important arena where HTS has been applied to great effect. The Cancer Genome Atlas (TCGA) and International Cancer Genome Consortium (ICGC) have performed genome and exome sequencing on thousands of tumor-normal pairs. These studies have described the mutational landscapes for over 20 cancer types, demonstrating that tumors can vary dramatically in both the type and quantity of mutations (Lawrence et al., 2014). These global descriptions have been integral to the development of background mutation rates that are necessary for the detection of cancer driver genes. For example, replication timing and gene expression were both found to be important covariates when determining if a gene is mutated at a rate higher than expected. Using these background models, TCGA-led projects discovered several novel cancer drivers, known drivers in new cancer types, and commonly disrupted pathways (Lawrence et al., 2014). Moreover, WGS of cancer samples has also identified high-frequency, non-coding mutations, such as activating mutations in the TERT promoter (Huang et al., 2013), a poorly characterized but highly relevant class of somatic variants.

The scale and sensitivity of HTS has also enabled global descriptions of tumor heterogeneity, clonal evolution, and the mechanisms underlying drug resistance. By tracking copy number aberrations in primary breast cancer cells using single-cell sequencing techniques, Navin and colleagues demonstrated that copy number rearrangements can occur in bursts, followed by persistent clonal expansion (Navin et al., 2011). Point mutations, in contrast, appear to accumulate more slowly over time, giving rise to more extensive clonal diversity, which may enable the tumor to adapt to diverse selective pressures (Wang et al., 2014). In addition to examining clonal diversity, HTS has also been used to compare primary tumors with relapse lesions, allowing characterization of the effects of chemotherapy as well as the molecular mechanisms underlying resistance to therapy (Van Allen et al., 2014; Ding et al., 2012). Together, these molecular portraits of cancer are forming the foundation of new paradigms for the diagnosis and treatment of cancer.

Limitations of Current HTS Technologies

It is becoming increasingly clear that while the technologies of today may be capable of providing population-level sequencing to both researchers and clinicians, key limitations remain. From a technological perspective, accuracy and coverage across the genome are still problematic, particularly for GC-rich regions and long homopolymer stretches (Ross et al., 2013). In addition, the short read lengths produced by most current platforms severely limit our ability to accurately characterize large repeat regions, many indels, and SV, leaving significant portions of the genome opaque or inaccurate (Snyder et al., 2010). The establishment of a gold standard genome, as envisioned by the Genome in a Bottle Consortium (Zook and Salit, 2011) as well as standards for data processing, variant calling, and reporting as set out in the CLARITY Challenge (Brownstein et al., 2014), will be valuable for comparing and reporting the accuracy of different platforms and studies. Given the limitations and biases of different platforms, it is also likely that accurate genome sequencing will use a combination of technologies.

In addition to genomes, quantitative analysis of complete transcriptomes, with individual allelic and spliced isoforms, is hindered by short reads. Improvements in the throughput and accuracy of current long-read technologies, such as Pacific Biosciences and Oxford Nanopore Technologies, as well as the use of “synthetic long-read methods” in which longer fragments can be sequenced and assembled from short reads will help overcome these limitations (Tilgner et al., 2015). Although both the research and medical communities are pressing forward with current technologies, these limitations will also continue to drive the innovation of new sequencing platforms (reviewed by Schadt et al., 2010).

HTS in the Coming Era of Personalized Medicine

To date, clinical HTS has most often been employed on focused regions of the genome or in the context of small pathogen identification. For instance, prenatal tests designed to non-invasively detect chromosomal abnormalities in cell-free DNA from maternal blood are clinically available (e.g., Ariosa Diagnostics' Harmony Test and BGI's NIFTY Test). Similarly, targeted HTS of clinically actionable mutations is being utilized to guide the diagnosis and treatment of cancer (e.g., Foundation Medicine's FoundationONE test). HTS has also been employed in clinical contexts to monitor pathogen outbreaks, such as methicillin-resistant *S. aureus* infections (Köser et al., 2012). The development and use of these focused assays will continue to expand, but the full promise of personalized medicine relies upon the routine clinical application of more comprehensive techniques, such as WGS, which still faces significant challenges.

In order for large-scale genomics to become fully integrated into the clinic, we need to reduce the costs and timescales associated with storage and interpretation of genome data. Most importantly, however, we must improve our ability to understand the biological and clinical consequences of variants of unknown significance. This class of alterations is the most common in personal genome sequences and includes novel variants that affect the coding sequence of known disease-causing genes but can also refer to variants in genes previously unlinked to disease or in regulatory regions of the genome. Interpretation of these var-

iants will benefit from additional genome sequencing as well as the data provided by large-scale genomics projects, such as ENCODE and GTEx, which enable the generation of more complete reference databases. Open access projects, such as the Personal Genomes Project and integrative Personal Omics Profiling (iPOP), will also provide valuable community resources for linking phenotypes to sequences (Chen et al., 2012; Church, 2005). The incorporation of high-throughput biochemical measurements of novel variation and detailed health records along with open data sharing will maximize our ability to both interpret personal genomes and better understand human health and disease.

ACKNOWLEDGMENTS

We would like to acknowledge C. Araya, C. Cenik, D. Webster, and P. Dumesic for helpful discussions regarding the manuscript. Work in Snyder lab is supported by NIH grants to M.S. (1P50HG00773501 and 8U54DK10255602). J.R. was supported by the Damon Runyon Cancer Research Foundation, and D.V.S. was supported by an NIH T32 fellowship (HG000044) and a Genentech Graduate Fellowship.

REFERENCES

- Abyzov, A., Urban, A.E., Snyder, M., and Gerstein, M. (2011). CNVnator: an approach to discover, genotype, and characterize typical and atypical CNVs from family and population genome sequencing. *Genome Res.* 21, 974–984.
- Ashton, P.M., Nair, S., Dallman, T., Rubino, S., Rabsch, W., Mwaigwisya, S., Wain, J., and O'Grady, J. (2015). MinION nanopore sequencing identifies the position and structure of a bacterial antibiotic resistance island. *Nat. Biotechnol.* 33, 296–300.
- Bentley, D.R., Balasubramanian, S., Swerdlow, H.P., Smith, G.P., Milton, J., Brown, C.G., Hall, K.P., Evers, D.J., Barnes, C.L., Bignell, H.R., et al. (2008). Accurate whole human genome sequencing using reversible terminator chemistry. *Nature* 456, 53–59.
- Bilgüvar, K., Öztürk, A.K., Louvi, A., Kwan, K.Y., Choi, M., Tatli, B., Yalnizoglu, D., Tüysüz, B., Çağlayan, A.O., Gökben, S., et al. (2010). Whole-exome sequencing identifies recessive WDR62 mutations in severe brain malformations. *Nature* 467, 207–210.
- Brownstein, C.A., Beggs, A.H., Homer, N., Merriman, B., Yu, T.W., Flannery, K.C., DeChene, E.T., Towne, M.C., Savage, S.K., Price, E.N., et al. (2014). An international effort towards developing standards for best practices in analysis, interpretation and reporting of clinical genome sequencing results in the CLARITY Challenge. *Genome Biol.* 15, R53.
- Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y., and Greenleaf, W.J. (2013). Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat. Methods* 10, 1213–1218.
- Buenrostro, J.D., Araya, C.L., Chircus, L.M., Layton, C.J., Chang, H.Y., Snyder, M.P., and Greenleaf, W.J. (2014). Quantitative analysis of RNA-protein interactions on a massively parallel array reveals biophysical and evolutionary landscapes. *Nat. Biotechnol.* 32, 562–568.
- Carneiro, M.O., Russ, C., Ross, M.G., Gabriel, S.B., Nusbaum, C., and DePristo, M.A. (2012). Pacific biosciences sequencing technology for genotyping and variation discovery in human data. *BMC Genomics* 13, 375.
- Chaisson, M.J.P., Huddleston, J., Dennis, M.Y., Sudmant, P.H., Malig, M., Hormozdiari, F., Antonacci, F., Surti, U., Sandstrom, R., Boitano, M., et al. (2015). Resolving the complexity of the human genome using single-molecule sequencing. *Nature* 517, 608–611.
- Chen, R., Mias, G.I., Li-Pook-Than, J., Jiang, L., Lam, H.Y.K., Chen, R., Miriami, E., Karczewski, K.J., Hariharan, M., Dewey, F.E., et al. (2012). Personal omics profiling reveals dynamic molecular and medical phenotypes. *Cell* 148, 1293–1307.

- Cheng, Y., Ma, Z., Kim, B.-H., Wu, W., Cayting, P., Boyle, A.P., Sundaram, V., Xing, X., Dogan, N., Li, J., et al.; Mouse ENCODE Consortium (2014). Principles of regulatory information conservation between mouse and human. *Nature* 515, 371–375.
- Chimpanzee Sequencing and Analysis Consortium (2005). Initial sequence of the chimpanzee genome and comparison with the human genome. *Nature* 437, 69–87.
- Chinwalla, A., Cook, L., Delehaunty, K., Fewell, G., Fulton, L., Fulton, R., Graves, T., Hillier, L., Mardis, E., and McPherson, J. (2002). Initial sequencing and comparative analysis of the mouse genome. *Nature* 420, 520–562.
- Chu, C., Qu, K., Zhong, F.L., Artandi, S.E., and Chang, H.Y. (2011). Genomic maps of long noncoding RNA occupancy reveal principles of RNA-chromatin interactions. *Mol. Cell* 44, 667–678.
- Church, G.M. (2005). The personal genome project. *Mol. Syst. Biol.* 1. Published online December 13, 2005.
- Churchman, L.S., and Weissman, J.S. (2011). Nascent transcript sequencing visualizes transcription at nucleotide resolution. *Nature* 469, 368–373.
- Cokus, S.J., Feng, S., Zhang, X., Chen, Z., Merriman, B., Haudenschild, C.D., Pradhan, S., Nelson, S.F., Pellegrini, M., and Jacobsen, S.E. (2008). Shotgun bisulphite sequencing of the Arabidopsis genome reveals DNA methylation patterning. *Nature* 452, 215–219.
- Core, L.J., Waterfall, J.J., and Lis, J.T. (2008). Nascent RNA Sequencing Reveals Widespread Pausing and Divergent Initiation at Human Promoters. *Science* 322, 1845–1848.
- Crawford, G.E., Holt, I.E., Whittle, J., Webb, B.D., Tai, D., Davis, S., Margulies, E.H., Chen, Y., Bernat, J.A., Ginsburg, D., et al. (2006). Genome-wide mapping of DNase hypersensitive sites using massively parallel signature sequencing (MPSS). *Genome Res.* 16, 123–131.
- Ding, L., Ley, T.J., Larson, D.E., Miller, C.A., Koboldt, D.C., Welch, J.S., Ritchey, J.K., Young, M.A., Lamprecht, T., McLellan, M.D., et al. (2012). Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature* 481, 506–510.
- Ding, Y., Tang, Y., Kwok, C.K., Zhang, Y., Bevilacqua, P.C., and Assmann, S.M. (2014). In vivo genome-wide profiling of RNA secondary structure reveals novel regulatory features. *Nature* 505, 696–700.
- Dixon, J.R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J.S., and Ren, B. (2012). Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* 485, 376–380.
- Djebali, S., Davis, C.A., Merkel, A., Dobin, A., Lassmann, T., Mortazavi, A., Tanzer, A., Lagarde, J., Lin, W., Schlesinger, F., et al. (2012). Landscape of transcription in human cells. *Nature* 489, 101–108.
- Dohm, J.C., Lottaz, C., Borodina, T., and Himmelbauer, H. (2008). Substantial biases in ultra-short read data sets from high-throughput DNA sequencing. *Nucleic Acids Res.* 36, e105.
- Doolittle, W.F. (2013). Is junk DNA bunk? A critique of ENCODE. *Proc. Natl. Acad. Sci. USA* 110, 5294–5300.
- Dostie, J., Richmond, T.a, Arnaout, R.a, Selzer, R.R., Lee, W.L., Honan, T.a, Rubio, E.D., Krumm, A., Lamb, J., Nusbaum, C., et al. (2006). Chromosome Conformation Capture Carbon Copy (5C): a massively parallel solution for mapping interactions between genomic elements. *Genome Res.* 16, 1299–1309.
- Drmanac, R., Sparks, A.B., Callow, M.J., Halpern, A.L., Burns, N.L., Kermani, B.G., Carnevali, P., Nazarenko, I., Nilsen, G.B., Yeung, G., et al. (2010). Human genome sequencing using unchained base reads on self-assembling DNA nanoarrays. *Science* 327, 78–81.
- Eid, J., Fehr, A., Gray, J., Luong, K., Lyle, J., Otto, G., Peluso, P., Rank, D., Baybayan, P., Bettman, B., et al. (2009). Real-time DNA sequencing from single polymerase molecules. *Science* 323, 133–138.
- ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in the human genome. *Nature* 489, 57–74.
- English, A.C., Richards, S., Han, Y., Wang, M., Vee, V., Qu, J., Qin, X., Muzny, D.M., Reid, J.G., Worley, K.C., and Gibbs, R.A. (2012). Mind the gap: upgrading genomes with Pacific Biosciences RS long-read sequencing technology. *PLoS ONE* 7, e47768.
- Fang, G., Munera, D., Friedman, D.I., Mandlik, A., Chao, M.C., Banerjee, O., Feng, Z., Losic, B., Mahajan, M.C., Jabado, O.J., et al. (2012). Genome-wide mapping of methylated adenine residues in pathogenic *Escherichia coli* using single-molecule real-time sequencing. *Nat. Biotechnol.* 30, 1232–1239.
- Flusberg, B.A., Webster, D.R., Lee, J.H., Travers, K.J., Olivares, E.C., Clark, T.A., Korlach, J., and Turner, S.W. (2010). Direct detection of DNA methylation during single-molecule, real-time sequencing. *Nat. Methods* 7, 461–465.
- Fredricks, D.N., Fiedler, T.L., and Marrazzo, J.M. (2005). Molecular identification of bacteria associated with bacterial vaginosis. *N. Engl. J. Med.* 353, 1899–1911.
- Fullwood, M.J., Liu, M.H., Pan, Y.F., Liu, J., Xu, H., Mohamed, Y.B., Orlov, Y.L., Velkov, S., Ho, A., Mei, P.H., et al. (2009). An oestrogen-receptor- α -bound human chromatin interactome. *Nature* 462, 58–64.
- German, M.A., Pillay, M., Jeong, D.-H., Hetawal, A., Luo, S., Janardhanan, P., Kannan, V., Rymarquis, L.A., Nobuta, K., German, R., et al. (2008). Global identification of microRNA-target RNA pairs by parallel analysis of RNA ends. *Nat. Biotechnol.* 26, 941–946.
- Gibbs, R.A., Weinstock, G.M., Metzker, M.L., Muzny, D.M., Sodergren, E.J., Scherer, S., Scott, G., Steffen, D., Worley, K.C., Burch, P.E., et al.; Rat Genome Sequencing Project Consortium (2004). Genome sequence of the Brown Norway rat yields insights into mammalian evolution. *Nature* 428, 493–521.
- Gilbert, L.A., Horlbeck, M.A., Adamson, B., Villalta, J.E., Chen, Y., Whitehead, E.H., Guimaraes, C., Panning, B., Ploegh, H.L., Bassik, M.C., et al. (2014). Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell* 159, 647–661.
- Guo, J., Xu, N., Li, Z., Zhang, S., Wu, J., Kim, D.H., Sano Marma, M., Meng, Q., Cao, H., Li, X., et al. (2008). Four-color DNA sequencing with 3'-O-modified nucleotide reversible terminators and chemically cleavable fluorescent dideoxynucleotides. *Proc. Natl. Acad. Sci. USA* 105, 9145–9150.
- Guttman, M., Amit, I., Garber, M., French, C., Lin, M.F., Feldser, D., Huarte, M., Zuk, O., Carey, B.W., Cassady, J.P., et al. (2009). Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature* 458, 223–227.
- Hansen, R.S., Thomas, S., Sandstrom, R., Canfield, T.K., Thurman, R.E., et al. (2010). Sequencing newly replicated DNA reveals widespread plasticity in human replication timing. *Proc. Natl. Acad. Sci. USA* 107, 139–144.
- Harris, T.D., Buzby, P.R., Babcock, H., Beer, E., Bowers, J., Braslavsky, I., Causey, M., Colonell, J., Dimeo, J., Efcavitch, J.W., et al. (2008). Single-molecule DNA sequencing of a viral genome. *Science* 320, 106–109.
- Huang, F.W., Hodis, E., Xu, M.J., Kryukov, G.V., Chin, L., and Garraway, L.A. (2013). Highly recurrent TERT promoter mutations in human melanoma. *Science* 339, 957–959.
- Human Microbiome Project Consortium (2012). Structure, function and diversity of the healthy human microbiome. *Nature* 486, 207–214.
- Ingolia, N.T., Ghaemmaghami, S., Newman, J.R.S., and Weissman, J.S. (2009). Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. *Science* 324, 218–223.
- International Human Genome Sequencing Consortium (2004). Finishing the euchromatic sequence of the human genome. *Nature* 431, 931–945.
- Jain, M., Fiddes, I.T., Miga, K.H., Olsen, H.E., Paten, B., and Akeson, M. (2015). Improved data analysis for the MinION nanopore sequencer. *Nat. Methods* 12, 351–356.
- Jin, F., Li, Y., Dixon, J.R., Selvaraj, S., Ye, Z., Lee, A.Y., Yen, C.-A., Schmitt, A.D., Espinoza, C.A., and Ren, B. (2013). A high-resolution map of the three-dimensional chromatin interactome in human cells. *Nature* 503, 290–294.
- Johnson, D.S., Mortazavi, A., Myers, R.M., and Wold, B. (2007). Genome-wide mapping of in vivo protein-DNA interactions. *Science* 316, 1497–1502.
- Kellis, M., Wold, B., Snyder, M.P., Bernstein, B.E., Kundaje, A., Marinov, G.K., Ward, L.D., Birney, E., Crawford, G.E., Dekker, J., et al. (2014). Defining

- functional DNA elements in the human genome. *Proc. Natl. Acad. Sci. USA* 111, 6131–6138.
- Kertesz, M., Wan, Y., Mazon, E., Rinn, J.L., Nutter, R.C., Chang, H.Y., and Segal, E. (2010). Genome-wide measurement of RNA secondary structure in yeast. *Nature* 467, 103–107.
- Khalil, A.M., Guttman, M., Huarte, M., Garber, M., Raj, A., Rivea Morales, D., Thomas, K., Presser, A., Bernstein, B.E., van Oudenaarden, A., et al. (2009). Many human large intergenic noncoding RNAs associate with chromatin-modifying complexes and affect gene expression. *Proc. Natl. Acad. Sci. USA* 106, 11667–11672.
- Kheradpour, P., Ernst, J., Melnikov, A., Rogov, P., Wang, L., Zhang, X., Alston, J., Mikkelsen, T.S., and Kellis, M. (2013). Systematic dissection of regulatory motifs in 2000 predicted human enhancers using a massively parallel reporter assay. *Genome Res.* 23, 800–811.
- Khodor, Y.L., Rodriguez, J., Abruzzi, K.C., Tang, C.H.A., Marr, M.T., and Rosbash, M. (2011). Nascent-seq indicates widespread cotranscriptional pre-mRNA splicing in *Drosophila*. *Genes Dev.* 25, 2502–2512.
- Kidd, J.M., Graves, T., Newman, T.L., Fulton, R., Hayden, H.S., Malig, M., Kallicki, J., Kaul, R., Wilson, R.K., and Eichler, E.E. (2010). A human genome structural variation sequencing resource reveals insights into mutational mechanisms. *Cell* 143, 837–847.
- Korbel, J.O., Urban, A.E., Affourtit, J.P., Godwin, B., Grubert, F., Simons, J.F., Kim, P.M., Palejev, D., Carriero, N.J., Du, L., et al. (2007). Paired-end mapping reveals extensive structural variation in the human genome. *Science* 318, 420–426.
- Koren, S., Schatz, M.C., Walenz, B.P., Martin, J., Howard, J.T., Ganapathy, G., Wang, Z., Rasko, D.A., McCombie, W.R., and Jarvis, E.D.; Adam M Phillippy (2012). Hybrid error correction and de novo assembly of single-molecule sequencing reads. *Nat. Biotechnol.* 30, 693–700.
- Köser, C.U., Holden, M.T.G., Ellington, M.J., Cartwright, E.J.P., Brown, N.M., Ogilvy-Stuart, A.L., Hsu, L.Y., Chewapreecha, C., Croucher, N.J., Harris, S.R., et al. (2012). Rapid whole-genome sequencing for investigation of a neonatal MRSA outbreak. *N. Engl. J. Med.* 366, 2267–2275.
- Kuleshov, V., Xie, D., Chen, R., Pushkarev, D., Ma, Z., Blauwkamp, T., Kertesz, M., and Snyder, M. (2014). Whole-genome haplotyping using long reads and statistical methods. *Nat. Biotechnol.* 32, 261–266.
- Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., et al.; International Human Genome Sequencing Consortium (2001). Initial sequencing and analysis of the human genome. *Nature* 409, 860–921.
- Landt, S.G., Marinov, G.K., Kundaje, A., Kheradpour, P., Pauli, F., Batzoglou, S., Bernstein, B.E., Bickel, P., Brown, J.B., Cayting, P., et al. (2012). ChIP-seq guidelines and practices of the ENCODE and modENCODE consortia. *Genome Res.* 22, 1813–1831.
- Lawrence, M.S., Stojanov, P., Mermel, C.H., Robinson, J.T., Garraway, L.A., Golub, T.R., Meyerson, M., Gabriel, S.B., Lander, E.S., and Getz, G. (2014). Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature* 505, 495–501.
- Levene, M.J., Korlach, J., Turner, S.W., Foquet, M., Craighead, H.G., and Webb, W.W. (2003). Zero-mode waveguides for single-molecule analysis at high concentrations. *Science* 299, 682–686.
- Li, J.B., Levanon, E.Y., Yoon, J.-K., Aach, J., Xie, B., LeProust, E., Zhang, K., Gao, Y., and Church, G.M. (2009). Genome-wide identification of human RNA editing sites by parallel DNA capturing and sequencing. *Science* 324, 1210–1213.
- Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B.R., Sabo, P.J., Dorschner, M.O., et al. (2009). Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* 326, 289–293.
- Liu, L., Li, Y., Li, S., Hu, N., He, Y., Pong, R., Lin, D., Lu, L., and Law, M. (2012). Comparison of next-generation sequencing systems. *J. Biomed. Biotechnol.* 2012, <http://dx.doi.org/10.1155/2012/251364>.
- Loomis, E.W., Eid, J.S., Peluso, P., Yin, J., Hickey, L., Rank, D., Mccalmon, S., Hagerman, R.J., Tassone, F., and Hagerman, P.J. (2013). Sequencing the unsequenceable: expanded CGG-repeat alleles of the fragile X gene. *Genome Res.* 23, 121–128.
- Margulies, M., Egholm, M., Altman, W.E., Attiya, S., Bader, J.S., Bembien, L.A., Berka, J., Braverman, M.S., Chen, Y.J., Chen, Z., et al. (2005). Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437, 376–380.
- Meissner, A., Mikkelsen, T.S., Gu, H., Wernig, M., Hanna, J., Sivachenko, A., Zhang, X., Bernstein, B.E., Nusbaum, C., Jaffe, D.B., et al. (2008). Genome-scale DNA methylation maps of pluripotent and differentiated cells. *Nature* 454, 766–770.
- Mellmann, A., Harmsen, D., Cummings, C.A., Zentz, E.B., Leopold, S.R., Rico, A., Prior, K., Szczepanowski, R., Ji, Y., Zhang, W., et al. (2011). Prospective genomic characterization of the German enterohemorrhagic *Escherichia coli* O104:H4 outbreak by rapid next generation sequencing technology. *PLoS ONE* 6, e22751.
- Metzker, M.L. (2010). Sequencing technologies - the next generation. *Nat. Rev. Genet.* 11, 31–46.
- Morey, M., Fernández-Marmiesse, A., Castiñeiras, D., Fraga, J.M., Couce, M.L., and Cocho, J.A. (2013). A glimpse into past, present, and future DNA sequencing. *Mol. Genet. Metab.* 110, 3–24.
- Nagalakshmi, U., Wang, Z., Waern, K., Shou, C., Raha, D., Gerstein, M., and Snyder, M. (2008). The transcriptional landscape of the yeast genome defined by RNA sequencing. *Science* 320, 1344–1349.
- Navin, N., Kendall, J., Troge, J., Andrews, P., Rodgers, L., McIndoo, J., Cook, K., Stepansky, A., Levy, D., Esposito, D., et al. (2011). Tumour evolution inferred by single-cell sequencing. *Nature* 472, 90–94.
- Ng, S.B., Bigham, A.W., Buckingham, K.J., Hannibal, M.C., McMillin, M.J., Gildersleeve, H.I., Beck, A.E., Tabor, H.K., Cooper, G.M., Mefford, H.C., et al. (2010). Exome sequencing identifies MLL2 mutations as a cause of Kabuki syndrome. *Nat. Genet.* 42, 790–793.
- Patwardhan, R.P., Hiatt, J.B., Witten, D.M., Kim, M.J., Smith, R.P., May, D., Lee, C., Andrie, J.M., Lee, S.-I., Cooper, G.M., et al. (2012). Massively parallel functional dissection of mammalian enhancers in vivo. *Nat. Biotechnol.* 30, 265–270.
- Qin, J., Li, R., Raes, J., Arumugam, M., Burgdorf, K.S., Manichanh, C., Nielsen, T., Pons, N., Levenez, F., Yamada, T., et al.; MetaHIT Consortium (2010). A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 464, 59–65.
- Quick, J., Quinlan, A.R., and Loman, N.J. (2014). A reference bacterial genome dataset generated on the MinION TM portable single-molecule nanopore sequencer. 3, <http://dx.doi.org/10.1186/2047-217X-3-22>.
- Rao, S.S.P., Huntley, M.H., Durand, N.C., Stamenova, E.K., Bochkov, I.D., Robinson, J.T., Sanborn, A.L., Machol, I., Omer, A.D., Lander, E.S., and Aiden, E.L. (2014). A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* 159, 1665–1680.
- Roadmap Epigenomics Consortium, Kundaje, A., Meuleman, W., Ernst, J., Bilenyk, M., Yen, A., Heravi-Moussavi, A., Kheradpour, P., Zhang, Z., Wang, J., et al. (2015). Integrative analysis of 111 reference human epigenomes. *Nature* 518, 317–330.
- Ross, M.G., Russ, C., Costello, M., Hollinger, A., Lennon, N.J., Hegarty, R., Nusbaum, C., and Jaffe, D.B. (2013). Characterizing and measuring bias in sequence data. *Genome Biol.* 14, R51.
- Rothberg, J.M., Hinz, W., Rearick, T.M., Schultz, J., Mileski, W., Davey, M., Leamon, J.H., Johnson, K., Milgrew, M.J., Edwards, M., et al. (2011). An integrated semiconductor device enabling non-optical genome sequencing. *Nature* 475, 348–352.
- Schadt, E.E., Turner, S., and Kasarskis, A. (2010). A window into third-generation sequencing. *Hum. Mol. Genet.* 19 (R2), R227–R240.
- Schloss, J.A. (2008). How to get genomes at one ten-thousandth the cost. *Nat. Biotechnol.* 26, 1113–1115.
- Selvaraj, S., R Dixon, J., Bansal, V., and Ren, B. (2013). Whole-genome haplotype reconstruction using proximity-ligation and shotgun sequencing. *Nat. Biotechnol.* 31, 1111–1118.

- Sephton, C.F., Cenik, C., Kucukural, A., Dammer, E.B., Cenik, B., Han, Y., Dewey, C.M., Roth, F.P., Herz, J., Peng, J., et al. (2011). Identification of neuronal RNA targets of TDP-43-containing ribonucleoprotein complexes. *J. Biol. Chem.* 286, 1204–1215.
- Sharon, D., Tilgner, H., Grubert, F., and Snyder, M. (2013). A single-molecule long-read survey of the human transcriptome. *Nat. Biotechnol.* 31, 1009–1014.
- Shendure, J., Porreca, G.J., Reppas, N.B., Lin, X., McCutcheon, J.P., Rose-nbaum, A.M., Wang, M.D., Zhang, K., Mitra, R.D., and Church, G.M. (2005). Accurate multiplex polony sequencing of an evolved bacterial genome. *Science* 309, 1728–1732.
- Smith, M.G., Gianoulis, T.A., Pukatzki, S., Mekalanos, J.J., Ornston, L.N., Gerstein, M., and Snyder, M. (2007). New insights into *Acinetobacter baumannii* pathogenesis revealed by high-density pyrosequencing and transposon mutagenesis. *Genes Dev.* 21, 601–614.
- Snyder, M., Du, J., and Gerstein, M. (2010). Personal genome sequencing: current approaches and challenges. *Genes Dev.* 24, 423–431.
- Su, Z., Łabaj, P.P., Li, S., Thierry-Mieg, J., Thierry-Mieg, D., Shi, W., Wang, C., Schroth, G.P., Setterquist, R.A., Thompson, J.F., et al.; SEQC/MAQC-III Consortium (2014). A comprehensive assessment of RNA-seq accuracy, reproducibility and information content by the Sequencing Quality Control Consortium. *Nat. Biotechnol.* 32, 903–914.
- The 1000 Genomes Project Consortium (2010). A map of human genome variation from population-scale sequencing. *Nature* 467, 1061–1073.
- Thurman, R.E., Rynes, E., Humbert, R., Vierstra, J., Maurano, M.T., Haugen, E., Sheffield, N.C., Stergachis, A.B., Wang, H., Vernot, B., et al. (2012). The accessible chromatin landscape of the human genome. *Nature* 489, 75–82.
- Tilgner, H., Jahanbani, F., Blauwkamp, T., Moshrefi, A., Jaeger, E., Chen, F., Harel, I., Bustamante, C., Rasmussen, M., and Snyder, M. (2015). Comprehensive transcriptome analysis using synthetic long read sequencing reveals molecular co-association of distant splicing events. *Nat. Biotechnol.* 33.
- Travers, K.J., Chin, C.S., Rank, D.R., Eid, J.S., and Turner, S.W. (2010). A flexible and efficient template format for circular consensus sequencing and SNP detection. *Nucleic Acids Res.* 38, e159.
- Turnbaugh, P.J., Ley, R.E., Mahowald, M.A., Magrini, V., Mardis, E.R., and Gordon, J.I. (2006). An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 444, 1027–1031.
- Turnbaugh, P.J., Hamady, M., Yatsunenko, T., Cantarel, B.L., Duncan, A., Ley, R.E., Sogin, M.L., Jones, W.J., Roe, B.A., Affourtit, J.P., et al. (2009). A core gut microbiome in obese and lean twins. *Nature* 457, 480–484.
- Uemura, S., Aitken, C.E., Korlach, J., Flusberg, B.A., Turner, S.W., and Puglisi, J.D. (2010). Real-time tRNA transit on single translating ribosomes at codon resolution. *Nature* 464, 1012–1017.
- Valouev, A., Ichikawa, J., Tonthat, T., Stuart, J., Ranade, S., Peckham, H., Zeng, K., Malek, J.A., Costa, G., McKernan, K., et al. (2008). A high-resolution, nucleosome position map of *C. elegans* reveals a lack of universal sequence-dictated positioning. *Genome Res.* 18, 1051–1063.
- Van Allen, E.M., Wagle, N., Sucker, A., Treacy, D.J., Johannessen, C.M., Goetz, E.M., Place, C.S., Taylor-Weiner, A., Whittaker, S., Kryukov, G.V., et al.; Dermatologic Cooperative Oncology Group of Germany (DeCOG) (2014). The genetic landscape of clinical resistance to RAF inhibition in metastatic melanoma. *Cancer Discov.* 4, 94–109.
- Venter, J.C., Adams, M.D., Myers, E.W., Li, P.W., Mural, R.J., Sutton, G.G., Smith, H.O., Yandell, M., Evans, C.A., Holt, R.A., et al. (2001). The sequence of the human genome. *Science* 291, 1304–1351.
- Voskoboinik, A., Neff, N.F., Sahoo, D., Newman, A.M., Pushkarev, D., Koh, W., Passarelli, B., Fan, H.C., Mantalas, G.L., Palmeri, K.J., et al. (2013). The genome sequence of the colonial chordate, *Botryllus schlosseri*. *Elife* 2, e00569.
- Wang, Z., Gerstein, M., and Snyder, M. (2009). RNA-Seq: a revolutionary tool for transcriptomics. *Nat. Rev. Genet.* 10, 57–63.
- Wang, Y., Waters, J., Leung, M.L., Unruh, A., Roh, W., Shi, X., Chen, K., Scheet, P., Vattathil, S., Liang, H., et al. (2014). Clonal evolution in breast cancer revealed by single nucleus genome sequencing. *Nature* 512, 155–160.
- Wang, Y., Yang, Q., and Wang, Z. (2015). The evolution of nanopore sequencing. *Front. Genet.* 5, 449.
- Wheeler, D.A., Srinivasan, M., Egholm, M., Shen, Y., Chen, L., McGuire, A., He, W., Chen, Y.-J., Makhijani, V., Roth, G.T., et al. (2008). The complete genome of an individual by massively parallel DNA sequencing. *Nature* 452, 872–876.
- Worthe, E.A., Mayer, A.N., Syverson, G.D., Helbling, D., Bonacci, B.B., Decker, B., Serpe, J.M., Dasu, T., Tschannen, M.R., Veith, R.L., et al. (2011). Making a definitive diagnosis: successful clinical application of whole exome sequencing in a child with intractable inflammatory bowel disease. *Genet. Med.* 13, 255–262.
- Yang, Y., Muzny, D.M., Reid, J.G., Bainbridge, M.N., Willis, A., Ward, P.A., Braxton, A., Beuten, J., Xia, F., Niu, Z., et al. (2013). Clinical whole-exome sequencing for the diagnosis of mendelian disorders. *N. Engl. J. Med.* 369, 1502–1511.
- Zhang, Z.D., Du, J., Lam, H., Abyzov, A., Urban, A.E., Snyder, M., and Gerstein, M. (2011). Identification of genomic indels and structural variations using split reads. *BMC Genomics* 12, 375.
- Zhao, J., Ohsumi, T.K., Kung, J.T., Ogawa, Y., Grau, D.J., Sarma, K., Song, J.J., Kingston, R.E., Borowsky, M., and Lee, J.T. (2010). Genome-wide identification of polycomb-associated RNAs by RIP-seq. *Mol. Cell* 40, 939–953.
- Zook, J.M., and Salit, M. (2011). Genomes in a bottle: creating standard reference materials for genomic variation - why, what and how? *Genome Biol.* 12, 31.