

Supplemental Information

**Transcription Factors Activate Genes
through the Phase-Separation Capacity
of Their Activation Domains**

Ann Boija, Isaac A. Klein, Benjamin R. Sabari, Alessandra Dall'Agnese, Eliot L. Coffey, Alicia V. Zamudio, Charles H. Li, Krishna Shrinivas, John C. Manteiga, Nancy M. Hannett, Brian J. Abraham, Lena K. Afeyan, Yang E. Guo, Jenna K. Rimel, Charli B. Fant, Jurian Schuijers, Tong Ihn Lee, Dylan J. Taatjes, and Richard A. Young

Table S1. Related to Figure 1. Table of reported transcription factor-mediator subunit interactions.

Mediator subunits	Transcription factors								
MED1	OCT3/4 ¹	SOX2 ¹	GATA2 ²	MYC ³	P53 ⁴⁻⁶	PPARG ⁷	RARA ^{8,9}	ER1 ¹⁰⁻¹²	CEBPA ¹³
	VDR ^{8,14}	RXRA ^{8,9}	GATA1 ^{15,16}	THRA/THRB ^{8,9}	HNF4 ¹⁷	AHR ¹⁸	AR ¹⁹	SREBP1 ²⁰	NR1I3 ²¹
	RORa ²²	GR ^{21,23}	FXR ²⁴	Pit-1 ²	PPARA ^{8,9}	ER2 ¹⁰⁻¹²	KLF4 ¹		
MED12	ER1 ¹⁰	SOX9 ²⁵	THRA ²⁶	GLI3 ²⁷	NANOG ²⁸	ER2 ¹⁰	JUNB ²⁹	VDR ¹⁴	SREBF1 ²⁰
	RTA ³⁰	B-CATENIN ³¹	AICD ³²						
MED14	PPARG ³³	ER1 ¹⁰	VDR ¹⁴	HNF4 ¹⁷	ER2 ¹⁰	JUNB ²⁹	NR1I3 ³⁴	GATA1 ¹⁵	SREBF1 ²⁰
	GR ²¹								
MED15	P65 ²⁰	JUNB ²⁹	SREBF1 ³⁵						
MED16	THRA ²⁶	DIF ³⁶	VDR ¹⁴	MYC ³	JUNB ²⁹				
MED17	SREBF1 ³⁷	P65 ³⁸	FOS ²⁹	HNF4A ¹⁷	DIF ^{36,39}	RXR ³⁹	JUNB ²⁹	NR1I3 ³⁴	HSF ⁴⁰
	VP16 ²⁶								
MED19	REST ⁴¹								
MED21	SREBF1 ³⁷	P53 ⁴²	VDR ²⁶	THRA/THRB ⁴³					
MED23	SREBF1 ²⁰	ELF3 ⁴⁴	NR1I3 ³⁴	RB1 ⁴⁵	DIF ³⁶	HIF ⁴⁶	VDR ¹⁴	C/EBP ⁴⁷	P65 ²⁰
	ELK1 ^{48,49}	HSF ³⁵	ESX ⁴⁴						
MED25	VP16 ^{50,51}	DIF ³⁵	HSF ³⁵	RARA ⁵²	HNF4 ⁵³	SOX9 ⁵⁴			
MED26	SREBF1 ²⁰								
MED29	JUNB ²⁹	DSX ⁵⁵							
CDK8	THRA ²⁶	MYC ⁵⁶							

Adapted from Borggreve and Xue, 2011⁵⁷.

References

1. Apostolou, E. *et al.* Genome-wide chromatin interactions of the Nanog locus in pluripotency, differentiation, and reprogramming. *Cell Stem Cell* **12**, 699–712 (2013).
2. Gordon, D. F. *et al.* MED220/thyroid receptor-associated protein 220 functions as a transcriptional coactivator with Pit-1 and GATA-2 on the thyrotropin-beta promoter in thyrotropes. *Mol. Endocrinol.* **20**, 1073–89 (2006).
3. Liu, X., Vorontchikhina, M., Wang, Y.-L., Faiola, F. & Martinez, E. STAGA recruits Mediator to the MYC oncoprotein to stimulate transcription and cell proliferation. *Mol. Cell. Biol.* **28**, 108–21 (2008).
4. Meyer, K. D., Lin, S., Bernecky, C., Gao, Y. & Taatjes, D. J. p53 activates transcription by directing structural shifts in Mediator. *Nat. Struct. Mol. Biol.* **17**, 753–760 (2010).
5. Drané, P., Barel, M., Balbo, M. & Frade, R. Identification of RB18A, a 205 kDa new p53 regulatory protein which shares antigenic and functional properties with p53. *Oncogene* **15**, 3013–3024 (1997).
6. Frade, R., Balbo, M. & Barel, M. RB18A, whose gene is localized on chromosome 17q12-q21.1, regulates in vivo p53 transactivating activity. *Cancer Res.* **60**, 6585–9 (2000).
7. Ge, K. *et al.* Transcription coactivator TRAP220 is required for PPARγ2-stimulated adipogenesis. *Nature* **417**, 563–567 (2002).
8. Yuan, C. X., Ito, M., Fondell, J. D., Fu, Z. Y. & Roeder, R. G. The TRAP220 component of a thyroid hormone receptor-associated protein (TRAP) coactivator complex interacts directly with nuclear receptors in a ligand-dependent fashion. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 7939–44 (1998).
9. Zhu, X. G., McPhie, P., Lin, K. H. & Cheng, S. Y. The differential hormone-dependent

transcriptional activation of thyroid hormone receptor isoforms is mediated by interplay of their domains. *J. Biol. Chem.* **272**, 9048–54 (1997).

10. Kang, Y. K., Guermah, M., Yuan, C.-X. & Roeder, R. G. The TRAP/Mediator coactivator complex interacts directly with estrogen receptors and through the TRAP220 subunit and directly enhances estrogen receptor function in vitro. *Proc. Natl. Acad. Sci.* **99**, 2642–2647 (2002).
11. Jiang, P. *et al.* Key roles for MED1 LxxLL motifs in pubertal mammary gland development and luminal-cell differentiation. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 6765–70 (2010).
12. Burakov, D., Wong, C. W., Rachez, C., Cheskis, B. J. & Freedman, L. P. Functional interactions between the estrogen receptor and DRIP205, a subunit of the heteromeric DRIP coactivator complex. *J. Biol. Chem.* **275**, 20928–34 (2000).
13. Li, H. *et al.* The Med1 Subunit of Transcriptional Mediator Plays a Central Role in Regulating CCAAT/Enhancer-binding Protein- β -driven Transcription in Response to Interferon- γ . *J. Biol. Chem.* **283**, 13077–13086 (2008).
14. Rachez, C. *et al.* Ligand-dependent transcription activation by nuclear receptors requires the DRIP complex. *Nature* **398**, 824–8 (1999).
15. Stumpf, M. *et al.* The mediator complex functions as a coactivator for GATA-1 in erythropoiesis via subunit Med1/TRAP220. *Proc. Natl. Acad. Sci.* **103**, 18504–18509 (2006).
16. Crawford, S. E. *et al.* Defects of the Heart, Eye, and Megakaryocytes in Peroxisome Proliferator Activator Receptor-binding Protein (PBP) Null Embryos Implicate GATA Family of Transcription Factors. *J. Biol. Chem.* **277**, 3585–3592 (2002).
17. Malik, S., Wallberg, A. E., Kang, Y. K. & Roeder, R. G. TRAP/SMCC/mediator-dependent transcriptional activation from DNA and chromatin templates by orphan nuclear receptor hepatocyte nuclear factor 4. *Mol. Cell. Biol.* **22**, 5626–37 (2002).
18. Wang, S., Ge, K., Roeder, R. G. & Hankinson, O. Role of mediator in transcriptional activation by the aryl hydrocarbon receptor. *J. Biol. Chem.* **279**, 13593–600 (2004).
19. Wang, Q., Sharma, D., Ren, Y. & Fondell, J. D. A Coregulatory Role for the TRAP-Mediator Complex in Androgen Receptor-mediated Gene Expression. *J. Biol. Chem.* **277**, 42852–42858 (2002).
20. Näär, A. M. *et al.* Composite co-activator ARC mediates chromatin-directed transcriptional activation. *Nature* **398**, 828–32 (1999).
21. Hittelman, A. B., Burakov, D., Iñiguez-Lluhí, J. A., Freedman, L. P. & Garabedian, M. J. Differential regulation of glucocorticoid receptor transcriptional activation via AF-1-associated proteins. *EMBO J.* **18**, 5380–5388 (1999).
22. Atkins, G. B. *et al.* Coactivators for the Orphan Nuclear Receptor ROR α . *Mol. Endocrinol.* **13**, 1550–1557 (1999).
23. Chen, W. & Roeder, R. G. The Mediator subunit MED1/TRAP220 is required for optimal glucocorticoid receptor-mediated transcription activation. *Nucleic Acids Res.* **35**, 6161–9 (2007).
24. Pineda Torra, I., Freedman, L. P. & Garabedian, M. J. Identification of DRIP205 as a Coactivator for the Farnesoid X Receptor. *J. Biol. Chem.* **279**, 36184–36191 (2004).
25. Zhou, T. & Chiang, C.-M. Sp1 and AP2 regulate but do not constitute TATA-less human TAF(II)55 core promoter activity. *Nucleic Acids Res.* **30**, 4145–57 (2002).
26. Ito, M. *et al.* Identity between TRAP and SMCC complexes indicates novel pathways for the function of nuclear receptors and diverse mammalian activators. *Mol. Cell* **3**, 361–70 (1999).
27. Zhou, H., Kim, S., Ishii, S. & Boyer, T. G. Mediator Modulates Gli3-Dependent Sonic Hedgehog Signaling. *Mol. Cell. Biol.* **26**, 8667–8682 (2006).
28. Tutter, A. V. *et al.* Role for Med12 in regulation of Nanog and Nanog target genes. *J. Biol. Chem.* **284**, 3709–18 (2009).
29. Hein, M. Y. *et al.* A human interactome in three quantitative dimensions organized by stoichiometries and abundances. *Cell* **163**, 712–23 (2015).
30. Gwack, Y. *et al.* Principal role of TRAP/mediator and SWI/SNF complexes in Kaposi's sarcoma-

- associated herpesvirus RTA-mediated lytic reactivation. *Mol. Cell. Biol.* **23**, 2055–67 (2003).
31. Kim, S., Xu, X., Hecht, A. & Boyer, T. G. Mediator is a transducer of Wnt/beta-catenin signaling. *J. Biol. Chem.* **281**, 14066–75 (2006).
 32. Xu, X., Zhou, H. & Boyer, T. G. Mediator is a transducer of amyloid-precursor-protein-dependent nuclear signalling. *EMBO Rep.* **12**, 216–222 (2011).
 33. Grøntved, L., Madsen, M. S., Boergesen, M., Roeder, R. G. & Mandrup, S. MED14 tethers mediator to the N-terminal domain of peroxisome proliferator-activated receptor gamma and is required for full transcriptional activity and adipogenesis. *Mol. Cell. Biol.* **30**, 2155–69 (2010).
 34. Huttlin, E. L. *et al.* The BioPlex Network: A Systematic Exploration of the Human Interactome. *Cell* **162**, 425–440 (2015).
 35. Yang, F. *et al.* An ARC/Mediator subunit required for SREBP control of cholesterol and lipid homeostasis. *Nature* **442**, 700–704 (2006).
 36. Kim, T. W. *et al.* MED16 and MED23 of Mediator are coactivators of lipopolysaccharide- and heat-shock-induced transcriptional activators. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 12153–8 (2004).
 37. Taatjes, D. J., Näär, A. M., Andel, F., Nogales, E. & Tjian, R. Structure, function, and activator-induced conformations of the CRSP coactivator. *Science* **295**, 1058–62 (2002).
 38. van Essen, D., Engist, B., Natoli, G. & Sacconi, S. Two Modes of Transcriptional Activation at Native Promoters by NF- κ B p65. *PLoS Biol.* **7**, e1000073 (2009).
 39. Park, J. M. *et al.* Signal-induced transcriptional activation by Dif requires the dTRAP80 mediator module. *Mol. Cell. Biol.* **23**, 1358–67 (2003).
 40. Park, J. M., Werner, J., Kim, J. M., Lis, J. T. & Kim, Y. J. Mediator, not holoenzyme, is directly recruited to the heat shock promoter by HSF upon heat shock. *Mol. Cell* **8**, 9–19 (2001).
 41. Ding, N. *et al.* MED19 and MED26 are synergistic functional targets of the RE1 silencing transcription factor in epigenetic silencing of neuronal gene expression. *J. Biol. Chem.* **284**, 2648–56 (2009).
 42. Gu, W. *et al.* A novel human SRB/MED-containing cofactor complex, SMCC, involved in transcription regulation. *Mol. Cell* **3**, 97–108 (1999).
 43. Nevado, J., Tenbaum, S. P. & Aranda, A. hSrb7, an essential human Mediator component, acts as a coactivator for the thyroid hormone receptor. *Mol. Cell. Endocrinol.* **222**, 41–51 (2004).
 44. Asada, S. *et al.* External control of Her2 expression and cancer cell growth by targeting a Ras-linked coactivator. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 12747–52 (2002).
 45. Lambert, J.-P., Tucholska, M., Go, C., Knight, J. D. R. & Gingras, A.-C. Proximity biotinylation and affinity purification are complementary approaches for the interactome mapping of chromatin-associated protein complexes. *J. Proteomics* **118**, 81–94 (2015).
 46. Galbraith, M. D. *et al.* HIF1A employs CDK8-mediator to stimulate RNAPII elongation in response to hypoxia. *Cell* **153**, 1327–39 (2013).
 47. Mo, X., Kowenz-Leutz, E., Xu, H. & Leutz, A. Ras induces mediator complex exchange on C/EBP beta. *Mol. Cell* **13**, 241–50 (2004).
 48. Cantin, G. T., Stevens, J. L. & Berk, A. J. Activation domain-mediator interactions promote transcription preinitiation complex assembly on promoter DNA. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 12003–8 (2003).
 49. Stevens, J. L. *et al.* Transcription Control by E1A and MAP Kinase Pathway via Sur2 Mediator Subunit. *Science* (80-.). **296**, 755–758 (2002).
 50. Mittler, G. *et al.* A novel docking site on Mediator is critical for activation by VP16 in mammalian cells. *EMBO J.* **22**, 6494–504 (2003).
 51. Yang, F., DeBeaumont, R., Zhou, S. & Näär, A. M. The activator-recruited cofactor/Mediator coactivator subunit ARC92 is a functionally important target of the VP16 transcriptional activator. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 2339–44 (2004).
 52. Lee, H.-K., Park, U.-H., Kim, E.-J. & Um, S.-J. MED25 is distinct from TRAP220/MED1 in cooperating with CBP for retinoid receptor activation. *EMBO J.* **26**, 3545–3557 (2007).
 53. Rana, R., Surapureddi, S., Kam, W., Ferguson, S. & Goldstein, J. A. Med25 is required for RNA

polymerase II recruitment to specific promoters, thus regulating xenobiotic and lipid metabolism in human liver. *Mol. Cell. Biol.* **31**, 466–81 (2011).

54. Nakamura, Y. *et al.* Wwp2 is essential for palatogenesis mediated by the interaction between Sox9 and mediator subunit 25. *Nat. Commun.* **2**, 251 (2011).
55. Garrett-Engle, C. M. *et al.* intersex, a gene required for female sexual development in *Drosophila*, is expressed in both sexes and functions together with doublesex to regulate terminal differentiation. *Development* **129**, 4661–75 (2002).
56. Eberhardy, S. R. & Farnham, P. J. Myc Recruits P-TEFb to Mediate the Final Step in the Transcriptional Activation of the *cad* Promoter. *J. Biol. Chem.* **277**, 40156–40162 (2002).
57. Borggreffe, T. & Yue, X. Interactions between subunits of the Mediator complex with gene-specific transcription factors. *Semin. Cell Dev. Biol.* **22**, 759–768 (2011).