
a
3–10 bp
11–50 bp
3–50 bp
SpliceAI (published basis)
SpliceAI (distance 50)
Pangolin
AlphaGenome splice score
Atlas combined score †
Atlas splice sites
Atlas splice-site usage
Atlas splice junctions
CADD
phyloP
phastCons
GPN-MSA
Nucleotide Transformer
not reached in-sample
1
10
100
1000
Held-out gene: likelihood ratio at the fitted Moderate threshold (dotted: the Moderate boundary, 4.33)
b
SpliceAI (published basis)
SpliceAI (distance 50)
Pangolin
AlphaGenome splice score
Atlas combined score †
Atlas splice sites
Atlas splice-site usage
Atlas splice junctions
CADD
phyloP
phastCons
GPN-MSA
Nucleotide Transformer
not reached in-sample
no evaluable fold
Held-out gene: likelihood ratio at the fitted Strong threshold (dotted: the Strong boundary, 18.7)
c
3–10 bp
11–50 bp
3–50 bp
Moderate
Strong
SpliceAI, published 0.2
SpliceAI, fitted Moderate
SpliceAI, fitted Strong
Pangolin, fitted Moderate
Pangolin, fitted Strong
not reached in-sample
External gene: likelihood ratio at the published cut point and the fitted Moderate and Strong thresholds
Held-out gene, clears the boundary
Held-out gene, below it
Median of held-out genes
DDX3X
CTCF
SFPQ
RAD51D
XRCC2
SBDS
TP53 (HCT116 cells, Nutlin-3a)
Open: below the boundary