

Math 340 HW 2

Asa Royal (ajr74) [collaborators: none]

January 26, 2024

1. (i) In this setup, we are essentially modelling 120 flips of a p-coin. Each $\omega \in \Omega$ is a sequence of 120 true/false values. Thus,

$$\begin{aligned}\mathbb{P}(< 3 \text{ calls}) &= \mathbb{P}(0 \text{ calls}) + \mathbb{P}(1 \text{ calls}) + \mathbb{P}(2 \text{ calls}) \\ &= \binom{120}{0}(0.05)^0(0.95)^{120-0} + \binom{120}{1}(0.05)^1(0.95)^{120-1} + \binom{120}{2}(0.05)^2(0.95)^{120-2}\end{aligned}$$

- (ii) In this setup, $\omega \in \mathbb{W}$, and we can use a Poisson distribution with parameter $\lambda = 6$ to model the nubmer of arrivals in an hour. We choose $\lambda = 6$ so that $p = \lambda/n = 6/120 = 0.05$, which the last problem stated was the probability of an arrival during an interval. We have proved earlier that as $n \rightarrow \infty$, the $\mathbb{P}(A_k)$ follows the probability mass function stated in * if $\lambda = pn$. Once again, the probability of seeing < 3 calls during a 1 hour interval can be calculated by summing the probabilties of seeing zero, one, or two calls during the hour:

$$\mathbb{P}(< 3 \text{ calls}) = \mathbb{P}(0 \text{ calls}) + \mathbb{P}(1 \text{ calls}) + \mathbb{P}(2 \text{ calls})$$

* Per the probability mass function of the Poisson distribution, $\mathbb{P}(A_k) = \frac{\lambda^k}{k!}e^{-\lambda}$, so

$$\begin{aligned}\mathbb{P}(< 3 \text{ calls}) &= \frac{6^0}{0!}e^{-6} + \frac{6^1}{1!}e^{-6} + \frac{6^2}{2!}e^{-6} \\ &= e^{-6} + 6e^{-6} + 16e^{-6} \\ &= 23e^{-6} \\ &\approx 0.057\end{aligned}$$

2. (i) What is the probability that after n tosses of a p-coin, you have not seen heads?
This is the probability of seeing n tails in a row. Each p-coin flip is independent, so

$$\mathbb{P}(n \text{ tails}) = \prod_{k=1}^n \rho(T) = \prod_{k=1}^n (1-p) = (1-p)^n$$

- (ii) What is the probability that you toss $n-1$ tails and then heads occurs for the first time on the n th toss.
The first $n-1$ tosses are independent of the n th toss, so $\mathbb{P}(A)$, where A is the event described above, is $\mathbb{P}(n-1 \text{ tails}) * \mathbb{P}(\text{head})$. Per 2a, $\mathbb{P}(n-1 \text{ tails}) = (1-p)^{n-1}$. $\mathbb{P}(\text{head}) = p$ by construction. Thus,

$$\mathbb{P}(A) = (1-p)^{n-1} * p$$

3. (i) B_1 and B_2 are indeed independent. $\mathbb{P}(B_1 \cap B_2) =$
- (ii) Suppose your coin lands heads k times (out of the n tosses). What is the probability that your coin is the j th coin.
Let X_i be the event that we have the i th coin. Let A_k be the event that we flip k heads. Per Bayes' Rule

$$\begin{aligned}\mathbb{P}(X_j|A_k) &= \frac{\mathbb{P}(A_k|X_j) * \mathbb{P}(X_j)}{\sum_{i=1}^M \mathbb{P}(A_k|X_i) * \mathbb{P}(X_i)} \\ &= \frac{\left[\binom{n}{k}(p_j)^k(1-p_j)^{n-k}\right] \frac{1}{M}}{\sum_{i=1}^M \left[\binom{n}{k}(p_i)^k(1-p_i)^{n-k}\right] \frac{1}{M}} \\ &= \frac{\binom{n}{k}(p_j)^k(1-p_j)^{n-k}}{\sum_{i=1}^M \binom{n}{k}(p_i)^k(1-p_i)^{n-k}}\end{aligned}$$

- (iii) Suppose your coin lands heads n times in a row. You suspect it is the most biased coin (i.e. with the largest $p = p_M$). How large would n have to be in order that the probability of it being the most biased coin is at least ten times as large as the probability of it being the next most biased (i.e. the one with $p = p_{M-1}$)?

Note that

$$\mathbb{P}(A_n|X_m) = \binom{n}{n} (p_m)^n (1-p_m)^0 = (p_m)^n \quad (1)$$

$$\mathbb{P}(A_n|X_{m-1}) = \binom{n}{n} (p_{m-1})^n (1-p_{m-1})^0 = (p_{m-1})^n \quad (2)$$

If we are at least ten times likelier to have coin m vs coin $m-1$, we can set $\mathbb{P}(A_n|X_m) \geq 10 * \mathbb{P}(A_n|X_{m-1})$ and solve to find the minimum satisfying n .

$$\begin{aligned} \mathbb{P}(X_m|A_n) &\geq 10 * \mathbb{P}(X_{m-1}|A_n) && \text{problem setup} \\ \frac{\mathbb{P}(A_n|X_m) * \mathbb{P}(X_m)}{\mathbb{P}(A_n)} &\geq 10 \left(\frac{\mathbb{P}(A_n|X_{m-1}) * \mathbb{P}(X_{m-1})}{\mathbb{P}(A_n)} \right) && \text{Bayes' Rule} \\ \mathbb{P}(A_n|X_m) &\geq 10 * \mathbb{P}(A_n|X_{m-1}) && \text{cancel terms. } \mathbb{P}(X_m) = \mathbb{P}(X_{m-1}) = 1/m \\ (p_m)^n &\geq 10(p_{m-1})^n && \text{plugging in (1) and (2)} \\ n \ln(p_m) &\geq \ln(10) + n \ln(p_{m-1}) && \text{natural log both sides} \\ n \ln(p_m) - n \ln(p_{m-1}) &\geq \ln(10) \\ n(\ln(p_m) - \ln(p_{m-1})) &\geq \ln(10) && \text{factor out } n \\ n &\geq \frac{\ln(10)}{\ln(p_m) - \ln(p_{m-1})} \\ &\geq \frac{\ln(10)}{\ln(\frac{p_m}{p_{m-1}})} && \text{diff of logs} \end{aligned}$$

4. Meester 1.7.36

A pack contains m cards, labelled $1, 2, \dots, m$. The cards are dealt out in a random order, one by one. Given that the label of the k th card dealt is the largest of the first k cards, what is the probability that it is also the largest in the whole pack?

Let A be the event that the k th card is the largest of the first k cards. Let B be the event that the k th card is the largest of all M cards. We wish to calculate $\mathbb{P}(B|A)$. Per the the definition of conditional probability,

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(B \cap A)}{\mathbb{P}(A)} \quad (3)$$

Note that $\mathbb{P}(A) = 1/k$, since all of the first k cards are equally likely to have the highest value cards. For the same reason, $\mathbb{P}(B) = 1/M$.

Now, since the intersection operation is commutative,

$$\mathbb{P}(B \cap A) = \mathbb{P}(A \cap B) = \mathbb{P}(A|B)\mathbb{P}(B) = (1)\frac{1}{M} \quad (4)$$

Plugging (4) into (3), we see

$$\mathbb{P}(B|A) = \frac{\frac{1}{M}}{\frac{1}{k}} = \frac{k}{M}$$

5. Meester 1.7.38

Let A_1, A_2, \dots be events. Show that

$$\mathbb{P}\left(\bigcap_{i=1}^n A_i\right) \geq \sum_{i=1}^n \mathbb{P}(A_i) - (n-1)$$

Proof.

$$\begin{aligned}
 \mathbb{P}\left(\bigcap_{i=1}^n A_i\right) &= 1 - \mathbb{P}\left(\bigcap_{i=1}^n A_i\right)^c \\
 &= 1 - \mathbb{P}\left(\bigcup_{i=1}^n (A_i)^c\right) \\
 &\geq 1 - \sum_{i=1}^n \mathbb{P}(A_i)^c \\
 &= 1 - \sum_{i=1}^n (1 - \mathbb{P}(A_i)) \\
 &= 1 - n + \sum_{i=1}^n \mathbb{P}(A_i) \\
 &= \sum_{i=1}^n \mathbb{P}(A_i) - (n - 1)
 \end{aligned}$$

$$\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$$

DeMorgan's

Union probability bound

$$\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$$

Split summation terms. $\sum_{i=1}^n 1 = n$

□