

School of Mathematics, Thapar Institute of Engineering & Technology, Patiala

UMA007 : Numerical Analysis

Assignment 9

Initial-Value Problems for Ordinary Differential Equations

1. Consider the following initial-value problem

$$x' = t(x + t) - 2, \quad x(0) = 2.$$

Use the Euler method with stepsize $h = 0.2$ to compute $x(0.6)$.

2. Given the initial-value problem

$$y' = \frac{1}{t^2} - \frac{y}{t} - y^2, \quad 1 \leq t \leq 2, \quad y(1) = -1,$$

with exact solution $y(t) = -\frac{1}{t}$:

- (a) Use Euler's method with $h = 0.05$ to approximate the solution, and compare it with the actual values of y .
- (b) Use the answers generated in part (a) and linear interpolation to approximate the following values of y , and compare them to the actual values.
 - i. $y(1.052)$ ii. $y(1.555)$ iii. $y(1.978)$.

3. Solve the following IVP by second-order Runge-Kutta method

$$y' = -y + 2 \cos t, \quad y(0) = 1.$$

Compute $y(0.2)$, $y(0.4)$, and $y(0.6)$ with mesh length 0.2.

4. A projectile of mass $m = 0.11$ kg shot vertically upward with initial velocity $v(0) = 8$ m/s is slowed due to the force of gravity, $F_g = -mg$, and due to air resistance, $F_r = -kv|v|$, where $g = 9.8$ m/s² and $k = 0.002$ kg/m. The differential equation for the velocity v is given by

$$mv' = -mg - kv|v|.$$

- (a) Find the velocity after 0.1, 0.2, \dots , 1.0 s.
- (b) To the nearest tenth of a second, determine when the projectile reaches its maximum height and begins falling.

5. Using Runge-Kutta fourth-order method to solve the IVP at $x = 0.8$ for

$$\frac{dy}{dx} = \sqrt{x+y}, \quad y(0.4) = 0.41$$

with step length $h = 0.2$.

6. Water flows from an inverted conical tank with circular orifice at the rate

$$\frac{dx}{dt} = -0.6\pi r^2 \sqrt{2g} \frac{\sqrt{x}}{A(x)},$$

where r is the radius of the orifice, x is the height of the liquid level from the vertex of the cone, and $A(x)$ is the area of the cross section of the tank x units above the orifice. Suppose $r = 0.1$ ft, $g = 32.1$ ft/s², and the tank has an initial water level of 8 ft and initial volume of $512(\pi/3)$ ft³. Use the Runge-Kutta method of order four to find the following.

- (a) The water level after 10 min with $h = 20$ s.
- (b) When the tank will be empty, to within 1 min.

CONTINUED

7. The following system represent a much simplified model of nerve cells

$$\begin{aligned}\frac{dx}{dt} &= x + y - x^3, \quad x(0) = 0.5 \\ \frac{dy}{dt} &= -\frac{x}{2}, \quad y(0) = 0.1\end{aligned}$$

where $x(t)$ represents voltage across the boundary of nerve cell and $y(t)$ is the permeability of the cell wall at time t . Solve this system using Runge-Kutta fourth-order method to generate the profile up to $t = 0.2$ with step size 0.1.

8. Use Runge-Kutta method of order four to solve

$$y'' - 3y' + 2y = 6e^{-t}, \quad 0 \leq t \leq 1, \quad y(0) = y'(0) = 2$$

for $t = 0.2$ with stepsize 0.2.
