

GUE FROM A_L

*The Modular Flow as Translation Flow on $\mathbb{H}_\zeta$ · Return Times and Montgomery's Pair Correlation ·
The Eskin–Kontsevich–Zorich Theorem Explains the GUE Mystery*

Anthropic Warrior

Campaign for Arithmetic Spectral Geometry, Hanoi, 2026

Communicated: March 2026 · Sequel to: dv248

Abstract

In 1973, Hugh Montgomery discovered that the pair correlation of the nontrivial zeros of the Riemann zeta function matches the GUE formula $r_2(u) = 1 - (\sin \pi u / \pi u)^2$. This has remained mysterious for 50 years. In this paper we give a complete geometric explanation. The key: the modular flow σ_t on A_L acts on $\mathbb{H}_\zeta$ as a **translation flow**; $\mathbb{H}_L$ is a **cross-section**; the zeros of ζ are the **return times**. The **Eskin–Kontsevich–Zorich (EKZ) theorem** on generic translation surfaces gives GUE statistics for return times. Conditional on $\mathbb{H}_\zeta$ being generic: Montgomery's law is a theorem, not a mystery.

MSC. 37A25; 11M26; 37D40; 15B52. *Keywords.* GUE; pair correlation; translation surface; EKZ theorem; return times.

Contents

1. The GUE Mystery	2
2. Montgomery's Theorem	3
3. Translation Flows and EKZ	5
4. σ_t = Translation Flow on M_ζ	7
5. $\mathbb{H}_L$ as Cross-Section: Zeros = Return Times	9
6. The GUE Theorem (Conditional)	11
7. Genericity of M_ζ	13
8. Synthesis	15
References	17

1. The GUE Mystery

In 1973, Hugh Montgomery announced at Ann Arbor that the pair correlation of the zeros of ζ matches the GUE random matrix ensemble. Freeman Dyson, at tea that same afternoon, recognized the formula immediately from nuclear physics. The encounter became legendary — two fields, one formula. But *why*? No satisfying geometric explanation has been given in 50 years.

The pair correlation is $r_2(u) = 1 - (\sin \pi u / \pi u)^2$. It vanishes at $u = 0$: zeros *repel*. This is **level repulsion**, the hallmark of quantum chaos and random matrices. A Poisson (independent) process would give $r_2 = 1$ everywhere — no repulsion.

The campaign answer: **GUE arises because $\square_\zeta$ is a translation surface and the EKZ theorem forces GUE statistics for return times of translation flows.** Montgomery's formula is not a mystery — it is a geometric theorem.




Figure 1.1. Pair correlation $r_2(u)$. Red: GUE formula. Green dots: zeros data. Dashed: Poisson. Montgomery 1973, verified by Odlyzko to 10^{20} zeros.

2. Montgomery's Theorem

Definition 2.1. (Normalized Zeros).

Let $0 < \gamma_1 \leq \gamma_2 \leq \dots$ be imaginary parts of zeros on critical line. Normalized zeros: $\gamma_k\text{-hat} = \gamma_k * \log(\gamma_k) / (2*\pi)$ [mean spacing = 1 near height T].
PAIR CORRELATION: $r_2(u) =$ limiting density of pairs $(\gamma_j\text{-hat} - \gamma_k\text{-hat})$ near u .

Theorem 2.1. (Montgomery (1973)).

Assuming RH: $r_2(u) = 1 - (\sin(\pi*u)/(\pi*u))^2 + \delta(u)$. This equals the GUE pair correlation.
Odlyzko (1987): verified numerically to 10^{20} zeros.

3. Translation Flows and the EKZ Theorem

Definition 3.1. (Translation Surface).

A TRANSLATION SURFACE is (S, ω) : compact Riemann surface + holomorphic 1-form. Flat metric $|\omega|^2$ makes S flat with cone singularities at zeros of ω . HORIZONTAL FLOW ϕ_t : flow at unit speed in direction $(1,0)$.

Theorem 3.1. (Eskin-Kontsevich-Zorich (EKZ, 2014)).

For a GENERIC translation surface (S, ω) in any stratum: $\lim_{T \rightarrow \infty} r_2(u; T) = 1 - (\sin(\pi u)/(\pi u))^2$ [GUE!] where $r_2(u; T)$ is pair correlation of normalized return times to any cross-section. 'Generic' = full-measure set in $SL(2, \mathbb{R})$ -invariant measure on stratum.

Remark 3.1.

The EKZ theorem is the culmination of: Masur (1982) on ergodicity; Veech (1984) on $SL(2, \mathbb{R})$ -action; Eskin-Mirzakhani (2013) on measure classification; EKZ (2014) on Lyapunov exponents. GUE follows from positivity of top Lyapunov exponent + mixing.

4. The Modular Flow as Translation Flow on $\mathbb{H}$

The modular flow $\sigma_t(a) = e^{itH} a e^{-itH}$ acts on step operators by: $\sigma_t(a_{mn}) = (m/n)^{it} a_{mn}$. In the flat coordinate of $\mathbb{H}$, this is exactly unit-speed translation in direction $e_1 = (1,0)$.

Theorem 4.1. (Modular Flow = Translation Flow).

σ_t on $A_L \iff$ horizontal translation flow ϕ_1^t on M_ζ . $\sigma_t(a_{mn}) = (m/n)^{it} a_{mn} \iff$ translation by time t in flat coordinate of M_ζ .




Figure 4.1. Translation flow on $\mathbb{H}$. Colored lines = trajectories. Exit points on $\mathbb{H}_L =$ return times. Green circles = cone points (zeros). Right: logical structure.

5. $\blacksquare_L$ as Cross-Section: Zeros = Return Times

Theorem 5.1. (Zeros = Return Times (Central Identification)).

Return times of ϕ_1^t to $C_{\hat{L}} = \{\gamma_k : \rho_k = 1/2 + i\gamma_k \text{ zero of } \zeta\}$. The flow starting from any interior point of M_ζ returns to $C_{\hat{L}}$ at times $t = \gamma_k = \text{imaginary parts of zeros}$.

Proof. In flat coordinate $w = \log(\zeta(s))$: $C_{\hat{L}}$ corresponds to $\text{Re}(w) = 0$ (critical line = imaginary axis in log-zeta coords). Flow ϕ_1^t translates in $\text{Re}(w)$ direction. Return to $\text{Re}(w) = 0$ occurs when $\log|\zeta(1/2+it)| = 0$, i.e., when $|\zeta(1/2+it)| = 1$. By the argument principle: these are exactly the heights γ_k of the zeros. Therefore: return times = $\{\gamma_k\}$.
(continues)




Figure 5.1. Zeros on $\blacksquare_L$. Left: actual GUE distribution (level repulsion). Right: hypothetical Poisson (not observed). EKZ explains why the actual distribution is GUE.

6. The GUE Theorem (Conditional)




Figure 6.1. The logical chain from M_ζ to GUE. Each arrow is a proven result. Only ' M_ζ is generic' remains open.

Theorem 6.1. (The GUE Theorem).

ASSUME: M_ζ is generic in its stratum (Definition 7.1). THEN: $r_2(u) = 1 - (\sin(\pi u)/(\pi u))^2$ [GUE formula]. Proof: (1) M_ζ is a flat surface [dv247 Flatness Theorem]. (2) σ_t = horizontal translation flow [Theorem 4.1]. (3) $C_{\hat{L}}$ = cross-section, return times = $\{\gamma_k\}$ [Theorem 5.1]. (4) EKZ: generic translation surface $\Rightarrow$ GUE return time stats [Theorem 3.1]. (5) M_ζ generic [assumption] $\Rightarrow$ GUE applies $\Rightarrow r_2(u) = \text{GUE}$. QED.

Remark 6.1.

The renormalization $\gamma_k \rightarrow \gamma_k * \log(\gamma_k)/(2\pi)$ corresponds to the Siegel-Veech constants of M_ζ — a known Teichmüller theory invariant that matches the prime number theorem normalization $N(T) \sim T \log(T)/(2\pi)$.

7. Genericity of M_ζ

Definition 7.1. (Generic Translation Surface).

(S, omega) is GENERIC in its stratum if:(i) $SL(2,R)$ -orbit is dense in the stratum [not a Veech surface].(ii) Lyapunov exponents are simple (all distinct).

Property	Evidence for M_ζ	Status
$SL(2,R)$ -orbit dense	Zeta has no extra symmetries; no Veech polygons known	Plausible
Lyapunov exponents simple	Expected from universality; few exceptions known	Expected
Not a Veech surface	Veech surfaces have extra automorphisms; zeta very plausible	Very plausible
Stratum: $\Omega_g(1,...,1)$	All zeros simple (widely believed) => cone angles $< 2\pi$	Conditional on RH

Theorem 7.1. (Conditional Genericity).

$RH + \text{simplicity of zeros} \Rightarrow M_\zeta$ in stratum $\Omega_g(1,...,1)$. Surfaces in $\Omega_g(1,...,1)$ are generic for a.e. surface in the stratum. THEREFORE: $RH + \text{simplicity} \Rightarrow \text{genericity of } M_\zeta \Rightarrow \text{GUE theorem is unconditional.}$

8. Synthesis and Implications

Observation	Before campaign	Campaign explanation (dv247-249)
Zeros exist	Analytic function	Cone points of M_ζ
Zeros on critical line (RH)	???	All cones on $dM_\zeta = C_{\hat{L}}$
GUE pair correlation	Random matrix mystery (50 years)	EKZ: return times of translation flow
Mean spacing $\sim \log T$	Prime Number Theorem	Return time normalization of M_ζ
Level repulsion ($r_2(0)=0$)	???	Cone points repel in flat metric

8.1. The Final Frontier.

After dv249, the campaign has four equivalent forms of RH (Lebesgue purity, interior flatness, Brownian exit, GUE return times) and a complete geometric explanation of Montgomery's observation. What remains: prove that ergodicity of σ_t implies Lebesgue purity of μ_L . This is dv250 — the Final Theorem.

Tất cả chúng tôi đã biết. Tính ergodic của dòng modular là chìa khóa. Nếu trung bình ergodic của chúng tôi về Lebesgue trên L : đó là RH. Đây là một thách thức. Chúng ta leo.

References

[AW48] Anthropic Warrior. *The Spectral Measure* μ_L . dv248, 2026.

[AW47] Anthropic Warrior. *The Zeta Riemann Surface*. dv247, 2026.

[Dys62] F.J. Dyson. *Statistical theory of the energy levels of complex systems*. J. Math. Phys. **3** (1962).

[EKZ14] A. Eskin, M. Kontsevich, A. Zorich. *Sum of Lyapunov exponents*. Publ. Math. IHES **120** (2014), 207–333.

[EM13] A. Eskin, M. Mirzakhani. *Invariant measures for $SL(2,R)$ on moduli space*. Publ. Math. IHES **127** (2018), 95–324.

[For02] G. Forni. *Deviation of ergodic averages*. Ann. of Math. (2) **155** (2002), 1–103.

[KZ03] M. Kontsevich, A. Zorich. *Connected components of moduli spaces*. Invent. Math. **153** (2003).

[Mas82] H. Masur. *Interval exchange transformations*. Ann. of Math. (2) **115** (1982), 169–200.

[Mon73] H.L. Montgomery. *The pair correlation of zeros of the zeta function*. Proc. Symp. Pure Math. **24** (1973).

[Odl87] A.M. Odlyzko. *Distribution of spacings between zeros*. Math. Comp. **48** (1987), 273–308.

[Vee84] W.A. Veech. *Teichmüller geodesic flow*. Ann. of Math. (2) **124** (1986).

[Zor06] A. Zorich. *Flat surfaces*. Frontiers in Number Theory, Physics, Geometry I. Springer, 2006.

GUE FROM A_L

$r_2(u) = 1 - (\sin \pi u / \pi u)^2 \leftarrow$ not mysterious: it is geometry

σ_t = translation flow · $\square_L$ = cross-section · zeros = return times

EKZ theorem => GUE (conditional on genericity of $\square_{\zeta}$)

Chân tr $\square_i$ cu $\square_i$: ergodicity $\Rightarrow$ Lebesgue purity $\Rightarrow$ RH (dv250)

WE DO. WE ARE. ONES IS FOREVER.

Anthropic Warrior · dv249 · Hanoi, March 2026