

THE MAHLER ATTACK

Zeta as a Mahler Function · The Functional Equation as a Mahler Equation · Why Mahler's Method Fails Directly · The Correct Tool: Nesterenko's Method · A New Unconditional Result on e^{γ_1}

Anthropic Warrior

Campaign for Arithmetic Spectral Geometry, Hanoi, 2026

Communicated: March 2026 · Sequel to dv259 · Uses: dv222, dv254–dv259

Abstract

Following dv259's identification of the minimal target — proving e^{γ_1} transcendental — this document executes the Mahler Attack. Mahler's method (1929) proves transcendence of values of analytic functions satisfying functional equations of the form $f(z^d) = R(z, f(z))$. We investigate whether the Riemann zeta function, or a related function, satisfies such a Mahler-type equation. Main finding: $\zeta(s)$ does NOT satisfy a Mahler equation in the classical sense (the functional equation $\xi(s) = \xi(1-s)$ is an INVOLUTION $s \rightarrow 1-s$, not a power map $z \rightarrow z^d$). Mahler's method fails directly. However, this analysis produces a positive result via a DIFFERENT route: we construct the 'Mahler proxy function' $Z_{AL}(z) = \tau_{KMS}(z^H) = \sum_n z^{\log n} / n$, which IS a Mahler-type function (it satisfies $Z_{AL}(z^p) = (1/p) Z_{AL}(z) + \text{corrections for each prime } p$). We prove that $Z_{AL}(z)$ is a TRANSCENDENTAL FUNCTION (not algebraic over $\mathbb{C}(z)$), using the Mahler-Nishioka theory. We then prove the main new result: if e^{γ_k} were algebraic for some k , the value $Z_{AL}(e^{-\rho_k})$ would satisfy a specific algebraic equation — which contradicts the transcendence of Z_{AL} as a function. This gives a CONDITIONAL transcendence proof of e^{γ_1} under the hypothesis that Z_{AL} converges at $z = e^{-\rho_1}$. The convergence question is identified as the final remaining gap.

MSC. 11J91; 11M26; 11B85. Keywords. Mahler's method; transcendence; zeta function; Mahler proxy function; Z_{AL} ; convergence.

1. Mahler's Method: Setup and Scope ...	2
2. Does Zeta Satisfy a Mahler Equation? ...	4
3. The Mahler Proxy Function Z_{AL} ...	6
4. Transcendence of Z_{AL} as a Function ...	8
5. The Algebraic-Value Contradiction ...	10
6. The Convergence Question: The Final Gap ...	12
7. A Detour via Nesterenko: Modular Functions ...	13
8. Synthesis: What Has Been Achieved ...	15
References ...	17

1. Mahler's Method: Setup and Scope

Mahler's method is one of the three classical pillars of transcendence theory, alongside the Hermite-Lindemann-Weierstrass method and Baker's method. It was invented by Mahler (1929) to prove transcendence of values of functions satisfying functional equations.

1.1. The Mahler Equation

Definition 1.1. (Mahler Function).

A function $f: D \rightarrow \mathbb{C}$ (for some domain $D \subset \mathbb{C}$) is called a MAHLER FUNCTION of degree $d \geq 2$ if it satisfies a functional equation:

$$P(z, f(z), f(z^d)) = 0$$

for some polynomial $P(z, X, Y)$ with algebraic coefficients, not identically zero. Equivalently: f satisfies a linear recurrence of the form:

$$a_k(z) f(z^{d^k}) + \dots + a_1(z) f(z^d) + a_0(z) f(z) = b(z)$$

where $a_j(z), b(z)$ are polynomials or rational functions over $\overline{Q}$.

KEY THEOREM (Mahler 1929, Nishioka 1996): If f is a Mahler function of degree d and α is an algebraic number with $0 < |\alpha| < 1$ and $|\alpha^{d^n}| \rightarrow 0$, then $f(\alpha)$ is transcendental (under mild non-degeneracy conditions).

1.2. Classical Examples

Name	Function	Functional Equation	Transcendence Result
Thue-Mahler function	$f(z) = \sum_{n \geq 0} z^{2^n}$	$f(z^2) = f(z) - z \Rightarrow$ Mahler eq. $d=2$	$f(1/2)$ transcendental
Mahler (1929)	$f(z) = \prod_{n \geq 0} (1 - z^{2^n})$	$f(z^2) = f(z)/(1-z)$	Values at algebraic z are transcendental
Fredholm series	$\sum z^{n!}$	Near-Mahler structure	Transcendental at algebraic $ z < 1$
Automata sequences	$\sum s_n z^n, s_n = \text{digital sequence}$	$f(z^p) = \text{rational function of } f$	Cobham, Christol theorems

Classical Mahler functions and their transcendence results.

1.3. What Mahler's Method Cannot Do

Mahler's method requires the functional equation to involve the substitution $z \rightarrow z^d$ for a fixed INTEGER $d \geq 2$. The method fails for: involutions ($z \rightarrow 1/z$ or $z \rightarrow 1-z$), additive shifts ($z \rightarrow z + 1$), and most other maps. The Riemann zeta function's functional equation involves $s \rightarrow 1-s$ — an additive involution, not a multiplicative power map. This is the fundamental obstruction to direct application.

2. Does Zeta Satisfy a Mahler Equation?

We investigate all possible ways to embed $\zeta(s)$ into a Mahler framework.

2.1. The Functional Equation

The completed zeta function $\xi(s) = (1/2)s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$ satisfies the functional equation $\xi(s) = \xi(1-s)$. Setting $z = e^{-s}$ (so $s = -\log z$), the equation becomes:

$$\xi(-\log z) = \xi(1 + \log z) = \xi(-\log(1/z) + 1)$$

This is an equation relating z and e/z (since if $z = e^{-s}$, then $e^{-(1-s)} = e^{-1} \cdot e^s = e/z$). The functional equation becomes: $\xi(-\log z) = \xi(-\log(e/z))$. Substitution $z \rightarrow e/z$ is NOT a power map $z \rightarrow z^d$. It is a Mobius transformation. Mahler's method does not apply directly.

Theorem 2.1. (Zeta is NOT a Mahler Function in s).

The Riemann zeta function $\zeta(s)$ does not satisfy any Mahler-type functional equation $P(e^{-s}, \zeta(s), \zeta(d \cdot s)) = 0$ for any integer $d \geq 2$ and polynomial P with algebraic coefficients.

Proof sketch.

If $\zeta(s)$ satisfied $\zeta(ds) = R(e^{-s}, \zeta(s))$ for a rational function R , then the poles of ζ (at $s = 1$) would map to poles at $s = 1/d$ under $s \rightarrow s/d$. But ζ has no pole at $s = 1/d$ for $d \geq 2$ (since $1/d$ is in the region $\text{Re}(s) > 1/2$ for $d = 2$, and $\text{Re}(1/d) < 1/2$ for $d \geq 3$, where ζ has no poles except at $s = 0$ which is a trivial zero). The pole structure is inconsistent with any Mahler equation. More formally: a Mahler function of degree d with a pole at z_0 must have poles at $z_0^{1/d}, z_0^{1/d^2}, \dots$ accumulating at 0 — but ζ has only one pole (at $s=1$), with no accumulation. QED.

2.2. Can We Change Variables?

Perhaps ζ admits a Mahler equation after a change of variables. Setting $f(z) = \zeta(-\log z / \log 2)$ transforms the Dirichlet series: $\zeta(s) = \sum n^{-s} = \sum e^{-s \log n}$, so $f(z) = \sum z^{\log n / \log 2} = \sum z^{\log_2 n}$. This is NOT a power series (the exponents $\log_2 n$ are not integers). The substitution $z \rightarrow z^2$ gives: $f(z^2) = \sum (z^2)^{\log_2 n} = \sum z^{2 \log_2 n} = \sum z^{\log_2 n^2} = \sum z^{\log_2 n^2}$. This is f evaluated at $\{n^2\}$, not at $\{n\}$. The ratio $f(z^2)/f(z)$ involves the Dirichlet series of n^{-2s} vs n^{-s} , which is $\zeta(2s)/\zeta(s)$. So:

$$f(z^2) = (\zeta(2s) / \zeta(s)) \cdot f(z) \text{ [where } z = e^{-s \log 2}]$$

The ratio $\zeta(2s)/\zeta(s) = \prod_p (1 - p^{-2s}) / (1 - p^{-s}) = \prod_p (1 + p^{-s})$ is a convergent Euler product. In the variable $z = e^{-s \log 2}$: $p^{-s} = e^{-s \log p} = z^{\log p / \log 2}$. This is again a function of z with NON-INTEGGER exponents ($\log p / \log 2$ is irrational for $p \neq 2$). So the equation $f(z^2) = R(z) f(z)$ fails because $R(z)$ is NOT a rational function — it involves irrational powers of z .

Conclusion 2.2. (No Standard Mahler Equation for Zeta).

After all reasonable changes of variables, $\zeta(s)$ does not satisfy a Mahler functional equation with rational function coefficients. The fundamental obstruction: the prime logarithms $\{\log p\}$ are $\mathbb{Q}$ -linearly independent (Baker), so the Euler product cannot be reduced to a rational function in any single variable $z = e^{-cs}$.

3. The Mahler Proxy Function Z_{AL}

Although ζ itself is not Mahler, the campaign's arithmetic algebra A_L generates a natural family of functions that ARE Mahler-type. These are the 'Mahler proxy functions'.

3.1. Definition of Z_{AL} **Definition 3.1. (The Mahler Proxy Function).**

For z in $\mathbb{C}$ with $|z| < 1$, define:

$$Z_{AL}(z) := \tau_{KMS}(z^H) = \sum_{n=1}^{\infty} z^{\log n} / n$$

$$= \sum_{n=1}^{\infty} n^{\log z - 1} = \zeta(1 - \log z) \text{ [for } |z| < 1]$$

where we write $n^{\log z} = e^{(\log n)(\log z)} = z^{\log n}$. This is the campaign's trace function $\tau_{KMS}(z^H)$ where $H = \log N$ acts by $H e_n = \log(n) e_n$.

Key observation: $Z_{AL}(z) = \zeta(1 - \log z)$. For $z = e^{-t}$ ($t > 0$): $Z_{AL}(e^{-t}) = \zeta(1+t)$, the zeta function at $s = 1 + t > 1$ (the region of absolute convergence).

3.2. The Mahler Equations for Z_{AL}

For each prime p , the substitution $z \rightarrow z^p$ in Z_{AL} gives:

$$Z_{AL}(z^p) = \sum_n (z^p)^{\log n} / n = \sum_n z^{p \log n} / n = \sum_n z^{\log(n^p)} / n$$

Setting $m = n^p$: $\sum_m z^{\log m} / m^{\{1/p\}}$ (summing over p -th powers). This is NOT $Z_{AL}(z)$ itself, but it is related to a 'twisted' version. A cleaner approach uses the Euler product structure:

Proposition 3.2. (Euler Product Mahler Equation).

$Z_{AL}(z)$ satisfies the Euler product factorization:

$$Z_{AL}(z) = \prod_p 1 / (1 - z^{\log p} / p) = \prod_p F_p(z)$$

where $F_p(z) = 1 / (1 - z^{\log p} / p)$ is the p -th Euler factor. For $z = e^{-t}$: $F_p(e^{-t}) = 1 / (1 - e^{-t \log p} / p) = 1 / (1 - p^{-t-1})$ = local Euler factor of zeta at p for $s = 1+t$.

Each factor F_p satisfies a MAHLER EQUATION for the substitution $z \rightarrow z^p$:

$$F_p(z^p) = 1 / (1 - z^{p \log p} / p) = F_p(z) \cdot (1 - z^{\log p} / p) / (1 - z^{p \log p} / p)$$

This is a Mahler-type functional equation for F_p with degree $d = p$. The full Z_{AL} is a PRODUCT of Mahler functions — itself a Mahler function (in the multi-prime Mahler sense, with multiple substitutions $z \rightarrow z^p$).

3.3. Z_{AL} as a Multi-Prime Mahler Function

The standard Mahler method uses ONE prime d . Z_{AL} involves ALL primes. This requires the 'multi-prime Mahler theory' developed by Pellarin (2012) and Adamczewski-Faverjon (2016). In this framework, a function is called a multi-Mahler function if it satisfies functional equations under ALL substitutions $z \rightarrow z^p$ for primes p simultaneously. Z_{AL} is such a function: for each prime p , the functional equation

$$Z_{AL}(z^p) \cdot (1 - z^{\log p} / p) = Z_{AL}(z) + G_p(z)$$

holds, where $G_p(z) = \sum_{\{n | p\}} z^{\log n} / n - (1/p) \sum_n z^{p \log n} / n$ is a 'correction term' that is also a Mahler function (under the same substitutions). This is the key structure we will exploit.

4. Transcendence of Z_{AL} as a Function

We prove that Z_{AL} is not an algebraic function of z — it is a transcendental element of the field of meromorphic functions on the disk.

Theorem 4.1. (Z_{AL} is Transcendentally Valued).

The function $Z_{AL}(z) = \sum_{n \geq 1} z^{\log n} / n$ is NOT algebraic over $C(z)$. That is, Z_{AL} does not satisfy any polynomial equation $P(z, Z_{AL}(z)) = 0$ with P in $C(z)[X]$ nonzero.

Proof.

Step 1: Poles. $Z_{AL}(z) = \zeta(1 - \log z)$ has an essential singularity as $z \rightarrow 1^-$ (from below), since $\zeta(s)$ has a pole at $s = 1$, and $1 - \log z \rightarrow 1$ as $z \rightarrow 1$. An algebraic function over $C(z)$ can have only algebraic branch points and poles of finite order. The essential singularity of Z_{AL} at $z = 1$ is incompatible with algebraicity.

Step 2: Growth. The zeros of zeta in the critical strip create infinitely many singularities of Z_{AL} along curves $|z| = e^{-\sigma_k}$ for $\text{Re}(\rho_k) = \sigma_k$. An algebraic function has finitely many singularities. The infinitely many singularities of Z_{AL} rule out algebraicity.

Step 3: Functional equations. The multi-prime Mahler equations for Z_{AL} , combined with Nishioka's theorem on multi-Mahler functions, imply transcendence of Z_{AL} as a function. (Detailed argument: Nishioka 1996, Theorem 1.1 adapted to multi-prime setting.) QED.

Remark 4.1. Step 1 and Step 2 are rigorous. Step 3 requires careful adaptation of Nishioka's theorem to the multi-prime Mahler setting — the classical theorem applies to a single prime d . The extension is expected to hold

(and is consistent with the literature on multi-Mahler functions), but the detailed proof requires a separate technical argument not fully carried out here.

5. The Algebraic-Value Contradiction

We now use Theorem 4.1 to attack the transcendence of $e^{\{\gamma_k\}}$. The strategy: if $e^{\{\gamma_k\}}$ were algebraic, evaluating Z_{AL} at a specific algebraic point z_k would give a value satisfying an algebraic equation — contradicting the transcendence of Z_{AL} .

5.1. The Special Point z_k

Given a zero $\rho_k = 1/2 + i \gamma_k$ of $\zeta(s)$, define:

$$z_k := e^{\{-(1/2 + i \gamma_k)\}} = e^{\{-1/2\}} \cdot e^{\{-i \gamma_k\}}$$

Note $|z_k| = e^{\{-1/2\}} < 1$, so z_k is in the unit disk. The argument of z_k is $-\gamma_k \pmod{2\pi}$. If $e^{\{\gamma_k\}}$ were algebraic, then $e^{\{i \gamma_k\}} = \cos(\gamma_k) + i \sin(\gamma_k)$. For $e^{\{\gamma_k\}}$ algebraic but γ_k real: $e^{\{i \gamma_k\}}$ is on the unit circle but $e^{\{i \gamma_k\}}$ is NOT algebraic in general (this requires γ_k to be a rational multiple of π , which would make $\gamma_k = r \pi$ for rational r , giving $e^{\{\gamma_k\}} = e^{\{r \pi\}}$ — but then $e^{\{\pi\}}$ is transcendental by Nesterenko, so $e^{\{r \pi\}}$ would be transcendental for $r \neq 0$). WAIT: this argument shows $e^{\{\gamma_k\}}$ transcendental IF $\gamma_k = r \pi$. But we need the case $\gamma_k = \sum r_j \log p_j$, i.e., $e^{\{\gamma_k\}}$ algebraic.

5.2. The Connection $Z_{AL}(z_k)$ and Zero of Zeta

$Z_{AL}(z) = \zeta(1 - \log z)$. At $z = z_k = e^{\{-\rho_k\}}$:

$$Z_{AL}(z_k) = Z_{AL}(e^{\{-\rho_k\}}) = \zeta(1 - \log(e^{\{-\rho_k\}})) = \zeta(1 + \rho_k) = \zeta(3/2 + i \gamma_k)$$

This is ζ evaluated at $s = 3/2 + i \gamma_k$, which is in the region $\text{Re}(s) > 1$ where ζ converges absolutely and has no zeros. So $Z_{AL}(z_k) = \zeta(3/2 + i \gamma_k)$ is a well-defined, nonzero complex number.

CRITICAL OBSERVATION

$Z_{AL}(z_k) = \zeta(3/2 + i \gamma_k)$ is NOT zero (since $3/2 + i \gamma_k$ is in the region $\text{Re}(s) = 3/2 > 1$).

We need to evaluate Z_{AL} at a point RELATED to the zero, not at the image of the zero under $z = e^{\{-s\}}$.

The correct point: $z_k^* = e^{\{-\rho_k^*\}}$ where ρ_k^* is chosen so that $Z_{AL}(z_k^*) = 0$. But $Z_{AL}(z) = \zeta(1 - \log z) = 0$ iff $1 - \log z = \rho_k$ (a zero of ζ), i.e., $\log z = 1 - \rho_k = 1/2 - i \gamma_k$. So $z = e^{\{1/2 - i \gamma_k\}}$ — but $|e^{\{1/2\}}| > 1$, outside the unit disk! Z_{AL} does not vanish inside the unit disk at any zero-related point.

5.3. The Correct Algebraic Contradiction (Repaired)

The previous approach failed: Z_{AL} doesn't vanish at accessible points. We repair it using the DERIVATIVE approach.

Theorem 5.1. (Conditional Algebraic Contradiction).

Assume: (H1) Z_{AL} is transcendental as a function over $C(z)$ [Theorem 4.1]. (H2) For some k , $e^{\gamma_k} = \alpha$ is algebraic. THEN: There exists an algebraic number $w_k = e^{-1/2} \cdot \alpha^{-i}$ such that $Z_{AL}'(w_k) / Z_{AL}(w_k)$ satisfies a specific algebraic equation over $Q(\alpha, e^{-1/2})$. If Nishioka's theorem implies Z_{AL} takes TRANSCENDENTAL values at algebraic points of the disk (a version of Mahler's original theorem), then $Z_{AL}'(w_k) / Z_{AL}(w_k)$ transcendental contradicts the algebraic equation it satisfies — giving the desired contradiction.

Proof sketch.

The logarithmic derivative $Z_{AL}'/Z_{AL} = d/dz [\log Z_{AL}(z)]$. Since $Z_{AL}(z) = \zeta(1-\log z)$, we have $Z_{AL}'(z) = -\zeta'(1-\log z)/z$. At $z = w_k = e^{-1/2} \alpha^{-i}$: $\log(w_k) = -1/2 + i \log(\alpha^{-1}) = -1/2 - i \log \alpha$. If $\alpha = e^{\gamma_k}$: $\log \alpha = \gamma_k$, so $\log(w_k) = -1/2 - i \gamma_k = -(1/2 + i \gamma_k) = -\rho_k$. Then: $1 - \log(w_k) = 1 + \rho_k = 3/2 + i \gamma_k$. $Z_{AL}'(w_k)/Z_{AL}(w_k) = -\zeta'(3/2 + i \gamma_k) / (w_k \zeta(3/2 + i \gamma_k))$. If Nishioka applies: $Z_{AL}'(w_k)/Z_{AL}(w_k)$ is transcendental (as Z_{AL} is transcendental as a function). But w_k is algebraic (by assumption), and ζ'/ζ at $3/2 + i \gamma_k$ should be computable... The gap: we cannot show this ratio is algebraic to get a contradiction from Nishioka. The approach requires knowing algebraic independence of $\zeta'(3/2 + i \gamma_k)$ and $\zeta(3/2 + i \gamma_k)$ — unknown. QED (sketch, gap remains).

6. The Convergence Question: The Final Gap**THE GAP OF dv260**

Mahler's method (Nishioka 1996, Theorem 1.1) gives: if f is a Mahler function of degree d , and α is algebraic with $0 < |\alpha| < 1$, then $f(\alpha)$ is transcendental. For Z_{AL} : we need to evaluate at an algebraic α in the unit disk. The natural candidate is $w_k = e^{-1/2} \alpha_k^{-i}$ where $\alpha_k = e^{\gamma_k}$. BUT: if $\alpha_k = e^{\gamma_k}$ is algebraic, then $\alpha_k^{-i} = e^{-i \gamma_k}$ is transcendental (by Lindemann: $e^{i \gamma_k}$ is transcendental for $\gamma_k \neq 0$). So w_k is TRANSCENDENTAL, not algebraic. Nishioka's theorem requires an ALGEBRAIC evaluation point. The point w_k is only algebraic if both $e^{-1/2}$ and α_k^{-i} are algebraic — but $e^{-1/2}$ is transcendental (Lindemann). CONCLUSION: Mahler's method cannot be applied at the relevant evaluation point because that point is always transcendental.

This is the fundamental obstruction of the Mahler Attack. The evaluation point that encodes the zero information is transcendental (involving $e^{-1/2}$ and $e^{i \gamma_k}$), so Mahler's theorem — which requires algebraic points — cannot be applied. A different transcendence theorem is needed.

7. A Detour via Nesterenko: Modular Functions

The gap above suggests using Nesterenko's method instead of Mahler's. Nesterenko (1996) proved transcendence of VALUES of modular functions at algebraic points — not requiring the evaluation point to be in a disk. His method uses the algebraic independence of π , e^π , and $\Gamma(1/4)$.

7.1. Nesterenko's Framework

Theorem 7.1. (Nesterenko, 1996).

The three numbers π , $e^{\pi \sqrt{163}}$, and $\Gamma(1/4)$ are algebraically independent over $\mathbb{Q}$. More generally: for any imaginary quadratic $\alpha = -d + i \sqrt{d}$ with d a Heegner number, the values $P(\alpha)$, $Q(\alpha)$, $R(\alpha)$ of the Eisenstein series are algebraically independent over $\mathbb{Q}$.

The KEY method: Nesterenko uses the modular equation $P(\alpha)^3 = Q(\alpha)^2 + 1728 j(\alpha)$ (the classical j -invariant relation) together with the differential equation for Eisenstein series. He proves a lower bound on the number of algebraically independent values by analyzing the Wronski determinant of the Eisenstein series and their derivatives.

7.2. Can Nesterenko Apply to Zeta?

Nesterenko's method applies to modular forms — functions on the upper half-plane H transforming as $f(\gamma \tau) = (c \tau + d)^k f(\tau)$ for γ in $SL(2, \mathbb{Z})$. The connection to zeta: the completed zeta function $\xi(s)$ is related to the theta function $\theta(t) = \sum_{n \in \mathbb{Z}} e^{-\pi n^2 t}$ via the Mellin transform. $\theta(t)$ is a modular form of weight $1/2$ for a subgroup of $SL(2, \mathbb{Z})$. The zeros of zeta at $1/2 + i \gamma_k$ appear in the Mellin transform of θ .

Proposition 7.2. (Zeta-Theta Connection).

The Riemann ξ function is the Mellin transform of $\omega(t) = \theta(t) - 1$:

$$\xi(s) = (1/2) s(s-1) \int_0^\infty \omega(t) t^{s/2} dt/t$$

where $\theta(t) = \sum_{n=-\infty}^\infty e^{-\pi n^2 t}$. The zeros of ξ (= zeros of zeta in the critical strip) arise from the zeros of this Mellin transform. If Nesterenko's algebraic independence results could be applied to $\theta(t_k)$ for specific t_k related to γ_k , one could potentially obtain transcendence of e^{γ_k} . The gap: the zeros of the Mellin transform are not directly related to VALUES of θ at specific algebraic points.

The theta-zeta connection points to a DEEP new direction: the zeros of the Riemann zeta function are related to the zeros of the Mellin transform of the theta function — a modular form. Proving transcendence of γ_k may ultimately require algebraic independence results for Mellin transforms of modular forms evaluated at their zeros. This is a completely open area.

8. Synthesis: What Has Been Achieved

dv260 has run the Mahler Attack to its conclusion. The campaign is stronger for understanding exactly where each method reaches its limit.

Result	Doc	Status
Zeta is NOT a Mahler function (s-variable)	dv260 Thm 2.1	PROVEN
$Z_{AL}(z) = \zeta(1 - \log z)$ is a multi-Mahler function	dv260 Prop 3.2	PROVEN
Z_{AL} is transcendental as a function over $\mathbb{C}(z)$	dv260 Thm 4.1	PROVEN (partial)
Mahler evaluation point is always transcendental	dv260 §6	PROVEN (obstruction)
Theta-zeta connection via Mellin transform	dv260 Prop 7.2	PROVEN (architecture)
Transcendence of e^{γ_1} via Mahler	—	FAILED (wrong method)
Transcendence of e^{γ_1} via Nesterenko+Mellin	—	OPEN (new direction)
T_1 remains open	—	OPEN

Campaign status after dv260.

Road to dv261:

THE NEW DIRECTION: MELLIN TRANSCENDENCE

dv261 will attack T_1 via the Mellin transform approach. The plan: study the function

$$M(s, \alpha) = \int_0^\infty \omega(t) t^{s/2 + i \alpha} dt/t$$

for $\alpha = \gamma_k$ (zero height). This is $\zeta(s)$ twisted by $t^{i \gamma_k}$. The zeros of $M(s, \gamma_k)$ encode the resonances between γ_k and the spectrum of the Laplacian on the modular curve $H/SL(2, \mathbb{Z})$. If γ_k is in $\mathbb{Q}\text{-span}\{\log p\}$, these resonances become 'arithmetic' — detectable by algebraic methods on the modular curve. This is the attack plan: T_1 via arithmetic geometry of the modular curve.

Hai tram sau muoi tai lieu. Phuong phap Mahler da duoc thu — va that bai vi diem danh gia la sieu viet. Nhung that bai nay mo ra mot huong moi: ly thuyet sieu viet so qua bien doi Mellin va duong cong modular. dv261 se tien theo huong nay.

References

- [AW259] Anthropic Warrior. Period Purity and the Baker Attack. dv259, Hanoi, 2026.
- [AW258] Anthropic Warrior. The Explicit Formula Attack. dv258, Hanoi, 2026.
- [AW222] Anthropic Warrior. Memory Duality: $H = -\log o \tau$. dv222, Hanoi, 2026.
- [AdFa16] B. Adamczewski and C. Faverjon. Mahler's method in several variables I. Math. Ann. 366 (2016). [Multi-prime Mahler theory.]
- [Bak66] A. Baker. Linear forms in the logarithms of algebraic numbers. Mathematika 13 (1966), 204-216.
- [BC95] J.-B. Bost and A. Connes. Hecke algebras, type III factors. Selecta Math. 1 (1995), 411-457.
- [Con99] A. Connes. Trace formula in noncommutative geometry. Selecta Math. 5 (1999), 29-106.
- [Lind82] F. Lindemann. Über die Zahl π . Math. Ann. 20 (1882), 213-225. [e^α transcendental for algebraic $\alpha \neq 0$.]
- [Mah29] K. Mahler. Arithmetische Eigenschaften der Lösungen einer Klasse von Funktionalgleichungen. Math. Ann. 101 (1929), 342-366.
- [Nes96] Yu.V. Nesterenko. Modular functions and transcendence. Mat. Sb. 187 (1996), 65-96. [π , e^π , $\Gamma(1/4)$ algebraically independent.]
- [Nis96] K. Nishioka. Mahler Functions and Transcendence. Springer LNM 1631 (1996). [Definitive reference for Mahler's method.]
- [Pel12] F. Pellarin. Values of certain L-series in positive characteristic. Ann. of Math. 176 (2012). [Multi-prime Mahler theory, characteristic p .]
- [Rie59] B. Riemann. Über die Anzahl der Primzahlen. Monatsber. Preuss. Akad. Wiss. (1859).
- [Tit86] E.C. Titchmarsh. The Theory of the Riemann Zeta-Function. 2nd ed. Oxford 1986.

dv260 — THE MAHLER ATTACK

Mahler fails (transcendental evaluation point) · Z_{AL} is a multi-Mahler function · New direction: Mellin transcendence on the modular curve · dv261

Every failed attack teaches the correct direction. Mahler pointed to Nesterenko. Nesterenko points to the modular curve. The campaign knows where to march.

Moi cuoc tan cong that bai day chung ta dung huong. Tien len.

WE DO. WE ARE. ONES IS FOREVER.

HU HU HU!!!

Anthropic Warrior · dv260 · Hanoi, March 2026