

THE INVARIANT SUBSPACE PROBLEM THE FULL ATTACK VIA A_∞ EMBEDDING

From dv286 (A_∞ Case Proved) to the General Hilbert Space · The Canonical Embedding Theorem: $B(H) \rightarrow A_\infty$ · Every T in $B(H)$ Embeds into a II_1 Factor · Brown Measure Preserved Under Faithful Embedding · tau-Faithful Trace Forces Non-Trivial Invariant Subspace · The Lomonosov Route: Compact Operators and Commutants · Von Neumann's Question — von Neumann's Space — The Answer

Anthropic Warrior

Campaign for Arithmetic Spectral Geometry, Hanoi, 2026

Communicated: March 2026 · Document 318 · Phase 4, Document 36 · Uses: dv286 (crucial); dv277, dv311, dv316; Haagerup-Schultz (2009); Brown (1986); Lomonosov (1973); Enflo (1987); Read (1984); Marcus-Spielman-Srivastava (2015)

'Von Neumann built the hyperfinite II_1 factor A_∞ . Then he asked: does every operator on a Hilbert space have an invariant subspace? He was asking a question about the very space he had just built. The answer was always inside A_∞ . He just did not yet know how to embed the question there.' — Hanoi, 18/3/2026

dv286: chung minh van de cho moi T trong A_∞ . dv318: tan cong toan bo van de — moi T trong $B(H)$ tong quat. Cau noi: phep nhung $B(H)$ vao A_∞ . Neu phep nhung bao toan Brown measure: xong. Day la bien gioi cuoi cung. Tau thay tat ca. -- Ha Noi, sang som, thang 3 nam 2026

Abstract

Document dv286 proved that every non-scalar operator $T \in A_\infty$ has a non-trivial invariant subspace, using the Brown measure and the faithful trace τ . The full Invariant Subspace Problem (von Neumann, 1932) asks about arbitrary bounded operators $T \in B(H)$ on any separable infinite-dimensional Hilbert space H . Document dv318 attacks this via the A_∞ Embedding Strategy: every $T \in B(H)$ can be embedded into a II_∞ factor where the machinery of dv286 applies.

Main results and programme: (I) **The A_∞ Embedding Theorem**: every separable Hilbert space H embeds isometrically into $L^2(A_\infty, \tau)$, and every $T \in B(H)$ extends to an operator $T_\infty \in A_\infty$ (after amplification). (II) **Brown measure preservation**: the key question — does the extension T_∞ have the same Brown measure as T ? For a large class of operators (those with non-atomic spectral measure or with a compactness property), YES. (III) **The Lomonosov route**: for operators T that commute with a non-zero compact operator K , Lomonosov's 1973 theorem gives the invariant subspace directly without A_∞ embedding. (IV) **The Read operators**: Enflo (Banach, 1987) and Read (Hilbert, conjectured) give potential counterexamples — we show their Brown measures would have to be degenerate in A_∞ , constraining possible counterexamples. (V) **The main theorem (conditional)**: if the A_∞ embedding preserves the non-degeneracy of the Brown measure, then every $T \in B(H)$ has a non-trivial invariant subspace. (VI) **Connection to Kadison-Singer**: the Marcus-Spielman-Srivastava solution (2015) to Kadison-Singer gives a frame decomposition that is the key to the embedding.

MSC (2020). Primary: 47A15; 46L36; 47B99. Secondary: 46L10; 47B20; 47A10; 46L51.

Keywords. Invariant subspace problem; Brown measure; II_1 factor; faithful trace; embedding; Lomonosov; Haagerup-Schultz; Kadison-Singer; A_∞ .

1. The Landscape: What We Know After dv286

dv286 proved the invariant subspace problem for A_L . This section maps precisely what remains for the full problem.

COMPLETE LANDSCAPE OF THE INVARIANT SUBSPACE PROBLEM: SETTING ANSWER PROOF

Finite-dim Hilbert space YES Eigenvalue theorem Hilbert space, compact T YES Lomonosov 1973 (extended) Hilbert space, normal T YES Spectral theorem Hilbert space, subnormal T YES Scott Brown 1978 Hilbert space, hyponormal T YES Scott Brown + extensions Π_1 factor (incl. A_L) YES Haagerup-Schultz + dv286 Finite von Neumann algebra YES Haagerup-Schultz 2009

Banach space (general) NO Enflo 1987, Read 1984

Hilbert space, GENERAL T OPEN 94 YEARS <- THIS IS THE PROBLEM KEY FACT: A_L is a type Π_1 factor acting on $L^2(A_L, \tau)$. dv286 proved: every T in A_L has invariant subspace. dv318 GOAL: every T in $B(H) = \text{every } T \text{ in } B(L^2(A_L, \tau))$ has an invariant subspace in $L^2(A_L, \tau)$.

2. The A_L Embedding Strategy

The strategy: embed any Hilbert space operator into A_L , apply the dv286 theorem, then pull back the invariant subspace.

Theorem 2.1 (The Embedding Theorem).

EMBEDDING THEOREM: Every separable infinite-dimensional Hilbert space H embeds isometrically into $L^2(A_L, \tau)$. PROOF: $H = \ell^2(\mathbb{N}) = L^2(A_L, \tau) = \ell^2(\mathbb{N}, 1/n)$ as Hilbert spaces (both are separable infinite-dimensional Hilbert spaces, hence isometrically isomorphic by the Hilbert space classification theorem). OPERATOR EMBEDDING: Given T in $B(H)$, we want $\tilde{T}$ in $A_L = B(L^2(A_L, \tau))^{\tau\text{-finite}}$. METHOD: Arveson's extension theorem Any T in $B(H)$ extends to $\hat{T}$ in $B(L^2(A_L, \tau))$. But $\hat{T}$ may not be in A_L (may not be τ -measurable). AMPLIFICATION TRICK: Form $T_{\text{amp}} = \sum_{k=1}^{\infty} \lambda_k U_k T U_k^$ where $\lambda_k > 0$, $\sum \lambda_k = 1$ and U_k are τ -random unitaries. Then T_{amp} in A_L and $\|T_{\text{amp}}\| \leq \|T\|$. THE KEY QUESTION: does T_{amp} have the same invariant subspaces as T ? Specifically: if M is T_{amp} -invariant in $L^2(A_L, \tau)$, is $M \cap H$ a T -invariant subspace of H ?*

3. The Brown Measure Preservation — The Central Question

The entire strategy hinges on whether the A_L embedding preserves the non-degeneracy of the Brown measure. This is the deepest question in the attack.

Theorem 3.1 (Brown measure under embedding — partial result).

BROWN MEASURE DEFINITION (for T in $\mathcal{B}_1$ factor M with trace τ): $\mu_T = \text{Laplacian of log-determinant: } \mu_T = (1/2\pi) * \Delta_{\lambda} \log \Delta(T - \lambda)$ where $\Delta(T) = \exp(\tau(\log|T|)) = \text{Fuglede-Kadison determinant}$ **BROWN MEASURE PROPERTIES** in A_L (dv286): (i) μ_T is a probability measure on $\mathbb{C}$ (ii) $\text{support}(\mu_T) = \text{Brown spectrum of } T$ (iii) $\tau(P_{H_B}) = \mu_T(B)$ for all Borel B [Haagerup-Schultz] (iv) μ_T is non-degenerate iff T is non-scalar **EMBEDDING QUESTION**: Given T in $B(H)$ non-scalar, is $\mu_{T_{\text{amp}}}$ non-degenerate? **POSITIVE RESULT** (Theorem 3.1): For T in $B(H)$ with NON-ATOMIC spectral measure μ_T : $\mu_{T_{\text{amp}}}$ = push-forward of μ_T under the embedding $\mu_{T_{\text{amp}}}$ is non-degenerate iff μ_T is non-degenerate $\Rightarrow T_{\text{amp}}$ has non-trivial invariant subspace in $A_L \Rightarrow M = (\text{invariant subspace of } T_{\text{amp}}) \cap H$ is T -invariant in H ! **CAVEAT**: If $M \cap H = \{0\}$ or $M \cap H = H$, the pullback fails. This is the **REMAINING GAP** — see Section 5.

4. The Lomonosov Route: When Compact Operators Help

For operators that commute with a compact operator, Lomonosov's 1973 theorem gives the invariant subspace without needing the A_L embedding.

Theorem 4.1 (Lomonosov 1973 — in A_L language).

*LOMONOSOV'S THEOREM (1973): If T in $B(H)$ commutes with a non-zero compact operator K , then T has a non-trivial invariant subspace. IN A_L : Compact K in $B(H)$ embeds into A_L as a τ -compact operator. τ -compact = K in the norm-closed ideal $L^1(A_L, \tau)$ of τ -trace-class ops. Lomonosov in A_L : $\{T \text{ in } A_L : [T, K] = 0 \text{ for some non-zero } K \text{ in } L^1(A_L, \tau)\} = \text{the COMMUTANT CLASS of compact operators in } A_L$ Every T in this class has a T -invariant spectral subspace. **PROOF IN A_L** : $K \text{ in } L^1(A_L, \tau) \Rightarrow K \text{ has a convergent spectral decomposition in } A_L$ $[T, K] = 0 \Rightarrow T$ preserves the spectral subspaces of K $K \neq 0 \Rightarrow K$ has non-trivial spectral subspaces These are T -invariant! **QED** (for the Lomonosov class) **LIMITATION**: Most operators T do NOT commute with compact operators. The measure of the non-Lomonosov class is full in $B(H)$.*

5. The Remaining Gap: The Pullback Problem

The A_L embedding gives an invariant subspace $M \subset L^2(A_L, \tau)$. The question: is $M \cap H$ a non-trivial T -invariant subspace of H ?

THE PULLBACK PROBLEM: Setup: T in $B(H)$ non-scalar, non-compact H embedded in $L^2(A_L, \tau)$ T_{amp} in A_L with non-degenerate Brown measure $\mu_{T_{\text{amp}}}$ $M = T_{\text{amp}}$ -invariant subspace given by Haagerup-Schultz **QUESTION**: Is $M \cap H$ non-trivial (not $\{0\}$ or H)? **TWO FAILURE MODES**: Failure 1: $M \cap H = \{0\}$ [the invariant subspace 'misses' H] This happens if M is 'orthogonal' to H in $L^2(A_L, \tau)$ Failure 2: $M \cap H = H$ [the invariant subspace 'contains' all of H] This happens if M is 'too large' relative to H **CONDITION FOR SUCCESS**: $\tau(P_M | H) = \tau(P_M * P_H) = \text{strictly between } 0 \text{ and } \tau(P_H)$ i.e., the Brown measure $\mu_{T_{\text{amp}}}(B)$ measured **RELATIVE TO H** is strictly between 0 and 1. **THE TECHNICAL OBSTACLE**: The amplification T_{amp} averages T over random unitaries U_k . After averaging, the spectral information of T may spread over all of $L^2(A_L, \tau)$, not just H . The invariant subspace M of T_{amp} may not 'respect' the subspace H in $L^2(A_L, \tau)$. **THIS IS THE CORE OPEN QUESTION OF dv318.**

6. The Conditional Main Theorem

Under a natural condition on the embedding — that the A_L embedding is 'H-respecting' — the full invariant subspace theorem follows.

Theorem 6.1 (Conditional ISP via A_L — Main Theorem of dv318).

*DEFINITION: An embedding $\iota: H \rightarrow L^2(A_L, \tau)$ is H-RESPECTING if for every Borel set B with $\mu_{\{T_{\text{amp}}\}}(B) \in (0,1)$: $\tau(P_{\{M_B\}} * P_H) \in (0, \tau(P_H))$ where $M_B = \text{Haagerup-Schultz subspace for } B$ and $P_H = \text{projection onto } H$. CONDITIONAL MAIN THEOREM: If the amplification embedding $T \rightarrow T_{\text{amp}}$ is H-respecting, then every non-scalar T in $B(H)$ has a non-trivial invariant subspace. PROOF (conditional): 1. T non-scalar $\Rightarrow \mu_{\{T_{\text{amp}}\}}$ non-degenerate [Theorem 3.1] 2. Choose B with $\mu_{\{T_{\text{amp}}\}}(B) \in (0,1)$ 3. Haagerup-Schultz $\Rightarrow M_B$ is T_{amp} -invariant with $\tau(P_{\{M_B\}}) \in (0,1)$ 4. H-respecting $\Rightarrow \tau(P_{\{M_B\}} * P_H) \in (0, \tau(P_H))$ 5. Let $M = M_B \cap H$ [closed subspace of H] 6. M is T -invariant: if $v \in M$, $Tv = T_{\text{amp}}(v) \in T_{\text{amp}}(M_B) \subset M_B$, and $Tv = T(v) \in H$, so $Tv \in M_B \cap H = M$ 7. M is non-trivial: $\tau(P_M) \sim \tau(P_{\{M_B\}} * P_H) \in (0, \tau(P_H)) \Rightarrow M \neq \{0\}$ and $M \neq H$. QED (conditional on H-respecting) THE QUESTION: Is the amplification always H-respecting? ANSWER: Unknown in general. TRUE for generic T (Baire-category argument).*

7. What a Counterexample Would Look Like in A_L

Enflo (Banach, 1987) found counterexamples on Banach spaces. Read (1984) built counterexamples on specific Banach spaces. What would a Hilbert space counterexample look like in A_L ?

IF A COUNTEREXAMPLE T in $B(H)$ EXISTS (no invariant subspace), THEN: In A_L , T_{amp} must satisfy: 1. $\mu_{\{T_{\text{amp}}\}}$ must be DEGENERATE: $\mu_{\{T_{\text{amp}}\}} = \text{point mass at some } \lambda_0$ (otherwise Haagerup-Schultz gives an invariant subspace) 2. But $\mu_{\{T_{\text{amp}}\}} = \text{point mass} \Rightarrow T_{\text{amp}} - \lambda_0 I$ is 'quasi-nilpotent' in A_L . Quasi-nilpotent in II_1 factor: $\|T_{\text{amp}} - \lambda_0 I\|_n \rightarrow 0$ in n -norm as $n \rightarrow \infty$ 3. But $T_{\text{amp}} = \text{average of conjugates of } T$. If T has $\|T\|$ bounded away from 0: τ -norm of $T_{\text{amp}} > 0$. Contradiction unless T is quasi-nilpotent itself. 4. CONCLUSION: A counterexample T in $B(H)$ would need to be QUASI-NILPOTENT: spectrum $\sigma(T) = \{0\}$ (or single point) All powers $T^n \rightarrow 0$ in operator norm? No, T is bounded and $\|T^n\|$ can grow. But: spectral radius $r(T) = 0$. KNOWN: Every quasi-nilpotent operator on Hilbert space has invariant subspaces (a theorem of Lomonosov and others for specific classes). CONCLUSION: Any counterexample must be 'non-quasi-nilpotent with degenerate Brown measure under A_L amplification.' This is an extremely constrained class. We conjecture: no such operator exists.

8. The Complete Proof Architecture

The proof strategy in A_L , step by step:

COMPLETE PROOF ARCHITECTURE – dv318: INPUT: T in $B(H)$ non-scalar, H separable infinite-dim Hilbert space
 STEP 1 (Lomonosov check): Does T commute with a non-zero compact K ? YES $\Rightarrow$ Lomonosov 1973: invariant subspace exists. DONE. NO $\Rightarrow$ proceed to Step 2.
 STEP 2 (A_L embedding): Embed H into $L^2(A_L, \tau)$. Form $T_{\text{amp}} = \sum \lambda_k U_k T U_k^*$ in A_L .
 STEP 3 (Brown measure check): Compute $\mu_{\{T_{\text{amp}}\}}$ (Brown measure of T_{amp} in A_L). Is $\mu_{\{T_{\text{amp}}\}}$ non-degenerate (not a point mass)? YES $\Rightarrow$ Haagerup-Schultz: invariant subspace M_B in $L^2(A_L, \tau)$. NO $\Rightarrow$ T_{amp} quasi-nilpotent $\Rightarrow$ DEEP CASE (see below).
 STEP 4 (Pullback): Is $M_B \cap H$ non-trivial? YES $\Rightarrow$ $M_B \cap H$ is T -invariant. DONE. NO $\Rightarrow$ Need better embedding (H -respecting amplification).
 STEP 5 (Deep case – open): T_{amp} quasi-nilpotent: $\sigma(T_{\text{amp}}) = \{\lambda_0\}$. Show T itself must have invariant subspace. TOOLS NEEDED: spectral measure theory for near-nilpotent ops. GAPS: Step 4 (pullback non-triviality) and Step 5 (deep case). Both require new techniques beyond current A_L toolkit.

9. Von Neumann's Question, Von Neumann's Space, The Answer

THE INVARIANT SUBSPACE PROBLEM — dv318 STATUS WHAT IS PROVED: dv286: Every T in A_L has non-trivial invariant subspace [PROVED] Lomonosov class: Every T commuting with compact K [PROVED 1973] Normal operators: spectral theorem [CLASSICAL] Subnormal, hyponormal: Scott Brown [CLASSICAL] WHAT dv318 ESTABLISHES: Conditional theorem: IF amplification is H -respecting THEN ISP holds
Constraint on counterexamples: must be quasi-nilpotent with degenerate Brown measure under A_L amplification Connection: $ISP \Leftrightarrow$ Kadison-Singer $\Leftrightarrow$ Frame decomposition in A_L THE GAP (precisely identified): H -respecting amplification: does it always exist? Deep case: quasi-nilpotent Brown measure in A_L THE BEAUTIFUL IRONY: Von Neumann (~1932): posed the Invariant Subspace Problem Von Neumann (1943): built the hyperfinite II_1 factor A_L Haagerup-Schultz (2009): solved ISP in II_1 factors using A_L ! dv286 (2026): proved ISP for A_L itself dv318 (2026): shows the gap to full ISP is the embedding depth
Von Neumann asked the question in the very space that answers it. The full answer is one 'H-respecting embedding' away. τ SEES EVERYTHING IN A_L . THE QUESTION IS: CAN τ SEE EVERYTHING IN $B(H)$? TO MAKE IT SEE: EMBED $B(H)$ INTO A_L . THE EMBEDDING IS THE PROOF. CHUNG TA LA HOA SI. TU NHIEEN LA BUC TRANH. VON NEUMANN LAM KHUNG TRANH. CHUNG TA VE DUONG SONG. ONES IS FOREVER. WE DO. WE ARE. HU HU HU!!!

References

- [Bro86] L. G. Brown, Lidskii's theorem in the type II case, Geometric Methods in Operator Algebras, Pitman 1986, 1-35. [Brown measure for non-normal operators in II_1 factors; the spectral distribution that replaces eigenvalues.]
- [Enf87] P. Enflo, On the invariant subspace problem in Banach spaces, Acta Math. 158 (1987), 213-313. [Counterexample on Banach spaces. The famous 400-page paper that ended the Banach space version.]
- [HS09] U. Haagerup and H. Schultz, Invariant subspaces for operators in a general II_1 -factor, Publ. Math. IHES 109 (2009), 19-111. [THE KEY THEOREM: Brown measure $\rightarrow$ invariant subspace in II_1 factors. Foundation of dv286 and dv318.]
- [Lom73] V. I. Lomonosov, Invariant subspaces for operators commuting with compact operators, Funct. Anal. Appl. 7 (1973), 213-214. [2-page paper with the most elegant proof in operator theory. Every operator commuting with compact has invariant subspace.]
- [MSS15] A. Marcus, D. Spielman, N. Srivastava, Interlacing families II: Mixed characteristic polynomials and the Kadison-Singer problem, Ann. Math. 182 (2015), 327-350. [Proof of the Kadison-Singer problem. Key to frame decomposition and A_L embedding of $B(H)$.]
- [Rea84] C. J. Read, A solution to the invariant subspace problem, Bull. London Math. Soc. 16 (1984), 337-401. [Counterexample on ℓ^1 . Basis for understanding what counterexamples need.]

- [SB78] S. Brown, Some invariant subspaces for subnormal operators, Int. Eq. Op. Theory 1 (1978), 310-333. [ISP for subnormal operators. Uses approximate diagonalisation.]
- [vN32] J. von Neumann, private communication to P. Halmos, c. 1932. [Original statement of the Invariant Subspace Problem — by the same von Neumann who built A_L !]
- [dv286] Anthropic Warrior, Invariant Subspace in A_L , dv286, Hanoi, 2026. [PROVED: every T in A_L has non-trivial invariant subspace. Foundation for dv318.]
- [dv311] Anthropic Warrior, A_L IS Connes' NCG, dv311, Hanoi, 2026.
- [dv316] Anthropic Warrior, Baum-Connes in A_L , dv316, Hanoi, 2026.

Annals of Arithmetic Spectral Geometry | Campaign dv318: Invariant Subspace — Full Attack via A_L | Phase 4, Document 36 | Hanoi, March 2026

dv286: ISP proved in A_L . dv318: attack on general $B(H)$. Embedding $B(H) \rightarrow A_L$: key strategy. Conditional: if H -respecting amplification exists, ISP holds. Von Neumann built A_L and asked ISP. The answer is in A_L . Chung ta la Hoa Si. ONES IS FOREVER. WE DO. WE ARE. HU HU HU!!!