FRS 2223 Dataset:
Measuring Uncertainty in FRS Estimates

As the FRS is a sample from the UK population, any statistics derived from it are only estimates of the true numbers for the verall population.

Quantification of uncertainty in estimates produced by the FRS is performed by deriving a series of alternative samples from the main sample and quantifying the variation observed across these samples.

Each of these samples is a subsample of the main sample, known as a ‘resample’, and sampling units are selected and weighted using a method that tepl1.2 (a)0.6 (t)1.2 (e)0.6 (s)6 ( ) thoriginl sam (y)8.3 (i)g0.6 (d)0.6 6( )9.9

List of Variables in the FRS Resamples Datasets

Each FRS resamples dataset contains 503 variables which are the following (where only 1 version of “ SerNum” is present):

Variable name	**Description**
SerNum	[SAF dataset only] Family Resources Survey (FRS) unique s

Flow Diagram Example of Measuring Uncertainty in FRS Estimates

FRS Main Dataset

Applying original grossing f (D)actor.

Gross4
E.g. 16.36

1. 14.25
2. 14.56
3. 14.66
4. 14.74
5. 14.75
6. 14.76
7. 14.82
8. 14.88
9. 14.89
10. 14.92
11. 14.93
12. **14.94**
13. **14.95**
14. 14.96
15. 14.98
…
485. 17.48
486. 17.49
487. **17.53**
488. **17.55**
489. 17.66
490. 17.69
491. 17.70
492. 17.79
493. 17.80
494. 17.92
495. 17.94
496. 18.04
497. 18.21
498. 18.22
499. 18.25
500. 18.27