

Homework №6

Author: David Oniani
Instructor: Tommy Occhipinti

October 12, 2018

43. Prove that for all $n \in \mathbb{Z}^+$, $11^n - 6$ is divisible by 5.

Let's prove this by induction. For this we have to have a base case and an inductive hypothesis.

Base case: if $n = 1$, $11^n - 6 = 11 - 6 = 5$ and thus, since 5 is divisible by 5, $11^n - 6$ is divisible by 5. Hence, the base case check is done.

Inductive case: suppose for all $k \in \mathbb{Z}^+$, $11^k - 6$ is divisible by 5 and prove that $11^{k+1} - 6$ is divisible by 5. Notice that

$$11^{k+1} - 6 = 11 \times 11^k - 6 = 10 \times 11^k + (11^k - 6)$$

Now, once again, notice that 10×11^k is divisible by 5 since 10 is a multiple of 5. According to our assumption, $11^k - 6$ is also divisible by 5. Finally, we get that $11^{k+1} - 6 = 10 \times 11^k + (11^k - 6)$ which means that $11^{k+1} - 6$ is the sum of two numbers which are both divisible by 5.

Q.E.D.

44. Prove that for $n \in \mathbb{Z}^+$, we have $1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$.

Let's prove this by taking an inductive approach.

Base case: if $n = 1$, $1^2 = \frac{1(1+1)(2+1)}{6} = \frac{1 \times 2 \times 3}{6} = 1$ and thus, the base case check is done.

Inductive hypothesis: suppose for $k \in \mathbb{Z}^+$, $1^2 + 2^2 + 3^2 + \dots + k^2 = \frac{k(k+1)(2k+1)}{6}$ and prove that $1^2 + 2^2 + 3^2 + \dots + k^2 + (k+1)^2 = \frac{(k+1)(k+2)(2k+3)}{6}$. According to our initial assumption, we have:

$$\begin{aligned}
1^2 + 2^2 + 3^2 + \dots + k^2 + (k+1)^2 &= (1^2 + 2^2 + 3^2 + \dots + k^2) + (k+1)^2 = \\
1^2 + 2^2 + 3^2 + \dots + k^2 + (k+1)^2 &= \frac{k(k+1)(2k+1)}{6} + (k+1)^2 = \\
&= \frac{k(k+1)(2k+1)}{6} + \frac{6(k+1)^2}{6} = \\
&= \frac{k(k+1)(2k+1) + 6(k+1)^2}{6} = \\
&= \frac{(k+1)(k(2k+1) + 6(k+1))}{6} = \\
&= \frac{(k+1)(2k^2 + k + 6k + 6)}{6} = \\
&= \frac{(k+1)(2k^2 + 7k + 6)}{6} = \\
&= \frac{(k+1)((k+2)(2k+3))}{6}
\end{aligned}$$

Thus, we proved that $1^2 + 2^2 + 3^2 + \dots + k^2 + (k+1)^2 = \frac{(k+1)(k+2)(2k+3)}{6}$ which finishes our proof.

Q.E.D.

45. Prove that for all $n \in \mathbb{Z}^+$, $2^{n+2} + 3^{2n+1}$ is divisible by 7.

To prove it using induction, we'll need a base case and an inductive hypothesis.

Base case: let $n = 1$, then $2^{1+2} + 3^{2 \times 1 + 1} = 2^3 + 3^3 = 8 + 27 = 35 = 7 \times 5$ thus is divisible by 7. And we've checked the base case.

Inductive hypothesis: now, we have to prove the inductive hypothesis. Here is the hypothesis: suppose that for all $k \in \mathbb{Z}^+$, $2^{k+2} + 3^{2k+1}$ is divisible by 7 and show that $2^{k+3} + 3^{2k+3}$ is also divisible by 7.

Notice that

$$\begin{aligned}
2^{k+3} + 3^{2k+3} &= \underbrace{2^{k+2} + 2^{k+2}}_{2^{k+3}} + \underbrace{3^{2k+1} + 3^{2k+1} + 7 \times 3^{2k+1}}_{3^{2k+3}} = \\
&= 2 \times (2^{k+2} + 3^{2k+1}) + 7 \times 3^{2k+1}
\end{aligned}$$

And now it's clear that since $2^{k+2} + 3^{2k+1}$ and $7 \times 3^{2k+1}$ are the multiples of 7, $2 \times (2^{k+2} + 3^{2k+1}) + 7 \times 3^{2k+1}$ is also a multiple of 7 and thus is divisible by 7. Hence, we proved the inductive hypothesis and it concludes our proof.

Q.E.D.

46. Let F_n denote the n^{th} Fibonacci number. Prove that $\sum_{k=1}^n F_k^2 = F_n F_{n+1}$.

Let's prove this using induction. For this we have to have a base case and inductive hypothesis.

Let's first look at the Fibonacci sequence. It is 1, 1, 2, 3, 5, 8, 13, 21 ... where each element is the sum of two previous elements.

Base case: let $k = 1$, then $\sum_{k=1}^1 F_k^2 = 1 \times 1 = 1$ and indeed the second element of the sequence is 1.

Inductive hypothesis: let F_m denote the m^{th} Fibonacci number and also suppose that $\sum_{j=1}^m F_j^2 = F_m F_{m+1}$.

Then, we have to prove that $\sum_{j=1}^{m+1} F_j^2 = F_{m+1} F_{m+2}$.

Notice that $\sum_{j=1}^{m+1} F_j^2 = F_1^2 + F_2^2 + F_3^2 + \dots + F_{m+1}^2$ and using our assumption, we get:

$$F_1^2 + F_2^2 + F_3^2 + \dots + F_{m+1}^2 = F_m F_{m+1} + F_{m+1}^2 = F_{m+1} (F_m + F_{m+1})$$

. Now, let's remember the way Fibonacci numbers are created. To get the next Fibonacci number, one must sum up the previous two thus,

$F_m + F_{m+1} = F_{m+2}$ and we finally get that $\sum_{j=1}^{m+1} F_j^2 = F_1^2 + F_2^2 + F_3^2 + \dots + F_{m+1}^2 = F_m F_{m+1} + F_{m+1}^2 = F_{m+1} (F_m + F_{m+1}) = F_{m+1} F_{m+2}$. Hence,

we proved that $\sum_{j=1}^{m+1} F_j^2 = F_{m+1} F_{m+2}$ and this concludes our inductive reasoning as well as the proof.

Finally, if F_n denotes the n^{th} Fibonacci number, $\sum_{k=1}^n F_k^2 = F_n F_{n+1}$.

Q.E.D.

47. Prove that for all $n \in \mathbb{Z}^+_{\geq 12}$ we have $n! \geq 5^n$.

To prove the statement above, we can use induction. We need to have a base case as well as an inductive hypothesis.

Base case: let $n = 12$, then we have $12! > 5^{12}$ which is true since $12! = 479001600$ and $5^{12} = 244140625$ (thus $12! > 5^{12}$).

Inductive hypothesis: suppose that for every $k \in \mathbb{Z}^+_{\geq 12}$ we have $k! \geq 5^k$ and prove that $(k+1)! > 5^{k+1}$.

HAVE TO FINISH THIS.

Q.E.D.