

Analysis of Melody Through Key Definition and Generation of Complementary Harmonies

Onkur Sen
Rice University, Houston, TX
onkur@rice.edu

Kurt Stallmann
Rice University, Houston, TX
stallmann@rice.edu

Abstract

Algorithmic music engages computational methods to create sound structures. Previous efforts in algorithmic music include automatic generation of music, sometimes through the application of biological models, e.g., genetic programming and genetic algorithms. Other notable methods use random walks and Markov chains.

Our focus is to create a mathematical model of theoretical notions found in common practice music. We provide a method of analysis that determines the key of an input melodic line by utilizing two different correlation metrics depending on consonance and scale membership, respectively.

We then generate a harmonic progression that supports the melody within the determined key. To do this, we reuse the previous metrics with additional constraints such as harmonic relations with the given key and transitions between chords.

We test this model using prominent compositions from musical literature as input and noting how the output of the algorithm meets expectations or provides viable alternatives.

1 Introduction

Algorithmic music is a relatively new form of music generation that utilizes computational approaches to create coherent musical structures. Much work has been done in this field within the context of a particular style. David Cope has previously worked with generating Western classical music in the style of Mozart. Microsoft’s Songsmith generates harmonies for a sung melody using Markov models. In particular, we aim to study how much information we can obtain about a work of tonal music given only its melodic line. To this effect, we wish to find the key of the piece and generate a complementary harmony that maintains the standards provided by common practice.

This problem presents multiple challenges in different areas of music. First, different melodies convey different amounts and types of information, and often times tonal ambiguity arises when considering only the melody. However, we can use this ambiguity to our advantage when considering the generation of a variation upon an original work. We can follow multiple paths that will produce different outcomes which are all musically viable in the context of common practice music.

Second, the presence of the harmony is interrelated with the melody towards defining a certain key. Thus, when the harmony is not given, we must choose a harmony that articulates the melody tonally. In particular, most harmonies support the melody in supportive or antagonistic ways through conjunct and disjunct motion. Therefore, we must select a type of harmony that properly reinforces the melody.

2 Determining the Key of an Input Score

The question we pose here is simple: How can we find the key of a piece of music given only its melody? We define the key of the piece as having two components, a **tonic**, the root of the key, and a **mode**, which is either major or minor. Our approach is broken up into the following steps:

2.1 Note-Counting

We assume the musical work is sampled at some regular time interval, i.e., we are assuming a discrete-time input signal M . Our first step in analysis is to simply count up the number of occurrences of each pitch in the chromatic scale. This indicates the dominant contributions among the scale to the character of the piece and signals a first step towards understanding the key of the piece.

Mathematically, we represent pitches on a chromatic scale as 12 values between 0 and 11, with 0 being C and 11 being B, in compliance with the commonly-used MIDI system. For our purposes, notes that differ by octaves are considered equivalent and thus undergo a modulo 12 operation. We can thus define the **note-counting operation** on a pitch p as:

$$n(p_i) = \#\{x \in S; x = p_i\}, \quad p_i = 0, 1, \dots, 11.$$

Alternatively, we can determine the importance of notes with **weights that represent the relative frequency** of notes. This is defined as follows:

$$w(p_i) = \frac{n(p_i)}{\sum_{j=0}^{11} n(p_j)}.$$

2.2 Defining Correlation Functions for Characterizing Keys

In order to measure the viability of a certain key with the given input score, we define two different **correlation metrics** with different criteria.

2.2.1 Consonance Correlation Function

The consonance correlation emphasizes how euphonious a pitch (or a sequence of pitches) is with respect to a certain key in accordance with common practice music. We first define the consonance of a note with respect to a given key in terms of the note's relation to the tonic of the key (see Table 1). These values were experimentally adjusted to produce correct results for a set of trial pieces.

The consonance correlation function C_C for a key is then simply defined as the sum of the consonance relations for each pitch on the chromatic scale weighted by its appearance in the piece:

$$C_C(key) = \sum_{i=0}^{11} w(p_i) \times \text{consonance}(p_i, key).$$

Note that for any key, $0 \leq C_C(key) \leq 1$, so that the consonance correlation is normalized.

Relation of Note to Tonic	$consonance(note, key)$
1^{st}	1
5^{th}	$\frac{2}{3}$
4^{th}	$\frac{1}{3}$
$3^{rd}, 6^{th}$ (same mode)	$\frac{1}{3}$
Other	0

Table 1: Consonance relations for a note with respect to a given key. For the 3^{rd} and 6^{th} , the note must be in the same mode as the key; else, the consonance relation is 0 (e.g., A^b with respect to C Major).

2.2.2 Membership Correlation Function

The membership correlation emphasizes how well a pitch (or a sequence of pitches) fits into a given scale which is associated with a key. We define the membership function between a note and a scale as simply the characteristic function of that scale. More explicitly:

$$membership(note, scale) = \begin{cases} 1 & \text{if } note \in scale \\ 0 & \text{if } note \notin scale \end{cases}$$

The membership correlation function C_M for a key is then the weighted sum, similar to the consonance correlation:

$$C_M(key) = \sum_{i=0}^{11} w(p_i) \times membership(p_i, scale(key)).$$

Here too, we see that the membership function is normalized, i.e., $0 \leq C_M(key) \leq 1$ for any key.

2.3 Final Steps: Using Correlation Functions to Determine Key

We first find the consonance and membership correlations for each key in the chromatic scale. To find the key of the piece, we wish to maximize both of those correlation functions. However, a perfect membership correlation is not viable in common practice music as we often have the introduction of accidentals as well as temporary deviations into other keys.

Thus, we settle on a threshold value of 0.9 for the membership correlation. All keys with a membership correlation lower than this value are discarded. The key of the piece is then chosen from the remaining options as that which maximizes the consonance correlation and would hence best characterize the piece.

3 Creating a Complementary Harmonic Progression

We have obtained a significant amount of information now that we know the key of the piece. How can we use this information to generate a harmony that is plausible in the realm of common practice? We set forth a few reasonable approximations to tremendously simplify the problem:

- We uniformly divide the piece into smaller subdivisions, usually representing beats or measures.

- We assume that each subdivision has one "key" or "chord" associated with it.
- We limit our choice of keys to (with respect to the key of the piece): the tonic, the dominant, the subdominant, and the relative minor/major (depending on the mode of the key). Note that when we attempted to find the key of the entire piece, our key space could not be narrowed down because we had no other information. However, now that we had arrived at a key for the entire piece, we can reasonably limit our key space to those keys that are strongly associated with it.
- We do *not* assume any positioning (e.g., inversions, octaves) for the chords associated with each subdivision.

Using these assumptions, we simply apply the procedure for determining the key of a piece as before, except with our scope limited to a subdivision as input and the choice of keys as listed above. After doing this for all subdivisions, we have a list of keys which also represent a accompanying **chord progression** for the piece.

4 Results and Discussion

4.1 Key Definition

We focused on applying this method of key definition to samples of Western classical music, particularly those from the Baroque and Classical periods. Overall, we were able to successfully determine keys for 15 different pieces, which included the following:

- Twinkle, Twinkle, Little Star (in C Major)
- Bach: Invention No. 1 in C Major; Minuet in G Major; Minuet in G minor
- Mozart: K. 545, 1st mvt. (C Major) and 2nd mvt. (G Major)
- Beethoven: Piano Sonata No. 1, 1st mvt. (F minor); Piano Sonata No. 8 (Pathétique), 2nd mvt. (C^{sharp} Major)

However, we observed limitations of our approach when considering pieces from the Romantic period, such as Chopin's Prelude in Em, where the melody simply consists of two notes. This suggests that in Chopin and in Romantic music in general, the harmony begins to play a more dominant role compared to the melody and becomes a much stronger component of the score.

Another interesting example was the Schumann Romance, where the key returned was the relative minor of the actual key. However, the melody was harmonized on parallel thirds, and when the lower melody was accounted for, the proper key was returned. Thus, we see that as music progresses historically, it becomes difficult to precisely determine the key using common practice conventions. Of course, we can utilize this to our advantage and produce a harmonic progression based around the same melody in another key for the sake of variation. Nevertheless, these contrary examples give us a much more realistic indication of the scope of our approach as well as the problem space.

4.2 Harmony Generation

The output of a sequence of keys or chords for the harmonic progression leaves a great deal of flexibility for the form of the accompaniment. This is both a boon and a burden because while it does leave us with an open problem that can be attacked from multiple angles, we can explore each of these paths and the variations on the original melody-harmony relationship that they develop. Thus, we must consider the type of relation of the harmony with respect to the melody that we wish to emphasize. Some examples include:

- We must consider both the “vertical” support that a chord can provide a melody as well as the “horizontal” path that the chord progression lays out in terms of expressing the piece.
- A key for a chord leaves room for variation in inversions as well as placements in different octaves. This can result in parallel, contrary, and oblique movement as well as conjunct or disjunct motion.
- We also can raise the question of voicing since the melody is not necessarily assigned to a particular hand (if we consider the piano as our instrument) and since the analysis of melody is independent of placement within a certain octave.

Furthermore, we attempted to glean time information from the harmonic progression by analyzing a **return to tonic**, i.e., the length of time it takes until a the chord assigned to a subdivision is in the key of the entire piece. With pieces that were in the sonata form, for instance, we were able to see that this length was approximately twice the length of a measure. This roughly gives us the time signature in terms of subdivisions, which we can then use as an additional tool on shaping the harmony to be more applicable for the melody. We can improve our work here by implementing pattern recognition, e.g., accommodating for harmonic rhythm by being able to recognize cadences and other changes in tonal structure.

5 Conclusion

Our research goal was to produce a computational analysis technique for determining the key of an input melody and generate a complementary harmonic progression in accordance with common practice music. While we were able to see the success of our methods in many instances of Western classical music from the Baroque and Classical periods, we also saw the limitations in applying key definition to music from the Romantic period. Furthermore, our work in harmony generation, while relatively accurate, leaves many doors open on the actual form of the harmony, a fact that can be exploited in the future for variational purposes.