

Module 5 Notes (MATH-211)

Lillie Donato

08 July 2024

General Notes (and Definitions)

- Maxima and Minima

Absolute Maximum: Assume a function f is defined on a set D , and $x = c$ is a point in D . Then, $y = f(c)$ is an **absolute maximum value** of f on D if $f(c) \geq f(x)$ for every x in D . Changing the set on which f is defined may change the absolute maximum value.

Absolute Minimum: Assume a function f is defined on a set D , and $x = c$ is a point in D . Then, $y = f(c)$ is an **absolute minimum value** of f on D if $f(c) \leq f(x)$ for every x in D . Changing the set on which f is defined may change the absolute minimum value.

Extreme Value Theorem: A function that is continuous on a closed interval is guaranteed to have both an absolute maximum value and an absolute minimum value.

A discontinuous function, or a function defined on an interval that is not closed, may still have absolute extrema.

Local Maximum and Minimum Values: Assume $x = c$ is an interior point (not an endpoint) of some interval I in the domain of f . Then, $y = f(c)$ is a **local maximum value** of f if $f(c) \geq f(x)$ for every x in I , and $y = f(c)$ is a **local minimum value** of f if $f(c) \leq f(x)$ for every x in I .

Critical Points: An interior point $x = c$ of the domain of f is called a **critical point** of f if either $f'(c) = 0$ or $f'(c)$ does not exist.

Local Extreme Value Theorem: If a function f has a local maximum or a local minimum at a point $x = c$, then either $f'(c) = 0$ or $f'(c)$ does not exist.

If f has a local extreme, it must occur at a critical point.

Not every critical point is the location of a local extreme value.

For a continuous function f on a closed interval $[a, b]$, absolute extremes are guaranteed to exist, and they must occur either at the endpoints of interval or at critical points of f within the interval.

Examples

1. Locate absolute maxima and minima from a graph
Absolute Maximum: $f(c)$ and occurs at $x = c$
Absolute Minimum: None, as the $f(b)$ does not exist

2. Locate local maxima and minima from a graph
Absolute Min at $(a, f(a))$
Absolute Max at $(p, f(p))$
Local Max at $(p, f(p))$
Local Max at $(r, f(r))$
Local Min at $(q, f(q))$
Local Min at $(s, f(s))$

3. Find critical points of a function

$$f(t) = t^2 - 2 \ln(t^2 + 1)$$

$$f'(t) = \frac{2t(t+1)(t-1)}{t^2 + 1}$$

Critical Point at $x = -1$

Critical Point at $x = 0$

Critical Point at $x = 1$

4. Find absolute extremes of a continuous function on a closed interval

$$f(x) = \frac{x}{(x^2 + 9)^5}$$

$$f'(x) = \frac{-9x^2 + 9}{(x^2 + 9)^6}$$

$$[-2, 2]$$

$$f(-2) \approx -0.000005$$

$$f(2) \approx 0.000005$$

$$f(-1) = -0.00001$$

$$f(1) = 0.00001$$

Absolute Min at $(-1, f(-1))$

Absolute Max at $(1, f(1))$

5. Application of finding absolute extreme values

$$P(x) = 2x + \frac{128}{x}$$

$$P'(x) = 2 + \frac{-128}{x^2}$$

$$(0, \infty)$$

$$f(8) = 18$$

Absolute min at $(8, 32)$ or a perimeter of 32 units

Related Exercises

1. (Section 4.1, Exercise 11)

Absolute Min at $x = c_2$

Absolute Max at $x = b$

2. (Section 4.1, Exercise 14)

Absolute Min at $x = c$

Absolute Max at $x = b$

3. (Section 4.1, Exercise 15)

Absolute Max at $x = b$

Absolute Min at $x = a$

Local Max at $x = p$

Local Max at $x = r$

Local Min at $x = q$

Local Min at $x = s$

4. (Section 4.1, Exercise 18) Absolute Max at $x = p$

Absolute Min at $x = u$

Local Max at $x = p$

Local Max at $x = r$

Local Max at $x = t$

Local Min at $x = q$

Local Min at $x = s$

Local Min at $x = u$

5. (Section 4.1, Exercise 35)

$$f(x) = \frac{1}{x} + \ln x$$

$$f'(x) = \frac{x-1}{x^2}$$

Critical Points at $x = 1$

6. (Section 4.1, Exercise 36)

$$f(t) = t^2 - 2\ln(t^2 + 1)$$
$$f'(t) = \frac{2t(t+1)(t-1)}{t^2 + 1}$$

Critical Points at $t = -1$, $t = 0$ and $t = 1$

7. (Section 4.1, Exercise 46)

$$f(x) = x^4 - 4x^3 + 4x^2$$
$$f'(x) = 4x^3 - 12x^2 + 8x$$
$$[-1, 3]$$
$$f(-1) = 9$$
$$f(0) = 0$$
$$f(1) = 1$$
$$f(2) = 0$$
$$f(3) = 9$$

Absolute Max at $(-1, 9)$ and $(3, 9)$

Absolute Min at $(0, 0)$ and $(2, 0)$

8. (Section 4.1, Exercise 52)

$$f(x) = 3x^{\frac{2}{3}}$$
$$f'(x) = \frac{2}{x^{\frac{1}{3}}}$$
$$[0, 27]$$
$$f(0) = 0$$
$$f(27) = 27$$

Absolute Min at $(0, 0)$

Absolute Max at $(27, 27)$

9. (Section 4.1, Exercise 73)

$$s(t) = -16t^2 + 64t + 192$$
$$s'(t) = -32t + 64$$
$$0 \leq t \leq 6$$
$$s(0) = 192$$
$$s(2) = 256$$
$$s(6) = 0$$

The stone will reach its maximum height at 2 seconds