

Module 6 Notes (MATH-211)

Lillie Donato

15 July 2024

General Notes (and Definitions)

- L'Hôpital's Rule

Indeterminate Form: An expression involving two components where the limit cannot be determined by evaluating the limits of the individual components.

L'Hôpital's Rule: Suppose f and g are differentiable functions on an open interval I containing the point $x = a$, with $g'(x) \neq 0$ on I when $x \neq a$.

If $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$ has any of the indeterminate forms: $\frac{0}{0}$, $\frac{\infty}{\infty}$, $-\frac{\infty}{\infty}$, then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

provided that one of the following is the case:

$$\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \in \mathbb{R}$$

$$\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = \infty$$

$$\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = -\infty$$

L'Hôpital's Rule is still valid if $x \rightarrow a$ is replaced by any of $x \rightarrow a^+$, $x \rightarrow a^-$, $x \rightarrow \infty$, or $x \rightarrow -\infty$. In the last two of these cases, there must be a greatest x -value beyond which both f and g are differentiable at every point.

Exponential Indeterminate forms: 1^∞ , 0^0 , ∞^0

Method for evaluating limits of indeterminate forms 1^∞ , 0^0 , ∞^0 :

Assume that $L = \lim_{x \rightarrow a} f(x)^{g(x)}$ has one of these indeterminate forms.

1. Use the fact that the natural logarithm and natural exponential functions are inverses to write

$$L = \lim_{x \rightarrow a} e^{\ln(f(x)^{g(x)})}$$

2. Use the power property of logarithm arguments to write

$$L = \lim_{x \rightarrow a} e^{g(x) \ln(f(x))}$$

3. Use continuity of the exponential function to write

$$L = e^{\lim_{x \rightarrow a} g(x) \ln(f(x))}$$

4. Rewrite multiplication as division by the reciprocal:

$$L = e^{\lim_{x \rightarrow a} \left(\frac{\ln(f(x))}{\frac{1}{g(x)}} \right)}$$

5. Use L'Hôpital's Rule to evaluate this limit expression

Growth Rates: Suppose f and g are functions with $\lim_{x \rightarrow \infty} f(x) = \infty$ and $\lim_{x \rightarrow \infty} g(x) = \infty$

1. If one of the following are true, f **grows faster than** g , and we use the notation $f \gg g$

$$\lim_{x \rightarrow \infty} \frac{g(x)}{f(x)} = 0 \quad (1)$$

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \infty \quad (2)$$

2. f and g **have comparable growth rates**, if there is some non-zero finite number M such that

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = M$$

Ranked Growth Rates as $x \rightarrow \infty$

For any base $b > 1$, and for any positive numbers p , q , r , and s

$$\ln^q x \ll x^p \ll x^p \ln^r x \ll x^{p+s} \ll b^x \ll x^x$$

- Antiderivatives

Antiderivative: A function F is an antiderivative of another function f on an interval I if for all x in I :

$$F'(x) = f(x)$$

Family of Antiderivatives: Let $F(x)$ be any antiderivative of $f(x)$ on an interval I . Then all antiderivatives of f on I have the form $F(x) + C$, where C is an arbitrary constant.

Differential Equations: Any equation involving an unknown function and its derivatives

- Infinite family of solutions
- No two solutions from the family pass through the same point
- Given an initial condition $f(a) = b$, we can identify the particular family member that solves the given problem by solving for C

Antiderivative Rules

- Power Rule

If $p \neq -1$ and C is an arbitrary constant:

$$\int x^p dx = \frac{x^{p+1}}{p+1} + C$$

- Integral of x^{-1}

$$\int x^{-1} dx = \int \frac{1}{x} dx = \ln |x| + C$$

- Constant Multiple and Sum Rules

If $c \in \mathbb{R}$:

$$\int cf(x) dx = c \int f(x) dx$$

$$\int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx$$

- Integral of e^x

$$\int e^x dx = e^x + C$$

- Integral of $\frac{1}{x}$

$$\int \frac{1}{x} dx = \ln |x| + C$$

Trigonometric (and inverse) Integrals

$$\int \cos(x) dx = \sin x + C \quad (1)$$

$$\int \sin(x) dx = -\cos x + C \quad (2)$$

$$\int \sec^2(x) dx = \tan x + C \quad (3)$$

$$\int \csc^2(x) dx = -\cot x + C \quad (4)$$

$$\int \sec(x) \tan(x) dx = \sec x + C \quad (5)$$

$$\int \csc(x) \cot(x) dx = -\csc x + C \quad (6)$$

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C \quad (7)$$

$$\int \frac{1}{1+x^2} dx = \tan^{-1} x + C \quad (8)$$

$$\int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1} |x| + C \quad (9)$$

Examples

1. Use L'Hôpital's Rule to evaluate a limit with indeterminate form $\frac{0}{0}$

$$\lim_{x \rightarrow 0} \frac{e^x - x - 1}{5x^2} = \lim_{x \rightarrow 0} \frac{e^x - 1}{10x} \quad (1)$$

$$= \lim_{x \rightarrow 0} \frac{e^x}{10} \quad (2)$$

$$= \frac{e^0}{10} \quad (3)$$

$$= \frac{1}{10} \quad (4)$$

2. Use L'Hôpital's Rule to evaluate a limit with indeterminate form $\frac{\infty}{\infty}$

$$\lim_{x \rightarrow 0^+} \frac{1 - \ln x}{1 + \ln x} = \lim_{x \rightarrow 0^+} \frac{-\frac{1}{x}}{\frac{1}{x}} \quad (1)$$

$$= \lim_{x \rightarrow 0^+} \frac{-\frac{1}{x}}{\frac{1}{x}} \quad (2)$$

$$= \frac{-1}{1} \quad (3)$$

$$= -1 \quad (4)$$

3. Use L'Hôpital's Rule to evaluate a limit with indeterminate form $0 \cdot \infty$

$$\lim_{x \rightarrow 1^-} (1-x) \tan\left(\frac{\pi x}{2}\right) = \lim_{x \rightarrow 1^-} \frac{(1-x)}{\cot\left(\frac{\pi x}{2}\right)} \quad (1)$$

$$= \lim_{x \rightarrow 1^-} \frac{-1}{-\frac{\pi}{2} \csc^2\left(\frac{\pi x}{2}\right)} \quad (2)$$

$$= \lim_{x \rightarrow 1^-} \frac{2}{\pi} \sin^2\left(\frac{\pi x}{2}\right) \quad (3)$$

$$= \frac{2}{\pi} \quad (4)$$

4. Use L'Hôpital's Rule to evaluate a limit with exponential indeterminate form

$$\lim_{x \rightarrow 0^+} x^{\tan x} = e^{\lim_{x \rightarrow 0^+} \frac{\ln x}{\tan x}} \quad (1)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{\ln x}{\cot x}} \quad (2)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{1}{-x \csc^2 x}} \quad (3)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{-\sin^2 x}{x}} \quad (4)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{-2 \sin x \cos x}{1}} \quad (5)$$

$$= e^{\lim_{x \rightarrow 0^+} -2 \sin x \cos x} \quad (6)$$

$$= e^0 \quad (7)$$

$$= 1 \quad (8)$$

5. Compare the growth rates of functions

$$f(x) = x^2 \ln x$$

$$g(x) = x \ln^2 x$$

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{x^2 \ln x}{x \ln^2 x} \quad (1)$$

$$= \lim_{x \rightarrow \infty} \frac{x}{\ln x} \quad (2)$$

$$= \lim_{x \rightarrow \infty} \frac{1}{\frac{1}{x}} \quad (3)$$

$$= \lim_{x \rightarrow \infty} x \quad (4)$$

$$= \infty \quad (5)$$

Since $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \infty$, $f \gg g$

6. Use knowledge of derivatives to find antiderivatives

$$f(x) = -4 \cos x - x$$

$$F(x) = -4 \sin x - \frac{1}{2}x^2$$

$$F'(x) = -4 \cos x - x$$

$$\int (-4 \cos x - x) dx = -4 \sin x - \frac{1}{2}x^2 + C$$

7. Determine indefinite integrals using antiderivative rules

$$\int \frac{3}{x^4} + 2 - 3x^2 dx = \int 3x^{-4} + 2 - 3x^2 dx = \frac{-1}{x^3} + 2x - x^3 + C$$

8. Rewrite an indefinite integral to find an antiderivative

$$\int \frac{2 + 3 \cos y}{\sin^2 y} dy = \int 2 \csc^2 y + 3 \cot y \csc y dy \quad (1)$$

$$= 2 \int \csc^2 y dy + 3 \int \cot y \csc y dy \quad (2)$$

$$= -2 \cot y - 3 \csc y + C \quad (3)$$

9. Solve an initial value problem

$$f'(u) = 4 \cos u - 4 \sin u$$

$$f(\pi) = 0$$

$$f(u) = 4 \int \cos u du - 4 \int \sin u du = 4 \sin u + 4 \cos u + C$$

$$4 \sin \pi + 4 \cos \pi + C = 0 \quad (1)$$

$$0 - 4 + C = 0 \quad (2)$$

$$C = 4 \quad (3)$$

$$f(u) = 4 \sin u + 4 \cos u + 4$$

10. Application of differential equations to linear motion

$$a(t) = 2 + 3 \sin t$$

$$v(0) = 1$$

$$s(0) = 10$$

$$v(t) = \int 2 + 3 \sin t \, dt \quad (1)$$

$$= 2 \int t^0 \, dt + 3 \int \sin t \, dt \quad (2)$$

$$= 2t - 3 \cos t + C \quad (3)$$

$$2(0) - 3 \cos 0 + C = 1 \quad (4)$$

$$-3 + C = 1 \quad (5)$$

$$C = 4 \quad (6)$$

$$v(t) = 2t - 3 \cos t + 4 \quad (7)$$

$$s(t) = \int 2t - 3 \cos t + 4 \, dt \quad (8)$$

$$= 2 \int t \, dt - 3 \int \cos t \, dt + 4 \int t^0 \, dt \quad (9)$$

$$= 2 \frac{t^2}{2} - 3 \sin t + 4t + C \quad (10)$$

$$= t^2 - 3 \sin t + 4t + C \quad (11)$$

$$0^2 - 3 \sin 0 + 4(0) + C = 10 \quad (12)$$

$$C = 10 \quad (13)$$

$$s(t) = t^2 - 3 \sin t + 4t + 10 \quad (14)$$

Related Exercises

1. (Section 4.7, Exercise 17)

$$\lim_{x \rightarrow 2} \frac{x^2 - 2x}{x^2 - 6x + 8} = \lim_{x \rightarrow 2} \frac{2x - 2}{2x - 6} \quad (1)$$

$$= \frac{2(2) - 2}{2(2) - 6} \quad (2)$$

$$= \frac{4 - 2}{4 - 6} \quad (3)$$

$$= \frac{2}{-2} \quad (4)$$

$$= -1 \quad (5)$$

2. (Section 4.7, Exercise 18)

$$\lim_{x \rightarrow -1} \frac{x^4 + x^3 + 2x + 2}{x + 1} = \lim_{x \rightarrow -1} \frac{4x^3 + 3x^2 + 2}{1} \quad (1)$$

$$= \lim_{x \rightarrow -1} 4x^3 + 3x^2 + 2 \quad (2)$$

$$= 4(-1)^3 + 3(-1)^2 + 2 \quad (3)$$

$$= -4 + 3 + 2 \quad (4)$$

$$= 1 \quad (5)$$

3. (Section 4.7, Exercise 36)

$$\lim_{x \rightarrow 0} \frac{e^x - x - 1}{5x^2} = \lim_{x \rightarrow 0} \frac{e^x - 1}{10x} \quad (1)$$

$$= \lim_{x \rightarrow 0} \frac{e^x}{10} \quad (2)$$

$$= \frac{e^0}{10} \quad (3)$$

$$= \frac{1}{10} \quad (4)$$

4. (Section 4.7, Exercise 39)

$$\lim_{x \rightarrow 0} \frac{e^x - \sin x - 1}{x^4 + 8x^3 + 12x^2} = \lim_{x \rightarrow 0} \frac{e^x - \cos x}{4x^3 + 24x^2 + 24x} \quad (1)$$

$$= \lim_{x \rightarrow 0} \frac{e^x + \sin x}{12x^2 + 48x + 24} \quad (2)$$

$$= \frac{e^0 + \sin 0}{12(0)^2 + 48(0) + 24} \quad (3)$$

$$= \frac{1 + 0}{24} \quad (4)$$

$$= \frac{1}{24} \quad (5)$$

5. (Section 4.7, Exercise 38)

$$\lim_{x \rightarrow \infty} \frac{e^{3x}}{3e^{3x} + 5} = \lim_{x \rightarrow \infty} \frac{3e^{3x}}{9e^{3x}} \quad (1)$$

$$= \lim_{x \rightarrow \infty} \frac{1}{3} \cdot \frac{e^{3x}}{e^{3x}} \quad (2)$$

$$= \lim_{x \rightarrow \infty} \frac{1}{3} \quad (3)$$

$$= \frac{1}{3} \quad (4)$$

6. (Section 4.7, Exercise 51)

$$\lim_{x \rightarrow \infty} \frac{x^2 - \ln \frac{2}{x}}{3x^2 + 2x} = \lim_{x \rightarrow \infty} \frac{2x + \frac{1}{x}}{6x + 2} \quad (1)$$

$$= \lim_{x \rightarrow \infty} \frac{2 - \frac{1}{x^2}}{6} \quad (2)$$

$$= \frac{2 - 0}{6} \quad (3)$$

$$= \frac{2}{6} \quad (4)$$

$$= \frac{1}{3} \quad (5)$$

7. (Section 4.7, Exercise 53)

$$\lim_{x \rightarrow 0} x \csc x = \lim_{x \rightarrow 0} \frac{x}{\sin x} \quad (1)$$

$$= \lim_{x \rightarrow 0} \frac{1}{\cos x} \quad (2)$$

$$= \frac{1}{\cos 0} \quad (3)$$

$$= \frac{1}{1} \quad (4)$$

$$= 1 \quad (5)$$

8. (Section 4.7, Exercise 63)

$$\lim_{x \rightarrow \infty} \left(x^2 - \sqrt{x^4 + 16x^2} \right) = \lim_{x \rightarrow \infty} \left(x^2 - \sqrt{x^4 \left(1 + \frac{16}{x^2} \right)} \right) \quad (1)$$

$$= \lim_{x \rightarrow \infty} \left(x^2 - x^2 \sqrt{1 + \frac{16}{x^2}} \right) \quad (2)$$

$$= \lim_{x \rightarrow \infty} x^2 \left(1 - \sqrt{1 + \frac{16}{x^2}} \right) \quad (3)$$

$$= \lim_{x \rightarrow \infty} \frac{1 - \sqrt{1 + \frac{16}{x^2}}}{\frac{1}{x^2}} \quad (4)$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{16}{x^3}}{\frac{-2}{x^3} \sqrt{1 + \frac{16}{x^2}}} \quad (5)$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{16}{x^3} \cdot \frac{x^3}{-2}}{\sqrt{1 + \frac{16}{x^2}}} \quad (6)$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{16}{-2} \cdot \frac{x^3}{x^3}}{\sqrt{1 + \frac{16}{x^2}}} \quad (7)$$

$$= \lim_{x \rightarrow \infty} \frac{-8}{\sqrt{1 + \frac{16}{x^2}}} \quad (8)$$

$$= \frac{-8}{\sqrt{1+0}} \quad (9)$$

$$= \frac{-8}{1} \quad (10)$$

$$= -8 \quad (11)$$

9. (Section 4.7, Exercise 64)

$$\lim_{x \rightarrow \infty} \left(x - \sqrt{x^2 + 4x} \right) = \lim_{x \rightarrow \infty} \left(x - \sqrt{x^2 \left(1 + \frac{4}{x} \right)} \right) \quad (1)$$

$$= \lim_{x \rightarrow \infty} \left(x - x \sqrt{1 + \frac{4}{x}} \right) \quad (2)$$

$$= \lim_{x \rightarrow \infty} x \left(1 - \sqrt{1 + \frac{4}{x}} \right) \quad (3)$$

$$= \lim_{x \rightarrow \infty} \frac{1 - \sqrt{1 + \frac{4}{x}}}{\frac{1}{x}} \quad (4)$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{2}{x^2}}{\frac{-1}{x^2} \sqrt{1 + \frac{4}{x}}} \quad (5)$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{2}{x^2} \cdot \frac{x^2}{-1}}{\sqrt{1 + \frac{4}{x}}} \quad (6)$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{2}{-1} \cdot \frac{x^2}{x^2}}{\sqrt{1 + \frac{4}{x}}} \quad (7)$$

$$= \lim_{x \rightarrow \infty} \frac{-2}{\sqrt{1 + \frac{4}{x}}} \quad (8)$$

$$= \frac{-2}{\sqrt{1+0}} \quad (9)$$

$$= -2 \quad (10)$$

10. (Section 4.7, Exercise 75)

$$\lim_{x \rightarrow 0^+} x^{2x} = e^{\lim_{x \rightarrow 0^+} \frac{\ln x}{\frac{1}{2x}}} \quad (1)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{\frac{1}{x}}{\frac{-1}{2x^2}}} \quad (2)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{1}{x} \cdot \frac{2x^2}{-1}} \quad (3)$$

$$= e^{\lim_{x \rightarrow 0^+} -\frac{2x^2}{x}} \quad (4)$$

$$= e^{\lim_{x \rightarrow 0^+} -2x} \quad (5)$$

$$= e^{-2(0)} \quad (6)$$

$$= e^0 \quad (7)$$

$$= 1 \quad (8)$$

11. (Section 4.7, Exercise 76)

$$\lim_{x \rightarrow 0} (1 + 4x)^{\frac{3}{x}} = e^{\lim_{x \rightarrow 0^+} \frac{\ln(1+4x)}{\frac{1}{\frac{3}{x}}}} \quad (1)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{\ln(1+4x)}{\frac{x}{3}}} \quad (2)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{\frac{4}{(1+4x)}}{\frac{1}{3}}} \quad (3)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{4}{(1+4x)} \cdot \frac{3}{1}} \quad (4)$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{12}{(1+4x)}} \quad (5)$$

$$= e^{\frac{12}{1}} \quad (6)$$

$$= e^{12} \quad (7)$$

12. (Section 4.7, Exercise 96)

$$f(x) = x^2 \ln x \quad (1)$$

$$g(x) = \ln^2 x \quad (2)$$

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{x^2 \ln x}{\ln^2 x} \quad (1)$$

$$= \lim_{x \rightarrow \infty} \frac{x^2}{\ln x} \quad (2)$$

$$= \lim_{x \rightarrow \infty} \frac{2x}{\frac{1}{x}} \quad (3)$$

$$= \lim_{x \rightarrow \infty} 2x^2 \quad (4)$$

$$= \infty \quad (5)$$

$$f \gg g$$

13. (Section 4.7, Exercise 100)

$$f(x) = x^2 \ln x \quad (1)$$

$$g(x) = x^3 \quad (2)$$

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{x^2 \ln x}{x^3} \quad (1)$$

$$= \lim_{x \rightarrow \infty} \frac{\ln x}{x} \quad (2)$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{1}{x}}{1} \quad (3)$$

$$= \lim_{x \rightarrow \infty} \frac{1}{x} \quad (4)$$

$$= \frac{1}{\infty} \neq \infty \quad (5)$$

$$g \gg f$$

14. (Section 4.7, Exercise 95)

$$f(x) = x^{10} \quad (1)$$

$$g(x) = e^{0.01x} \quad (2)$$

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{x^{10}}{e^{0.01x}} \quad (1)$$

$$= \lim_{x \rightarrow \infty} \frac{10x^9}{0.01e^{0.01x}} \quad (2)$$

$$= \lim_{x \rightarrow \infty} \frac{90x^8}{0.01^2 e^{0.01x}} \quad (3)$$

$$= \lim_{x \rightarrow \infty} \frac{7200x^7}{0.01^3 e^{0.01x}} \quad (4)$$

$$= \lim_{x \rightarrow \infty} \frac{50400x^6}{0.01^4 e^{0.01x}} \quad (5)$$

$$= \lim_{x \rightarrow \infty} \frac{302400x^5}{0.01^5 e^{0.01x}} \quad (6)$$

$$= \lim_{x \rightarrow \infty} \frac{1512000x^4}{0.01^6 e^{0.01x}} \quad (7)$$

$$= \lim_{x \rightarrow \infty} \frac{6048000x^3}{0.01^7 e^{0.01x}} \quad (8)$$

$$= \lim_{x \rightarrow \infty} \frac{18144000x^2}{0.01^8 e^{0.01x}} \quad (9)$$

$$= \lim_{x \rightarrow \infty} \frac{36288000x}{0.01^9 e^{0.01x}} \quad (10)$$

$$= \lim_{x \rightarrow \infty} \frac{36288000}{0.01^{10} e^{0.01x}} \quad (11)$$

$$= \frac{36288000}{\infty} \neq \infty \quad (12)$$

$$g \gg f$$

15. (Section 4.7, Exercise 101)

$$f(x) = x^{20} \quad (1)$$

$$g(x) = 1.00001^x \quad (2)$$

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{x^{20}}{1.00001^x} \quad (1)$$

$$= \frac{2432902008176640000}{\infty} \neq \infty \quad (2)$$

$$g \gg f$$

16. (Section 4.9, Exercise 12)

$$f(x) = 11x^{10}$$

$$\int 11x^{10} dx = 11 \int x^{10} dx \quad (1)$$

$$= 11 \frac{x^{11}}{11} + C \quad (2)$$

$$= x^{11} + C \quad (3)$$

17. (Section 4.9, Exercise 13)

$$f(x) = 2 \sin x + 1$$

$$\int 2 \sin x + 1 dx = 2 \int \sin x + \int x^0 \quad (1)$$

$$= -2 \cos x + x + C \quad (2)$$

18. (Section 4.9, Exercise 24)

$$\int 3u^{-2} - 4u^2 + 1 \, du = 3 \int u^{-2} \, du - 4 \int u^2 \, du + \int u^0 \, du \quad (1)$$

$$= 3 \frac{u^{-1}}{-1} - 4 \frac{u^3}{3} + u + C \quad (2)$$

$$= -\frac{3}{u} - \frac{4u^3}{3} + u + C \quad (3)$$

19. (Section 4.9, Exercise 25)

$$\int 4\sqrt{x} - \frac{4}{\sqrt{x}} \, dx = 4 \int x^{\frac{1}{2}} \, dx - 4 \int x^{-\frac{1}{2}} \, dx \quad (1)$$

$$= 4 \frac{x^{\frac{3}{2}}}{\frac{3}{2}} - 4 \frac{x^{\frac{1}{2}}}{\frac{1}{2}} + C \quad (2)$$

$$= \frac{8x^{\frac{3}{2}}}{3} - 8x^{\frac{1}{2}} + C \quad (3)$$

$$= \frac{8x\sqrt{x}}{3} - 8\sqrt{x} + C \quad (4)$$

20. (Section 4.9, Exercise 31)

$$\int (3x + 1)(4 - x) \, dx = \int -3x^2 + 11x + 4 \, dx \quad (1)$$

$$= -3 \int x^2 \, dx + 11 \int x \, dx + 4 \int x^0 \, dx \quad (2)$$

$$= -3 \frac{x^3}{3} + 11 \frac{x^2}{2} + 4x + C \quad (3)$$

$$= -x^3 + \frac{11x^2}{2} + 4x + C \quad (4)$$

21. (Section 4.9, Exercise 35)

$$\int \frac{4x^4 - 6x^2}{x} \, dx = \int 4x^3 - 6x \, dx \quad (1)$$

$$= 4 \int x^3 \, dx - 6 \int x \, dx \quad (2)$$

$$= 4 \frac{x^4}{4} - 6 \frac{x^2}{2} + C \quad (3)$$

$$= x^4 - 3x^2 + C \quad (4)$$

22. (Section 4.9, Exercise 40)

$$\int (\csc^2 \theta + 1) \, d\theta = \int \csc^2 \theta \, d\theta + \int \theta^0 \, d\theta \quad (1)$$

$$= -\cot \theta + \theta + C \quad (2)$$

23. (Section 4.9, Exercise 41)

$$\int \frac{2 + 3 \cos y}{\sin^2 y} \, dy = \int \left(\frac{2}{\sin^2 y} + \frac{3 \cos y}{\sin^2 y} \right) \, dy \quad (1)$$

$$= \int (2 \csc^2 y + 3 \cot y \csc y) \, dy \quad (2)$$

$$= 2 \int \csc^2 y \, dy + 3 \int \cot y \csc y \, dy \quad (3)$$

$$= -2 \cot y - 3 \csc y + C \quad (4)$$

24. (Section 4.9, Exercise 47)

$$\int (3t^2 + 2 \csc^2 t) dt = \int 3t^2 dt + \int 2 \csc^2 t dt \quad (1)$$

$$= 3 \int t^2 dt + 2 \int \csc^2 t dt \quad (2)$$

$$= 3 \frac{t^3}{3} - 2 \cot t + C \quad (3)$$

$$= t^3 - 2 \cot t + C \quad (4)$$

25. (Section 4.9, Exercise 45)

$$\int (\sec^2 \theta + \sec \theta \tan \theta) d\theta = \int \sec^2 \theta d\theta + \int \sec \theta \tan \theta d\theta \quad (1)$$

$$= \tan \theta + \sec \theta + C \quad (2)$$

26. (Section 4.9, Exercise 50)

$$\int \frac{\csc^3 x + 1}{\csc x} dx = \int \frac{\csc^3 x}{\csc x} + \frac{1}{\csc x} dx \quad (1)$$

$$= \int \csc^2 x + \sin x dx \quad (2)$$

$$= \int \csc^2 x dx + \int \sin x dx \quad (3)$$

$$= -\cot x + -\cos x + C \quad (4)$$

27. (Section 4.9, Exercise 51)

$$\int \frac{1}{2y} dy = \frac{1}{2} \int \frac{1}{y} dy \quad (1)$$

$$= \frac{1}{2} \ln |y| + C \quad (2)$$

$$= \frac{\ln |y|}{2} + C \quad (3)$$

28. (Section 4.9, Exercise 53)

$$\int \frac{6}{\sqrt{4-4x^2}} dx = \int \frac{6}{\sqrt{4(1-x^2)}} dx \quad (1)$$

$$= \int \frac{6}{2\sqrt{1-x^2}} dx \quad (2)$$

$$= 3 \int \frac{1}{\sqrt{1-x^2}} dx \quad (3)$$

$$= 3 \sin^{-1} x + C \quad (4)$$

29. (Section 4.9, Exercise 59)

$$\int \frac{t+1}{t} dt = \int \left(\frac{t}{t} + \frac{1}{t} \right) dt \quad (1)$$

$$= \int \frac{t}{t} dt + \int \frac{1}{t} dt \quad (2)$$

$$= \int t^0 dt + \ln |t| + C \quad (3)$$

$$= t + \ln |t| + C \quad (4)$$

30. (Section 4.9, Exercise 61)

$$\int e^{x+2} dx = \int e^2 e^x dx \quad (1)$$

$$= e^2 \int e^x dx \quad (2)$$

$$= e^2 e^x \quad (3)$$

$$= e^{x+2} \quad (4)$$

31. (Section 4.9, Exercise 54)

$$\int \frac{v^3 + v + 1}{1 + v^2} dv = \int \frac{v^3 + v}{1 + v^2} + \frac{1}{1 + v^2} dv \quad (1)$$

$$= \int v + \frac{1}{1 + v^2} dv \quad (2)$$

$$= \int v dv + \int \frac{1}{1 + v^2} dv \quad (3)$$

$$= \frac{v^2}{2} + \tan^{-1} v + C \quad (4)$$

32. (Section 4.9, Exercise 62)

$$\int \frac{10t^5 - 3}{t} dt = \int \left(\frac{10t^5}{t} - \frac{3}{t} \right) dt \quad (1)$$

$$= \int 10t^4 dt - \int \frac{3}{t} dt \quad (2)$$

$$= 10 \int t^4 dt - 3 \int \frac{1}{t} dt \quad (3)$$

$$= 10 \frac{t^5}{5} - 3 \ln |t| + C \quad (4)$$

$$= 2t^5 - 3 \ln |t| + C \quad (5)$$

33. (Section 4.9, Exercise 78)

$$g'(x) = 7x^6 - 4x^3 + 12 \quad (1)$$

$$g(1) = 24 \quad (2)$$

$$g(x) = \int 7x^6 - 4x^3 + 12 dx \quad (3)$$

$$= 7 \int x^6 dx - 4 \int x^3 dx + 12 \int x^0 dx \quad (4)$$

$$= 7 \frac{x^7}{7} - 4 \frac{x^4}{4} + 12x + C \quad (5)$$

$$= x^7 - x^4 + 12x + C \quad (6)$$

$$(1)^7 - (1)^4 + 12(1) + C = 24 \quad (7)$$

$$1 - 1 + 12 + C = 24 \quad (8)$$

$$12 + C = 24 \quad (9)$$

$$C = 12 \quad (10)$$

$$g(x) = x^7 - x^4 + 12x + 12 \quad (11)$$

34. (Section 4.9, Exercise 83)

$$y'(t) = \frac{3}{t} + 6 \quad (1)$$

$$y(1) = 8, t > 0 \quad (2)$$

$$y(t) = \int \left(\frac{3}{t} + 6 \right) dt \quad (3)$$

$$= 3 \int \frac{1}{t} dt + 6 \int t^0 dt \quad (4)$$

$$= 3 \ln |t| + 6t + C \quad (5)$$

$$3 \ln 1 + 6(1) + C = 8 \quad (6)$$

$$6 + C = 8 \quad (7)$$

$$C = 2 \quad (8)$$

$$y(t) = 3 \ln |t| + 6t + 2 \quad (9)$$

35. (Section 4.9, Exercise 105)

$$v(t) = \sin t \quad (1)$$

$$s(0) = 0 \quad (2)$$

$$s(t) = \int \sin t \, dt \quad (3)$$

$$= -\cos t + C \quad (4)$$

$$-\cos 0 + C = 0 \quad (5)$$

$$-1 + C = 0 \quad (6)$$

$$C = 1 \quad (7)$$

$$s(t) = -\cos t + 1 \quad (8)$$

$$V(t) = \cos t \quad (1)$$

$$S(0) = 0 \quad (2)$$

$$S(t) = \int \cos t \, dt \quad (3)$$

$$= \sin t + C \quad (4)$$

$$\sin 0 + C = 0 \quad (5)$$

$$C = 0 \quad (6)$$

$$S(t) = \sin t \quad (7)$$

36. (Section 4.9, Exercise 106)

$$v(t) = e^t \quad (1)$$

$$s(0) = 0 \quad (2)$$

$$s(t) = \int e^t \, dt \quad (3)$$

$$= e^t + C \quad (4)$$

$$e^0 + C = 0 \quad (5)$$

$$1 + C = 0 \quad (6)$$

$$C = -1 \quad (7)$$

$$s(t) = e^t - 1 \quad (8)$$

$$V(t) = 2 + \cos t \quad (1)$$

$$S(0) = 3 \quad (2)$$

$$S(t) = \int 2 + \cos t \, dt \quad (3)$$

$$= \int 2 \, dt + \int \cos t \, dt \quad (4)$$

$$= 2 \int 1 \, dt + \sin t + C \quad (5)$$

$$= 2t + \sin t + C \quad (6)$$

$$2(0) + \sin 0 + C = 3 \quad (7)$$

$$C = 3 \quad (8)$$