

The distribution of Genus *Neotrygon* around the world based on BoldSystem database

Onny Nurrahman Marwayana

Introduction

A genus of stingrays in the family of Dasyatidae is *Neotrygon*, which is mostly distributed in the Indo-West Pacific area. This genus is also called as the Maskrays since it has a special color pattern around the eyes, considering as a mask. *Neotrygon kuhlii* (Figure 1) is a species in this genus. Many researchers originally put the species under this genus in the genus of *Dasyatis*. However, some morphological and molecular analysis in taxonomy studies showed that the genus was supposed to be splitted into another genus so that genus *Neotrygon* was created (Last & White, 2008).

To do both analysis, researchers really need to observe and refer to the holotype and paratype of the samples that actually have been preserved in some reference collections, such as Australian National Fish Collection - CSIRO. There is a web page called **BoldSystem**¹, which provides some species database of living organisms around the world which have been collected, recorded, and preserved at some reference collections. The Barcode of Life Data Systems (BOLD) is definitely an informatics workbench aiding the acquisition, storage, analysis, and publication of DNA barcode records. By assembling molecular, morphological, and distributional data, it bridges a traditional bioinformatics chasm (Ratnasingham & Hebert, 2007).

The purpose of this project are to collect some information about this genus regarding the database of genus *Neotrygon* downloaded from BoldSystem and use it to answer some research questions: 1). how many institutions have stored the specimens, 2). how many specimens are stored in each institution, 3). where the specimens mostly have been stored. Finally, we will make a dictionary that contains the answer of the questions. At the end of this project, we will get the information regarding the collections of genus *Neotrygon* in BoldSystem database.

¹<http://www.boldsystems.org/index.php/databases>



2

²<https://en.wikipedia.org/wiki/Maskray>

Material and Methods

The material that we use in this project is a specimen database of genus *Neotrygon*, which has been downloaded from BoldSystem. The database has been created and published by some researchers (Cerutti-Pereyra et al., 2012; Borsa et al., 2013; Pavan-Kumar et al., 2013) By using some commands in UNIX shell and iPython, we manipulate the database to achieve the purpose. The commands are showed below.

These are some commands in UNIX shell to answer the questions:

```
cd /Desktop/eeb-177/eeb-174-final-project/final_project_upload
ls
cut -f 6 CP_NK_bold.txt | tail -n +2 | sort | uniq | wc -l
cut -f 6 CP_NK_bold.txt | tail -n +2 | sort | uniq
cut -f 6 CP_NK_bold.txt | tail -n +2 | sort | uniq -c
cut -f 21 CP_NK_bold.txt | tail -n +2 | sort | uniq | wc -l
cut -f 21 CP_NK_bold.txt | tail -n +2 | sort | uniq -c
touch new_database.txt #Make a new .txt file
#finally, end up the information on the new file
```

And this is some commands in iPython to create the dictionary:

```
neotrygon = open("../final_project_upload/formatted_bold_data.csv")
 #(for a complete directory address, please look above at cd /Desktop ...)
neotrygon.readline()
neotrygon_dict = {}
for line in neotrygon:
    species = line.split(",")[20]
    institution = line.split(",")[5]
    country = line.split(",")[37]
    values = [institution, country]
    neotrygon_dict[species] = values
print(neotrygon_dict)
```

Result and Discussion

By running the commands on UNIX shell, we get some information explained below:

```
## bash: line 0: cd: /Desktop/eeb-177/eeb-174-final-project/final_project_upload: No such file or directory
## 4
## Biodiversity Institute of Ontario
## Central Institute of Fisheries Education
## CSIRO, Australian National Fish Collection
## Mined from GenBank, NCBI
##      11 Biodiversity Institute of Ontario
##       1 Central Institute of Fisheries Education
##      40 CSIRO, Australian National Fish Collection
##      32 Mined from GenBank, NCBI
## 7
##      3 Neotrygon annotata
##      5 Neotrygon cf. annotata
##     56 Neotrygon kuhlii
##      8 Neotrygon leylandi
##      3 Neotrygon ningalooensis
##      8 Neotrygon picta
##      1 Neotrygon trigonoides
```

The result explains that there are 4 institutions around the world which have preserved some specimens of genus *Neotrygon*. Biodiversity Institute of Ontario stored 11 specimens. Central Institute of Fisheries Education preserved one specimen, while CSIRO - Australian National Fish Collection and GenBank-NCBI saved 40 and 32 specimens. Based on the result, we know that the specimens of *Neotrygon* from around the world mostly have been preserved at the Australian National Fish Collection - CSIRO. This information indicates that we should refer to the collection preserved at the institution if we want to study the conventional taxonomy of genus *Neotrygon* and analyze the morphological characters of the genus. Moreover, the holotypes and paratypes of the genus have probably preserved at the institution.

In addition, there are 85 specimens of 7 species that have been recorded on the BoldSystem web page. The species are *N. annotata*, *N. cf. annotata*, *N. kuhlii*, *N. leylandi*, *N. ningalooensis*, *N. picta*, and *N. trigonoides*. Based on the database, *N. kuhlii* is the most popular species of genus *Neotrygon*, which was collected around the world. This information describes that the most distributed species of genus *Neotrygon* is *N. kuhlii*. Puckridge and colleagues state that *N. kuhlii* has been broadly distributed in Japan, the South East Asia region, Northern Australia, and the Western coast of Africa (Puckridge et al., 2013).

```
## {'Neotrygon ningalooensis': ['Mined from GenBank, NCBI', 'Australia'], 'Neotrygon leylandi': ['Central Institute of Fisheries Education', 'Australia']}
```

This is a dictionary constructed from the database using Python program. Based on the dictionary, every single specimen of genus *Neotrygon* has been collected from a specific country and preserved at a specific reference collection. For example, the specimens of

N. kuhlii were mostly collected from India, while *N. trigonoides* were collected from New Caledonia. In South East Asia region, Indonesia represents the region that contributes some specimens of *N. cf. annotata*. This information truly indicates that the species in genus *Neotrygon* has been vastly distributed from the South Pacific region until Indian Ocean. That is like what Puckridge and colleagues state in their article (Puckridge et al., 2013).

Conclusion

Based on the data set, there are 85 specimens of 7 species in genus *Neotrygon*, which have been preserved at 4 reference collections around the world; they are Biodiversity Institute of Ontario, Central Institute of Fisheries Education, Australian National Fish Collection - CSIRO, and GenBank - NCBI. The highest species number of genus *Neotrygon* based on BoldSystem database is *N. kuhlii*, which were mostly collected from India.

Furthermore, we can manipulate some database to answer some questions that we were wondering to know by using some commands and functions in **UNIX shell** and **iPython**. This article also was created by using **R markdown** in **R studio**. Therefore, some computer programming could ease some researcher in manipulating their data to achieve the target of their research.

GitHub address

This is my GitHub address for final project: https://github.com/onnysmu3/eeb-174-final-project/tree/master/FIX_final_project_upload

References

- Borsa, P., Arlyza, I. S., Chen, W.-J., Durand, J.-D., Meekan, M. G. & Shen, K.-N. (2013) Resurrection of New Caledonian Maskray Neotrygon Trigonoides (Myliobatoidei: Dasyatidae) from Synonymy with *N. kuhlii*, Based on Cytochrome-Oxidase I Gene Sequences and Spotting Patterns. *Comptes Rendus Biologies* **336**, 221–232.
- Cerutti-Pereyra, F., Meekan, M. G., Wei, N.-W. V., Shea, O. O., Bradshaw, C. J. A. & Austin, C. M. (2012) Identification of Rays Through DNA Barcoding: An Application for Ecologists. *PLoS ONE* **7**, e36479.
- Last, P. & White, W. (2008) Resurrection of the Genus Neotrygon Castelnau (Myliobatoidei: Dasyatidae) with the Description of Neotrygon Picta Sp. Nov., a New Species from Northern Australia. *Descriptions of New Australian Chondrichthyans: CSIRO Marine and Atmospheric Research Paper* 315–325.
- Pavan-Kumar, A., Gireesh-Babu, P., Babu, P. P. S., Jaiswar, A. K., Prasad, K. P., Chaudhari, A., Raje, S. G., Chakraborty, S. K., Krishna, G. & Lakra, W. S. (2013) DNA Barcoding of Elasmobranchs from Indian Coast and Its Reliability in Delineating Geographically Widespread Specimens. *Mitochondrial DNA* **26**, 92–100.
- Puckridge, M., Last, P. R., White, W. T. & Andreakis, N. (2013) Phylogeography of the Indo-West Pacific Maskrays (Dasyatidae, Neotrygon): A Complex Example of Chondrichthyan Radiation in the Cenozoic. *Ecology and evolution* **3**, 217–232.
- Ratnasingham, S. & Hebert, P. D. (2007) BOLD: The Barcode of Life Data System ([Http://Www. Barcodinglife. Org](http://www.barcodinglife.org)). *Molecular ecology notes* **7**, 355–364.