1

Assignment 5

R.OOHA

Download all python codes from

https://github.com/ooharapolu/ASSIGNMNT5/ Assignment5.py

and latex-tikz codes from

https://github.com/ooharapolu/ASSIGNMNT5/main .tex

1 Question No.2.158

Find the area lying in the first quadrant and by the circle $\mathbf{x}^T \mathbf{x} = \mathbf{4}$ and the lines $\mathbf{x} = \mathbf{0}$ and $\mathbf{x} = \mathbf{2}$.

2 Solution

The given lines of the vector form is,

$$\begin{pmatrix} 1 & 0 \end{pmatrix} \mathbf{x} = 0 \tag{2.0.1}$$

$$\begin{pmatrix} 1 & 0 \end{pmatrix} \mathbf{x} = 2 \tag{2.0.2}$$

The general equation of the circle is,

$$\mathbf{x}^T \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + \mathbf{f} = 0 \tag{2.0.3}$$

$$\|\mathbf{x}\|^2 + 2\mathbf{u_1}^T\mathbf{x} + \mathbf{f_1} = 0$$
 (2.0.4)

$$\mathbf{x}^T \mathbf{x} - \mathbf{4} = 0 \tag{2.0.5}$$

$$\mathbf{u_1} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{f_1} = -\mathbf{4} \tag{2.0.6}$$

To find points A and B The parametric form is,

$$\mathbf{B} = \mathbf{q} + \lambda \mathbf{m} \tag{2.0.7}$$

$$= \begin{pmatrix} 0 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 1 \end{pmatrix} \tag{2.0.8}$$

From the intersection of circle and line, the value of λ can be found by,

$$\lambda^{2} = \frac{-\mathbf{f}_{1} - ||\mathbf{q}||^{2}}{||\mathbf{m}||^{2}} = \frac{4 - 0}{1} = 4$$
 (2.0.9)

$$\implies \lambda = \pm 2$$
 (2.0.10)

They are parallel to y-axis. So they make 90 degrees Point of intersection Sub equation (2.0.10) in (2.0.8) Put, $\lambda = 2$

$$\mathbf{B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + 2 \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 2 \end{pmatrix} \tag{2.0.11}$$

$$\implies \mathbf{B} = \begin{pmatrix} 0 \\ 2 \end{pmatrix} \tag{2.0.12}$$

and, $\lambda = -2$

$$\mathbf{B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ -2 \end{pmatrix} \tag{2.0.13}$$

$$\implies \mathbf{B} = \begin{pmatrix} 0 \\ -2 \end{pmatrix} \tag{2.0.14}$$

As given in question as first quadrant

$$\implies \mathbf{B} = \begin{pmatrix} 0 \\ 2 \end{pmatrix} \tag{2.0.15}$$

Similarly, to find point A, The parametric form of the line is,

$$\mathbf{A} = \mathbf{q} + \lambda \mathbf{m} \tag{2.0.16}$$

$$= \begin{pmatrix} 2\\0 \end{pmatrix} + \lambda \begin{pmatrix} 0\\1 \end{pmatrix} \tag{2.0.17}$$

$$\lambda^{2} = \frac{-\mathbf{f_{1}} - ||\mathbf{q}||^{2}}{||\mathbf{m}||^{2}}$$
 (2.0.18)

$$=\frac{4-4}{1}=0\tag{2.0.19}$$

$$\implies \lambda = 0 \tag{2.0.20}$$

Sub (2.0.20) in (2.0.17)

$$\mathbf{A} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} + 0 \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \tag{2.0.21}$$

$$\implies \mathbf{A} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \tag{2.0.22}$$

The equation of a circle can be expressed as,

$$\mathbf{x}^T \mathbf{x} - 2\mathbf{u}^T \mathbf{x} + \mathbf{f} = 0 \tag{2.0.23}$$

comparing equation (??) with the circle equation given,

$$\mathbf{x}^T \mathbf{x} = \mathbf{4} \tag{2.0.24}$$

$$\implies \mathbf{u} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{f} = -\mathbf{4} \tag{2.0.25}$$

$$\implies \mathbf{r} = \sqrt{\mathbf{u}^T \mathbf{x} - \mathbf{f}} = \sqrt{4} \tag{2.0.26}$$

$$\implies r = 2 \tag{2.0.27}$$

Using equation (2.0.27) and, the area of the sector is obtained as,

$$\implies \frac{\theta}{360^{\circ}} \pi \mathbf{r}^2 = \frac{90^{\circ}}{360^{\circ}} \pi \left(2\right)^2 = \pi \qquad (2.0.28)$$

Plot of the given -




Fig. 2.1: The Circle of the lines