

百年京张 AI 创新带 · 空间结构总图

主脊（京张铁路遗址公园）· 三带 · 两

北翼 · 研发孵化翼（清华·北大）

N10
绿楔缝合点

N7
信号塔

N6
知春里

创新带范围示意 43.6 km²

南翼 · 成果转化与展示翼（至北京北站）

坐标基于真实地名近似（provisional），非官方红线；官方数据发布

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported to be the most common serotype of *Shigella* isolated from children with shigellosis [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1980s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [12]. In the 1990s, *S. flexneri* was the most common serotype of *Shigella* isolated from children with shigellosis in the United Kingdom [13].

The aim of this study was to determine the prevalence of *S. flexneri* in children with shigellosis in the United Kingdom. The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high.

Methods

Study area

The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high.

Study design

The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high.

Study population

The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high.

Study procedures

The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high.

Study results

The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high. The study was conducted in the United Kingdom, where the incidence of shigellosis is high.