## Reconciliation of results and studies included.

	Current (August, 2020) PMID <u>32805168</u>	<b>Giustozzi, May 2020</b> PMID <u>31104194</u>	Fuentes, 2019 <sup>a</sup> PMID <u>31685262</u>	Wongsaengsak, 2019 DOI <u>10.1182/blood-</u> <u>2019-125473</u>	<b>Brunetti, 2019</b> PMID <u>31654194</u>
Outcome(s)					
		Venous thromboem	bolism recurrence		
Effect size	OR: 0.59 (0.41-0.86)	RR: 0.62 (0.43-0.91)	OR: 0.48 (0.24-0.96)	RR: 0.48 (0.26-0.87)	RR: 0.56 (0.40-0.79)
Heterogeneity	25%	30%	46%	56%	59%
		Major bl	eeding	<u>.                                      </u>	
Effect size	OR: 1.34 (0.83-2.18)	RR: 1.31 (0.83-2.08)	OR: 1.70 (1.04-2.78)	RR: 1.55 (0.79-3.04)	RR: 1.53 (0.95-2.47)
Heterogeneity	28%	23%	2%	29%	28%
Summary of important differences or findings	Included findings at 12 months for HOKUSAI VTE trial.				
Studies included	<u> </u>			·	
Raskob (HOKUSAI-VTE), 2018 PMID 29231094	Yes*	Yes	Yes	Yes	Yes
Young (SELECT- D), 2018 PMID <u>29746227</u>	Yes	Yes	Yes	Yes	Yes
McBane (ADAM-VTE), 2019 PMID <u>28837207</u>	Yes	Yes	Yes	Yes	Yes
Agnelli (CaraVaggio), 2020 PMID <u>32223112</u> Notes:	Yes	Yes	NA	NA	NA

<sup>\*</sup> Outcomes reported at 12 months for HOKUSAI VTE trial were used in the current analysis.