



## Ecological Approaches to Pest Management



P Srikala<sup>1\*</sup> and R Saranya<sup>2</sup>

<sup>1\*2</sup>Department of Agricultural Entomology, Tamil Nadu Agricultural University,  
Coimbatore - 641003, Tamil Nadu, India.

<sup>1\*</sup>Corresponding author E-mail: [srikala1104@gmail.com](mailto:srikala1104@gmail.com)

### Abstract

Ecological approaches to pest management are grounded in population ecology and trophic interactions that regulate insect pests within agro-ecosystems. These approaches exploit bottom-up factors such as host plant quality and cropping patterns, and top-down forces including predators, parasitoids, and entomopathogens, to maintain pest densities below the economic injury level. Cultural practices disrupt pest colonization and continuity, while botanical insecticides act through antifeedant, growth-regulatory, and oviposition-deterrent mechanisms with low non-target effects. Biological control provides density-dependent suppression mediated by natural enemy dynamics, further strengthened through habitat manipulation and resource provisioning. Integrated with pest monitoring and threshold-based decision making, ecological strategies enhance system stability, delay resistance evolution, and promote sustainable crop protection under changing climatic conditions.

### Introduction

Ecological approaches to pest management are founded on the understanding that insect pests are regulated by interactions among host plants, natural enemies, climate, and agricultural practices. These approaches aim to manipulate agro-ecosystem processes so that pest populations remain below the economic injury level (EIL), rather than achieving complete elimination. Such strategies reduce pesticide dependence, delay resistance development, and conserve ecosystem services.

### Ecological principles underlying pest management

#### Pest population regulation

Pest populations are governed by

- Bottom-up regulation (host plant nutrition, crop phenology)
- Top-down regulation (predators, parasitoids, pathogens)
- Environmental modulation (temperature, humidity)
- Anthropogenic interventions (cropping pattern, pesticide use)

Eg: Nitrogen-rich crops favour aphid multiplication, while predator-rich systems suppress aphid outbreaks.

#### Biodiversity-stability relationship

Agro-ecosystems with greater species and functional diversity exhibit:

- Lower pest outbreak frequency
- Greater resilience to climatic variability

Eg.: Polycropped vegetable systems show reduced whitefly pressure compared to monocultures.

#### Cultural methods as ecological pest management tools

##### Manipulation of crop phenology

- **Synchronized planting** reduces availability of vulnerable stages

*Example:* Community rice planting reduces brown planthopper (*Nilaparvata lugens*) population buildup.

**Host availability disruption**

- **Crop rotation** prevents pest carryover

*Example:* Cotton-non-host cereal rotation suppresses *Helicoverpa armigera*.

**Habitat diversification**

- **Intercropping** interferes with pest orientation and enhances enemies

*Example:* Maize-legume intercropping reduces stem borer incidence.

**Behavioural manipulation**

- **Trap cropping** exploits pest preference

*Example:* Marigold attracts *Helicoverpa* away from tomato.

**Botanical pesticides in ecological pest management****Mode of action in ecological context**

Botanicals act mainly as:

- Antifeedants
- Insect growth regulators
- Oviposition deterrents

**Crop-specific examples**

- Neem: disrupts moulting and fecundity in *S. litura* on groundnut
- Pongamia oil: reduces feeding in aphids and jassids in vegetables
- Pyrethrum: rapid knockdown of flies with minimal persistence

**Biological control as an ecological regulation mechanism****Predators (Generalist Regulation)**

- *Coccinella septempunctata* → aphids in wheat
- *Chrysoperla carnea* → thrips and whiteflies in cotton

**Parasitoids (Density-Dependent Suppression)**

- *Trichogramma chilonis* → eggs of *Helicoverpa armigera*
- *Bracon hebetor* → larvae of stored product moths

**Entomopathogens (Epizootic Control)**

- *Beauveria bassiana* → whiteflies in cotton
- *Metarhizium anisopliae* → termites and soil pests

These agents operate through host–pathogen dynamics, especially under favourable microclimates.

**Habitat manipulation and conservation biological control****Resource provisioning**

- Flowering plants supply nectar and pollen to parasitoids

*Example:* Buckwheat strips increase parasitoids longevity.

**Structural refugia**

- Bund vegetation and reduced tillage conserve predators

*Example:* Spiders in rice ecosystems reduce leafhopper populations.

**Ecological approaches under climate change**

Climate change increases pest voltinism and range expansion. Ecological strategies buffer these impacts by

- Maintaining enemy diversity
- Enhancing ecosystem redundancy

*Example:* Diversified cotton systems show stable pest–enemy ratios despite fluctuating temperatures.



Integration with IPM Framework

Ecological approaches form the preventive foundation of IPM, complemented by

- Pest surveillance
- Economic threshold-based interventions
- Selective insecticides when necessary

Table 1: Ecological Pest Management Approaches - Sub-Topics and Examples

| Main Approach      | Sub-topic        | Ecological Function     | Example                        |
|--------------------|------------------|-------------------------|--------------------------------|
| Cultural           | Crop rotation    | Host discontinuity      | Cotton vs <i>Helicoverpa</i>   |
| Cultural           | Intercropping    | Habitat diversification | Maize–cowpea vs stem borer     |
| Cultural           | Trap cropping    | Behavioural diversion   | Marigold–tomato                |
| Botanical          | Neem products    | Growth regulation       | <i>Spodoptera</i> in groundnut |
| Botanical          | Plant oils       | Feeding deterrence      | Aphids in vegetables           |
| Biological         | Predators        | Continuous suppression  | Ladybird vs aphids             |
| Biological         | Parasitoids      | Reproductive inhibition | <i>Trichogramma</i> vs eggs    |
| Habitat management | Flowering strips | Enemy conservation      | Cotton ecosystems              |

Conclusion

Ecological approaches to pest management emphasize ecosystem regulation rather than chemical domination. By strengthening natural control mechanisms, these approaches ensure long-term pest suppression, environmental safety, and agricultural sustainability.

References

Altieri, M.A. 1999. The ecological role of biodiversity in agroecosystems. *Agriculture, Ecosystems & Environment* 74: 19–31.

Dent, D. 2000. *Insect Pest Management*. 2nd ed. CABI Publishing, UK.

Kogan, M. 1998. Integrated pest management: Historical perspectives and contemporary developments. *Annual Review of Entomology* 43: 243–270.

Metcalf, R.L., and W.H. Luckmann. 1994. Introduction to *Insect Pest Management*. 3rd ed. Wiley, New York.

Pedigo, L.P., and M.E. Rice. 2009. *Entomology and Pest Management*. 6th ed. Pearson Education.