



Red Palm Mite (*Raoiella indica*): A Tiny Invader Threatening Global Agriculture

Parvati, B. Morabad^{1*} and Kishore S.M¹

^{1*,1}Department of Entomology, College of Agriculture, Shivamogga Keladi Shivappa Nayaka University of Agricultural and Horticultural Sciences, Shivamogga

Corresponding author: *Email I.D.: parvatimorabad17@gmail.com

Abstract

The red palm mite, *Raoiella indica* Hirst (Acari: Tenuipalpidae), is a tiny arthropod that has emerged as a major invasive pest, affecting economically important crops like coconut, arecanut, banana and ornamental palms across tropical and subtropical regions. Native to Southern Asia, this mite’s global spread has been facilitated by human activities and favourable environmental conditions. Its specialized feeding through stomata disrupts photosynthesis and respiration, leading to reduced productivity and significant economic losses. This article reviews its morphology, life cycle, feeding habits, host range, natural enemies and management strategies, emphasizing the importance of integrated pest control and farmer awareness, particularly in India.

Keywords *Raoiella indica*, red palm mite, invasive pest, Arecanut, coconut, host range, morphology and biological control.

1. Introduction

Mites are tiny arthropods that can inflict severe damage to crops worldwide. Among them, the red palm mite (*Raoiella indica*) is a particularly destructive pest, originally described in Coimbatore, India, on coconut (*Cocos nucifera*) by Hirst in 1924. It remained restricted to parts of Asia, Africa and the Middle East until its sudden appearance in the Caribbean in 2004 (Flechtmann and Etienne, 2004). Since then, it has rapidly expanded across the Americas and Africa, affecting vital agricultural zones.

Raoiella indica is a polyphagous pest with a wide host range that includes coconut, arecanut (*Areca catechu*), banana (*Musa* spp.), date palm, heliconia and several ornamental palms. In India, its infestation on arecanut leaves, particularly in young plantations during hot and humid seasons, has been increasingly reported (Patel and Rao, 1958; Chavan *et al.*, 2020).

Its notoriety stems from its invasive potential and the ecological damage it causes. It feeds through stomata, interfering with photosynthesis and respiration, leading to stunted growth and premature leaf fall

(Ochoa *et al.*, 2011). Human activities such as international trade and transport of plant materials have accelerated its spread, making quarantine and integrated management essential in affected regions like India.

2. What Does *Raoiella indica* Look Like? (Morphology)

The red palm mite is easily identifiable under magnification due to its striking red colour and oval shape. The adult female measures 0.23 to 0.30 mm in length and 0.17 to 0.19 mm in width, with smooth dorsal surfaces punctuated by sculptured depressions (Chavan *et al.*, 2020). Males are slightly smaller, ranging from 0.15 to 0.20 mm long and tapering at the posterior end.

Eggs are smooth, ovoid, orange to reddish and adhere to the underside of leaves with a slender hair-like structure. Newly hatched larvae are reddish with three pairs of legs and dark posterior markings after feeding. The protonymph and deutonymph stages develop four pairs of legs and gradually increase in size, eventually maturing into adults.

Infestations are visible as reddish patches or colonies on the underside of palm leaves, particularly near veins. Observations under a magnifying glass reveal dense aggregations of mites feeding simultaneously, causing discoloration and leaf deformation.

3. The Life Cycle: From Egg to Adult

Raoiella indica undergoes a complete life cycle consisting of five stages: egg, larva, protonymph, deutonymph and adult (Chavan *et al.*, 2020). Under laboratory conditions of $27 \pm 2^\circ\text{C}$ temperature and $79 \pm 5\%$ relative humidity, the egg incubation period lasts about 9-11 days, followed by larval development lasting 6-9 days.

The protonymph stage lasts 4-6 days, while the deutonymph stage takes 6-7 days. Adults live up to 30 days, with mated females laying an average of 8-10 eggs per day during their reproductive phase.

The mite thrives under warm and humid conditions, which are prevalent in tropical climates such as those in India. A single female can initiate a colony, with rapid development and short generation times allowing population explosions within weeks.

4. Feeding Habits and Host Range

Raoiella indica feeds by inserting its specialized mouthparts through or around the stomata of palm leaves, extracting sap and affecting vital plant processes like photosynthesis and respiration (Kane *et al.*, 2005; Ochoa *et al.*, 2011). Infested leaves turn yellow, exhibit necrotic patches and eventually wither, reducing crop yield and market value.

The pest’s host range includes:

- Coconut (*Cocos nucifera*)
- Arecanut (*Areca catechu*)
- Banana (*Musa* spp.)
- Date palm (*Phoenix dactylifera*)
- Heliconia spp.
- Ornamental palms

In India, Arecanut and coconut plantations are severely affected, especially during hot seasons when mite populations surge. Damage leads to reduced fruit production, poor growth and increased vulnerability to secondary infections.

5. Natural Enemies and Ecological Balance

Predatory mites like *Amblyseius largoensis* play an important role in keeping red palm mite populations in check (Carrillo *et al.*, 2010). These predators feed on all life stages of *Raoiella indica*, reducing their reproductive capacity and spread.

However, indiscriminate pesticide use disrupts this natural balance by killing beneficial predators, inadvertently worsening infestations. Integrated pest management emphasizes conserving natural enemies while using targeted control measures.

Global Spread and Invasive Potential

Although native to Southern Asia, *Raoiella indica* spread rapidly after being reported in the Caribbean in 2004 (Flechtmann and Etienne, 2004). Human-assisted movement of infested plant material, combined with wind dispersal, has allowed its expansion across tropical and subtropical regions.

It has been reported in India, the Middle East, Mauritius, Reunion, the Caribbean islands, South America and parts of Africa (CABI/EPPO, 2007; Melo *et al.*, 2018). Its capacity to establish in new regions is fuelled by its reproductive strategy, adaptability and the absence of specialized predators in invaded areas.

Why Should We Care? (Impact and Challenges)

Infestations of *Raoiella indica* in India threaten both smallholder and commercial farmers. Coconut and arecanut, major crops with cultural and economic importance, suffer substantial yield reductions.

Challenges include:

- Difficult detection due to small size
- Rapid population growth
- Inadequate awareness and training among farmers
- Overuse of pesticides harming ecological balance

Unchecked spread could disrupt rural livelihoods, strain agricultural economies and lead to long-term ecological consequences.

The Way Forward: Managing the Red Palm Mite

Effective management requires a combination of approaches:

Cultural Practices

- Use of clean planting material
- Regular monitoring and pruning of infested leaves

Biological Control

- Encouraging predatory mites like *Amblyseius largoensis*
- Avoiding broad-spectrum pesticides

Research and Awareness

- Extension services educating farmers
- Developing environmentally safe control strategies

Sustainable management emphasizes coexistence rather than eradication, balancing pest control with ecological preservation.

Conclusion: Living with a Tiny Invader

Raoiella indica may be tiny, but its impact on agriculture is profound. Understanding its morphology, life cycle, feeding habits and spread is essential for developing informed and sustainable management strategies. Farmers, researchers and policymakers must work together to promote eco-friendly approaches that protect livelihoods while conserving biodiversity. With vigilance, awareness and sciencebased interventions, it is possible to live with this invasive pest while minimizing its harm.

References

Cabi/Eppo. (2007). *Raoiella Indica* - Red Palm Mite. Crop Protection Compendium.

Chavan, I., Pradeep, S., Manjunatha, M., Narayanaswamy, H. And Sridhara, S. (2020). Biology of *Raoiella Indica* Hirst (Acari: Tenupalpidae) On Arecanut Leaves Under Laboratory Conditions. Agric. Sci. Digest., 40(3): 295-298.

Flechtmann, C. H. W. And Etienne, J. (2004). First Report of *Raoiella Indica* In the Caribbean. Exp. Appl. Acarol., 32(1): 123-124.

Hirst, S. (1924). Description of *Raoiella Indica* From Coconut in Coimbatore, India. Bull. Entomol. Res., 15: 87-90.

Kane, E. C., Ochoa, R. And Hernández, A. (2005). Feeding Mechanism of *Raoiella Indica* On Palm Leaves. Int. J. Acarol., 31(3): 179-185.

Melo, J. W. S., Navia, D., Mendes, J. A., Filgueiras, R. M. C., Teodoro, A. V., Ferreira, J. M. S., Guzzo, E. C., De Souza, I. V., De Mendonça, R. S., Calvet, É. C., Paz Neto, A. A., Gondim Jr., M. G. C., De Moraes, E. G. F., Godoy, M. S., Dos Santos, J. R., Silva, R. I. R., Da Silva, V. B., Norte, R. F., Oliva, A. B., ... Domingos, C. A. (2018). The Invasive Red Palm Mite, *Raoiella Indica*, In Brazil: Range Extension and Arrival into The Most Threatened Area. Int. J. Acarol., 44(4-5): 146-149. <https://doi.org/10.1080/01647954.2018.1474945>

Ochoa, R., Beard, J. J., Baughan, G. R., Welbourn, W. C. And Pooley, C. (2011). Stomatal Feeding Behavior Of *Raoiella Indica*. Am. Entomol., 57(1): 26-29.

Patel, K. D. And Rao, K. R. (1958). First Report of *Raoiella Indica* On Arecanut In India. Indian J. Agric. Sci., 28(6): 289-294.

