

Molecular Chaperone Functions in Protein Folding and Proteostasis

Yujin E. Kim,* Mark S. Hipp,* Andreas Bracher,
Manajit Hayer-Hartl, and F. Ulrich Hartl

Department of Cellular Biochemistry, Max Planck Institute of Biochemistry,
82152 Martinsried, Germany; email: uhartl@biochem.mpg.de

Annu. Rev. Biochem. 2013. 82:323–55

The *Annual Review of Biochemistry* is online at
biochem.annualreviews.org

This article's doi:
10.1146/annurev-biochem-060208-092442

Copyright © 2013 by Annual Reviews.
All rights reserved

*These authors contributed equally.

Keywords

chaperonin, degradation, Hsc70/Hsp70, Hsp90, protein folding,
trigger factor

Abstract

The biological functions of proteins are governed by their three-dimensional fold. Protein folding, maintenance of proteome integrity, and protein homeostasis (proteostasis) critically depend on a complex network of molecular chaperones. Disruption of proteostasis is implicated in aging and the pathogenesis of numerous degenerative diseases. In the cytosol, different classes of molecular chaperones cooperate in evolutionarily conserved folding pathways. Nascent polypeptides interact cotranslationally with a first set of chaperones, including trigger factor and the Hsp70 system, which prevent premature (mis)folding. Folding occurs upon controlled release of newly synthesized proteins from these factors or after transfer to downstream chaperones such as the chaperonins. Chaperonins are large, cylindrical complexes that provide a central compartment for a single protein chain to fold unimpaired by aggregation. This review focuses on recent advances in understanding the mechanisms of chaperone action in promoting and regulating protein folding and on the pathological consequences of protein misfolding and aggregation.

Contents

INTRODUCTION	324
Protein Folding and Aggregation	324
Protein Folding In Vivo	326
THE MOLECULAR CHAPERONE	327
CONCEPT	327
Chaperone-Mediated Folding	
by Kinetic Partitioning	327
Role in Protein Evolution	328
CHAPERONE PATHWAYS IN	
THE CYTOSOL	328
Chaperone Action on the	
Ribosome	328
Chaperones Acting Downstream	
of the Ribosome	330
CHAPERONE PARADIGMS	331
Trigger Factor	331
The Hsp70 System	332
The Chaperonin System	334
The Hsp90 System	338
PROTEOSTASIS NETWORK	340
Significance of Conformational	
Maintenance	341
Degradation	341
Proteostasis Collapse in Aging	
and Disease	343
The Proteostasis Network as a	
Drug Target	344
OUTLOOK	344

INTRODUCTION

The successful execution of cellular processes depends on the coordinated interactions of proteins. In humans, an estimated 20,000 to 25,000 different proteins are responsible for most aspects of biological function. Following synthesis on ribosomes as linear sequences of amino acids, the vast majority of proteins must fold into well-defined three-dimensional structures (their native states) to attain functionality. Although some newly translated proteins are able to fold spontaneously, a substantial fraction of proteins are less efficient at folding and vulnerable to misfolding, a problem exacerbated by the highly crowded cellular environment (1). In

particular, large proteins with complex structures may expose hydrophobic amino acid residues to the solvent during folding, rendering them susceptible to nonnative (off-pathway) interactions that lead to aggregation (2, 3). To counteract these nonnative interactions, cells have a network of molecular chaperones that assist in de novo folding and maintain preexisting proteins in their native states (4–7). A key role of molecular chaperones is preventing protein aggregation, especially under conditions of cellular stress. Moreover, the molecular chaperone network functions in diverse aspects of protein quality control, including protein unfolding and disaggregation and targeting terminally misfolded proteins for proteolytic degradation.

Imbalances in protein homeostasis (proteostasis) are observed in an increasing number of disease states, emphasizing the importance of cellular protein quality control (8). The predominant feature of these disorders is protein misfolding as manifested by the formation of intracellular and/or extracellular deposits of aggregated proteins. Examples include the formation of intracellular inclusions containing aggregated α -synuclein in Parkinson's disease or huntingtin in Huntington's disease, as well as the extracellular β -amyloid plaques in Alzheimer's disease (9). Deficiencies in proteostasis are also observed in many other age-related diseases, such as type II diabetes, peripheral amyloidosis, cancer, and cardiovascular diseases. Indeed, studies using model organisms demonstrate that a gradual decline in cellular proteostasis capacity occurs with aging (10).

Here, we review recent advances in understanding the role of molecular chaperones in protein folding and proteostasis maintenance. We focus our discussion on the cytosolic chaperone networks and the pathological consequences of their disruption. For a discussion of protein folding in the secretory pathway and the folding of membrane proteins, we refer the reader to recent reviews (11, 12, 13).

Protein Folding and Aggregation

The folded three-dimensional structures of most proteins represent a compromise between

thermodynamic stability and the conformational flexibility required for function. Consequently, proteins are often marginally stable in their physiological environment and thus susceptible to misfolding and aggregation (2, 3). In addition, a substantial fraction of proteins in eukaryotic cells (~30%) are classified as intrinsically unstructured and contain regions thought to adopt ordered structure only upon interaction with binding partners (14). Such proteins may be metastable and prone to aggregation.

The pioneering studies of Anfinsen revealed that small denatured proteins refold spontaneously *in vitro*, demonstrating that the three-dimensional structure of a protein is encoded in its amino acid sequence (2, 3). Much progress in recent years has helped us understand how exactly the linear sequence of amino acids encodes the native state of a protein and directs its folding process. Because the number of possible conformations a protein chain can adopt is very large, folding reactions are highly complex and heterogeneous, relying on the cooperation of multiple weak, noncovalent interactions. Among these, hydrophobic forces are critical in driving chain collapse and the burial of nonpolar amino acid residues within the interior of the folding protein. Polypeptide chains are thought to explore funnel-shaped potential energy surfaces as they progress toward the native structure along several downhill paths rather than a single defined pathway (Figure 1) (2). Rapid chain collapse and the incremental formation of native contacts limit the conformational space that must be searched en route to the native state. However, the rugged free-energy surface navigated during folding often requires molecules to cross substantial kinetic barriers. As a result, kinetically trapped folding intermediates and misfolded states may be transiently populated. Misfolded states are characterized by the presence of nonnative interactions that must be resolved prior to correct folding. Productive folding intermediates may display a high degree of configurational flexibility, increasing the search time required for the formation of native intra-chain contacts. The propensity of proteins to




Figure 1

Competing reactions of protein folding and aggregation. Proteins fold by sampling various conformations in a folding energy landscape. Energetically favorable intramolecular interactions (green) are stabilized as the protein progresses on a downhill path through the landscape toward the native state. Energetically favorable but nonnative conformations result in populations of kinetically trapped states that occupy low-energy wells (partially folded states or misfolded states). Chaperones assist these states in overcoming free energy barriers and prevent intermolecular interactions (red) leading to aggregation (amorphous aggregates, β -sheet-rich oligomers, and amyloid fibrils), thus promoting folding to the native state. Figure adapted and modified from References 17 and 237.

populate such entropically stabilized intermediates increases with larger, more topologically complex domain folds that are stabilized by many long-range interactions (such as α/β domain architectures). Such proteins are often highly chaperone dependent (15).

Partially folded or misfolded states typically expose hydrophobic amino acid residues and regions of unstructured polypeptide backbone to the solvent, the features that mediate aggregation in a concentration-dependent manner (Figure 1) (9, 16). Although aggregation primarily leads to amorphous structures largely driven by hydrophobic forces, it may also lead to the formation of amyloid-like fibrils (Figure 1). These fibrillar aggregates are characterized by β -strands that run perpendicular to the long

Protein folding: the process by which the extended, newly synthesized polypeptide chain collapses into its functional three-dimensional conformation

fibril axis (cross- β structure) and are associated with diseases of aberrant protein folding. Many proteins can adopt these highly ordered, thermodynamically stable structures in vitro, but molecular chaperones restrict their formation in vivo (9, 16, 17). The formation of soluble oligomeric intermediates often accompanies the formation of fibrillar aggregates (**Figure 1**). The pronounced toxicity of these less-ordered and rather heterogeneous forms likely correlates with the exposure of interaction-prone hydrophobic surfaces, which in the fibril are integrated into a compact cross- β core (18). We can view certain oligomers as kinetically trapped intermediates that must undergo considerable structural rearrangement to form fibrils, the thermodynamic end state of aggregation (**Figure 1**). The relative impact of fibrillar and oligomeric aggregates on proteostasis and cell health is currently under intense investigation.

Protein Folding In Vivo

Attempts to understand protein folding in vivo must take into consideration the dramatic differences in physical properties that exist between the cellular environment and the conditions of test-tube refolding (19, 20). Compared with the dilute conditions in vitro, the cellular environment is highly crowded, containing concentrations of 300–400 mg ml⁻¹ of protein and other macromolecules (1, 21). Macromolecular crowding results in excluded volume effects, limiting the entropic freedom of the folding polypeptide chains and favoring compact nonnative states (1, 22). In addition, macromolecular crowding enhances protein aggregation (amorphous and fibrillar) by increasing the affinities between interacting macromolecules including folding intermediates (1).

Protein folding in vivo is further complicated by its coupling with translation and by the fact that many newly synthesized polypeptides must be transported into subcellular compartments, such as the endoplasmic reticulum (ER) or the mitochondria (23), prior to folding. The vectorial translation process from the N

terminus to C terminus places considerable restrictions on the energy landscape of in vivo folding (24, 25). The exit tunnel of the large ribosomal subunit, ~100 Å long and ~20 Å wide, precludes folding beyond the formation of α -helices or small tertiary structural elements that may begin to form near the tunnel exit (26–29). Thus, the C-terminal 30–40 amino acid residues of the nascent chain cannot participate in the long-range interactions necessary for cooperative domain folding. Consequently, productive folding is delayed until a complete protein domain (~50–300 amino acid residues), or substantial segments thereof, has emerged from the ribosome (30–35). Whereas single-domain proteins complete folding posttranslationally (after chain termination and release from the ribosome), proteins consisting of multiple domains may fold cotranslationally as the domains emerge sequentially from the ribosome. Sequential folding of the domains prevents the formation of unproductive intermediates resulting from nonnative interactions between concomitantly folding domains (36, 37). For the multidomain protein firefly luciferase, sequential domain folding results in a dramatic acceleration of folding speed (6, 7, 36). The slower translation speed in eukaryotes (~4 amino acids s⁻¹) compared with bacteria (~20 amino acids s⁻¹) (38), together with evolutionary adaptations of the chaperone machinery, may facilitate cotranslational folding for domains with slower folding kinetics and thus may have contributed to the explosive evolution of multidomain proteins in eukaryotes. Although domain size was conserved during evolution, the average size of proteins increased from ~35 kDa in bacteria such as *Escherichia coli* to ~52 kDa in humans (36). Translational pausing may also enhance the efficiency of cotranslation folding, but the significance of this phenomenon in vivo is still under investigation (34, 39–42).

The fastest translation speeds are slow compared with the rapid kinetics of folding observed for small protein domains in vitro, some of which fold on the microsecond to millisecond timescale (2, 3). A nascent chain

of average length (\sim 300 amino acid residues in *E. coli*) will be exposed on the ribosome in an unfolded state for \sim 15 s, during which it may engage in nonnative intra- and inter-chain interactions. Contrary to the previous belief that polysomes enhance aggregation by increasing the local concentration of nascent chains, recent studies suggest that the three-dimensional organization of individual ribosomes in polysomes maximizes the distance between nascent chains, thus reducing the probability of unproductive interactions (43).

THE MOLECULAR CHAPERONE CONCEPT

We define a molecular chaperone as any protein that interacts with and aids in the folding or assembly of another protein without being part of its final structure (4). Chaperones are classified into different groups on the basis of sequence homology. Many are stress proteins or heat shock proteins (Hsps), as their synthesis is induced under conditions of stress (e.g., heat shock or oxidative stress), which structurally destabilize a subset of cellular proteins. Members of the various groups of chaperones were initially named according to their molecular weight: Hsp40s, Hsp60s, Hsp70s, Hsp90s, Hsp100s, and the small Hsps. Besides their fundamental role in de novo protein folding, chaperones are involved in various aspects of proteome maintenance, including assistance in macromolecular complex assembly, protein transport and degradation, and aggregate dissociation and refolding of stress-denatured proteins.

Chaperone-Mediated Folding by Kinetic Partitioning

Chaperones that function broadly in de novo folding and refolding (i.e., the chaperonins, Hsp70s, and Hsp90s) are ATP regulated and recognize segments of exposed hydrophobic amino acid residues, which are later buried in the interior of the natively folded protein. Binding to hydrophobic segments

enables these chaperones to recognize the nonnative states of many different proteins. Folding is then promoted during ATP- and cochaperone-regulated cycles of binding and release of nonnative protein (Figure 2). In this mechanism of kinetic partitioning, (re)binding to chaperones blocks aggregation and reduces

Assembly: the association of two or more protein molecules in a functional complex




Figure 2

Molecular chaperones promote protein folding through kinetic partitioning. Under physiological conditions, the unfolded state of a protein undergoes rapid collapse to a partially folded, compact intermediate. Although chain collapse restricts the conformational space that must be searched en route to the native state, the collapsed folding intermediates are often aggregation-prone, kinetically trapped states. Many molecular chaperones switch between high- and low-affinity states for unfolded and partially folded proteins in a manner regulated by ATP binding and hydrolysis. Folding proceeds efficiently when the folding rate constant (K_{fold}) is greater than the rate constants for chaperone binding (K_{on}) and aggregation (K_{agg}). Binding or rebinding of nonnative protein to chaperone allows kinetic partitioning, preventing aggregation and favoring folding. When K_{on} is greater than K_{fold} , the chaperone system functions as a holdase, stabilizing the protein in a nonaggregated state for transfer to other chaperone systems or for degradation. During exposure to conformational stress, K_{agg} becomes greater than K_{on} , resulting in aggregation unless the cell upregulates its chaperone capacity through stress response pathways. K_{off} is the rate constant for protein release from chaperone and P_i is inorganic phosphate. Figure adapted and modified from Reference 38.

Stress proteins:

cellular proteins whose expression is induced or increased under conditions of conformational stress; not all molecular chaperones are stress proteins

Aggregate: the association of two or more protein molecules in a nonfunctional state

Chaperonin: a class of structurally related molecular chaperones forming large, double-ring complexes that transiently enclose substrate protein for folding

TF: trigger factor

NAC: nascent-chain-associated complex

Nascent polypeptide: the polypeptide chain emerging from the ribosome during translation

RAC: ribosome-associated complex

the concentration of free folding intermediates, whereas transient release of bound hydrophobic regions is necessary for folding to proceed (**Figure 2**). ATP-independent chaperones, such as the small Hsps, may function as additional holdases that buffer aggregation. Efficient folding is achieved when the rate of folding is faster than the rates of aggregation or chaperone rebinding. If folding is slower, then the protein may be transferred to a chaperone system with different mechanistic properties, as exemplified by the sequential cooperation between Hsp70 chaperones and the cylindrical chaperonins (discussed below). Alternatively, transfer to the degradation machinery may occur. Aggregation occurs if the concentration of folding intermediates exceeds the available chaperone capacity. Such a situation generally results in the induction of the cellular stress response, which increases the abundance of stress-regulated chaperones.

Role in Protein Evolution

The general function of chaperones in assisting protein folding is significant in facilitating the structural evolution of proteins. By maintaining nonnative proteins in a soluble, folding-competent state, chaperones are thought to buffer mutations in proteins that would otherwise preclude their folding, thus broadening the range of mutant proteins subject to Darwinian selection (44, 45). After selection of a mutant protein with favorable functional properties, secondary mutations may improve its folding efficiency and solubility, allowing the protein to become less chaperone dependent and increase in abundance. Interestingly, proteins that depend highly on a specific chaperone system, such as the *E. coli* chaperonin GroEL, are of less than average abundance and often have nonessential functions (15). Conversely, highly abundant proteins with essential functions tend to be less dependent on a specific chaperone and may use multiple chaperone systems to optimize folding yield (46). Notable exceptions include the photosynthetic enzyme ribulose-1,5-bisphosphate carboxylase

oxygenase (RuBisCO) as well as the cytoskeletal proteins actin and tubulin. Although highly abundant, these proteins are obligate substrates of the chaperonin system for folding (4, 6). Presumably, RuBisCO, actin, and tubulin reside in an evolutionarily trapped state in which further mutations that reduce chaperonin dependence are incompatible with function.

CHAPERONE PATHWAYS IN THE CYTOSOL

The general organization of cytosolic chaperone pathways is highly conserved throughout evolution (**Figure 3**). In all domains of life (bacteria, archaea, and eukarya), ribosome-binding chaperones [e.g., trigger factor (TF), nascent-chain-associated complex (NAC), and specialized Hsp70s] interact first with the nascent polypeptide, followed by a second tier of chaperones without direct affinity for the ribosome (the classical Hsp70 system). Folding may begin cotranslationally and finish posttranslationally upon chain release from the ribosome or after transfer to downstream chaperones (e.g., the chaperonins and Hsp90 system) (**Figure 3**). Recent system-wide and bioinformatic approaches identified the substrate interactome of several major chaperones, revealing the cooperative organization of the chaperone network (15, 47–57).

Chaperone Action on the Ribosome

As discussed above, the nascent polypeptide chain is topologically restricted until a complete protein domain is synthesized and emerges from the ribosomal tunnel. Ribosome-binding chaperones prevent emerging nascent chains from engaging in unfavorable intra- and intermolecular interactions during translation, typically by shielding exposed hydrophobic segments. The ribosome-associated molecular chaperones include TF (in prokaryotes) and specialized Hsp70 complexes such as ribosome-associated complex (RAC; in *Saccharomyces cerevisiae*), MPP11 and Hsp70L1




Figure 3

Organization of chaperone pathways in the cytosol. In bacteria (*a*), archaea (*b*), and eukarya (*c*), ribosome-bound chaperones [trigger factor (TF) in bacteria, nascent-chain-associated complex (NAC) in archaea and eukarya] aid folding cotranslationally by binding to hydrophobic segments on the emerging nascent chains. For longer nascent chains, members of the heat shock protein (Hsp)70 family (DnaK in bacteria and Hsp70 in eukarya), together with Hsp40s and nucleotide exchange factors (NEFs), mediate co- and posttranslational folding. In archaea lacking the Hsp70 system, prefoldin (PFD) assists in folding downstream of NAC. Partially folded substrates may be transferred to the chaperonins [GroEL-GroES in bacteria, thermosome in archaea, and tailless complex polypeptide-1 (TCP-1) ring complex (TRiC)/chaperonin-containing TCP-1 (CCT) in eukarya]. The Hsp90 system also receives its substrates from heat shock cognate 70 (Hsc70) and mediates their folding with additional cofactors. The insert in panel *c* shows the ribosome-binding chaperone system, the ribosome-associated complex (RAC), in fungi. RAC consists of Ssz1 (a specialized Hsp70) and zuotin (Hsp40) and assists nascent chain folding together with another Hsp70 isoform, Ssb. Percentages indicate the approximate protein flux through the various chaperones. Figure adapted and modified from Reference 7.

(in mammals), and NAC (in archaea and eukaryotes) (Figure 3) (17, 58, 59).

TF binds to the large ribosomal subunit at the opening of the ribosomal exit tunnel (60–62) and interacts with most newly synthesized cytosolic proteins as well as a subset of secretory proteins (55, 56, 58, 59). *In vitro*, TF binds to nascent chains as short as ~60 amino acid residues, presumably when the first hydrophobic segment of the chain has emerged (58, 63). *In vivo*, TF preferentially binds ribosomes when nascent chains have reached ~100 amino acids in length (55), thereby permitting prior interactions of ribosome-binding targeting factors (e.g., signal recognition particle)

(64, 65) and modifying enzymes (e.g., peptide deformylase) (66) with the nascent chains. Release of TF from the nascent chain is ATP independent and permits folding or polypeptide transfer to downstream chaperones such as DnaK, the major Hsp70 in bacteria. Although the combined deletion of the genes encoding TF and DnaK is lethal at temperatures above 30°C, *E. coli* cells tolerate individual deletions of TF or DnaK, indicating that these proteins are functionally redundant (56, 58, 67).

In eukaryotes, RAC and NAC may fulfill a role similar to TF, although they are not structurally related to TF. RAC comprises the Hsp70-like protein Ssz1 (also known as Pdr13)

NEF: nucleotide exchange factor

TRiC: tailless complex polypeptide-1 ring complex

and the Hsp70 cochaperone zuotin (Hsp40) (**Figure 3c**) (58, 59, 68–71). In fungi, RAC cooperates with ribosome-binding isoforms of Hsp70, Ssb1, or Ssb2. NAC, a heterodimeric complex of α - (33 kDa) and β - (22 kDa) subunits, also associates with ribosomes (via the β -subunit) and short nascent chains (54, 59, 72, 73). The exact role of NAC in folding or protein quality control is not established. In yeast, NAC function appears to be partially redundant with that of Ssb (71), reminiscent of the interplay between TF and DnaK in bacteria. Both Ssb-RAC and NAC may have a role in ribosome biogenesis (71). Archaea have a homolog of α -NAC.

Chaperones Acting Downstream of the Ribosome

In bacteria and eukaryotic cells, the classical Hsp70s have a central role in the cytosolic chaperone network (6, 17). They interact with a multitude of nascent and newly synthesized polypeptides but have no direct affinity for the ribosome (56, 74). The nascent-chain-interacting Hsp70 chaperones include DnaK in bacteria and some archaea, Ssa1–4 in yeast, and the constitutively expressed heat shock cognate 70 (Hsc70) in metazoan and mammalian cells (4, 58). Hsp70 chaperones function with cochaperones of the Hsp40 family (also known as DnaJ proteins or J proteins) and nucleotide exchange factors (NEFs) (**Figure 3**) (75). In addition to protecting nascent chains against aberrant interactions, the Hsp70-Hsp40 chaperone systems assist folding co- or posttranslationally through ATP-regulated cycles of substrate binding and release (by kinetic partitioning; **Figure 2**) and mediate polypeptide chain transfer to downstream chaperones. Remarkably, most species of archaea lack the Hsp70 chaperone system. The chaperone prefoldin (also known as the Gim complex, GimC) may substitute for Hsp70 in these cases (7). Prefoldin, a hexamer of ~14–23 kDa α - and β -subunits with long α -helical coiled coils, binds in an ATP-independent manner to certain nascent chains and mediates their transfer to the cylindrical chaperonin complex for the

final stages of folding. In eukaryotes, prefoldin participates in the chaperonin-assisted folding of actin and tubulin (6, 7).

Proteins that are unable to utilize Hsp70 for folding are transferred to the chaperonin or the Hsp90 system. Chaperonins (also referred to as Hsp60s) are large double-ring complexes of 800–1,000 kDa with a central cavity, which permits protein molecules to fold in an isolated compartment protected from the aggregation-promoting cytosol (**Figure 3**) (4, 5). The chaperonins are structurally classified into group I and group II (76, 77). Group I chaperonins include GroEL in bacteria, Hsp60 in mitochondria, and Cpn60 in chloroplasts. They cooperate with lid-shaped cochaperones (GroES, Hsp10, Cpn10/20) to encapsulate substrates. The group II chaperonins include the archaeal thermosome and its eukaryotic homolog tailless complex polypeptide-1 (TCP-1) ring complex (TRiC), also known as chaperonin-containing TCP-1 (CCT), which have a built-in lid. GroEL-GroES assist folding posttranslationally, whereas TRiC may assist folding co- and posttranslationally (6). TRiC binds to nascent chains and cooperates with Hsp70 in the cotranslational folding of multidomain proteins (78). Investigators have demonstrated a direct interaction between Hsp70 and TRiC (79). The chaperonins interact with 10–15% of newly synthesized polypeptides in bacteria and archaea (15, 52, 53) and 5–10% in eukarya (50). The obligate substrates of GroEL typically include proteins with complex domain folds, which tend to populate kinetically trapped folding intermediates (15, 53). Established TRiC substrates include the cytoskeletal proteins actin and tubulin as well as several proteins with β -propellers/WD40 repeats (49, 50).

In the eukaryotes, many signaling proteins are transferred from Hsp70 to the ATP-dependent Hsp90 chaperone system for completion of folding and conformational regulation (**Figure 3c**) (45, 48, 80). Examples of Hsp90-mediated conformational regulation include nuclear hormone receptors, which Hsp90 stabilizes in a conformation poised for hormone binding and activation. Substrate

transfer to Hsp90 is mediated by the Hsp90 organizing protein (HOP; also known as St1 and p60), which uses multiple tetratricopeptide repeat (TPR) domains to bridge Hsp70 and Hsp90 (81). Various cochaperones facilitate the interaction of Hsp90 with steroid hormone receptors and multiple protein kinases (45, 48). Accordingly, Hsp90 affects many key cellular processes, including cell cycle progression, steroid signaling, calcium signaling, protein trafficking, protein secretion, the immune response, and the heat shock response (HSR) (45, 48, 82). Pharmacologic inhibition of Hsp90 with geldanamycin and derivatives results in the downregulation of many kinases (83) and is in clinical development for cancer therapy (84).

The molecular chaperone network is central to cellular protein quality control through its involvement in the conformational maintenance of proteins, the dissociation of aggregates, and the degradation of misfolded proteins. In yeast and other fungi, the cytosolic Hsp70 system cooperates with the AAA+ (ATPases associated with various cellular activities) chaperone Hsp104 in dissociating and refolding aggregated proteins. Hsp104 is homologous to bacterial ClpB, which functions with DnaK in protein disaggregation (85). Various cochaperones of Hsp70 and Hsp90 escort terminally misfolded proteins to the protein degradation machinery (ubiquitin-proteasome system or autophagy) (86).

CHAPERONE PARADIGMS

Research has defined several mechanistic paradigms of chaperone function in protein folding. In the following sections we discuss TF as an ATP-independent chaperone as well as the Hsp70 system, the chaperonins, and Hsp90 as ATP-dependent paradigms. Structural and functional data strongly support the mechanistic models of each of these systems.

Trigger Factor

Bacterial TF is an abundant ~50 kDa protein that binds to ribosomes and interacts with most nascent polypeptides (7, 55, 56, 59, 63, 87,

88). The crystal structure of *E. coli* TF reveals an elongated structure consisting of three domains, an N-terminal ribosome-binding domain, a peptidylprolyl *cis/trans* isomerase (PPIase) domain, and a C-terminal domain (positioned between the N-terminal and PPIase domains) (Figure 4a) (61). The N-terminal

PPIase:
peptidylprolyl *cis/trans* isomerase




Figure 4

Structure and functional cycle of trigger factor (TF). (a) Structure of TF. The N-terminal domain of TF (magenta) contains the Phe-Arg-Lys (FRK) ribosome-binding loop and connects by a long linker to the peptidylprolyl *cis/trans* isomerase (PPIase) domain (blue). The C-terminal domain (green) is in the center of the protein and provides the main binding site for the nascent chain substrate [Protein Data Bank (PDB) 1W26]. (b) TF reaction cycle. Free TF exists in the cytosol as a dimer in rapid equilibrium with monomers. Monomeric TF binds to ribosomes with translating nascent chains. Hydrophobic sequence motifs in the nascent chain regulate TF-nascent chain interaction. Binding to the elongating nascent chain may persist as TF dissociates from the ribosome (with $t_{1/2} \sim 10$ s) (63), allowing a second TF molecule to bind to the ribosome at the polypeptide exit site. Concurrent with release from TF, the newly synthesized polypeptide folds into its native state or is transferred to downstream chaperone systems.

NBD:

nucleotide-binding domain

SBD:

substrate-binding domain

domain mediates binding of monomeric TF to the large ribosomal subunit at proteins L23 and L29 in close proximity to the polypeptide exit site (60, 61, 89, 90). L23 is essential for TF binding and is thought to signal the progression of the nascent chain through the ribosomal exit tunnel (89), whereas L29 has an auxiliary function (60). The PPIase domain connects to the N domain via a long linker and is most distal to the ribosome docking site. It belongs to the family of FK506 binding proteins and recognizes stretches of eight amino acids that are enriched in basic and aromatic residues (7). Studies have measured PPIase activity *in vitro*, but its contribution to folding *in vivo* has remained unclear, as the domain is dispensable for TF function (59, 67, 91, 92). Cross-linking experiments showed that the PPIase domain interacts with longer nascent chains, presumably representing an auxiliary substrate recognition site (62, 87). The centrally located C-terminal domain is structurally similar to the N-terminal domain of the periplasmic chaperone SurA and provides the primary binding site for the nascent chain (87, 93).

Cells contain an excess of TF compared with ribosomes; the non-ribosome-bound fraction is in rapid monomer-dimer equilibrium (**Figure 4b**) (63, 94). TF monomer aids de novo folding through ATP-independent cycles of binding and release from both the ribosome and the nascent chain. Binding to the ribosome (mean residence time of ~10–15 s) is a prerequisite for interaction with the nascent chain. The disposition of the bound peptide to bury hydrophobic regions during folding likely drives the eventual release of TF from the nascent chain. Accordingly, TF slows hydrophobic chain collapse and delays cotranslational folding (37, 95, 96). Furthermore TF may remain bound to certain polypeptides after their release from the ribosome, which is consistent with a role for TF as a holdase in ribosome assembly (97, 98). Several studies indicate that TF retains a high degree of conformational flexibility on the ribosome during interactions with the emerging nascent chain (62, 63, 89, 90). This flexibility likely

enables TF to accommodate a wide range of polypeptides (63, 88). Although it primarily binds hydrophobic chain segments (63, 88), TF also interacts with small basic proteins, including many ribosomal proteins (56, 88, 98).

The Hsp70 System

Hsp70 chaperones are a ubiquitous class of proteins. They are involved in a wide range of protein quality control functions, including *de novo* protein folding, refolding of stress-denatured proteins, protein transport, membrane translocation, and protein degradation.

Structure and reaction cycle. Hsp70 consists of an N-terminal nucleotide-binding domain (NBD) and a C-terminal substrate-binding domain (SBD) connected by a highly conserved hydrophobic linker region (**Figure 5a**). The N-terminal domain has an actin-like fold; it consists of two lobes, each containing two subdomains, with the nucleotide-binding cleft situated in between (5, 99). The SBD consists of a β -sandwich subdomain and an α -helical lid with the substrate binding site located in the β -sandwich subdomain (5). The SBD binds to 5–7-residue peptide segments enriched in hydrophobic amino acids and typically flanked by positively charged residues. The peptide binds in an extended conformation mediated by hydrogen bonds between the SBD and the peptide backbone and by van der Waals contacts with the hydrophobic side chains (5, 100).

Conformational changes in the NBD upon ATP binding and hydrolysis are allosterically coupled to the SBD, regulating peptide binding and release. Binding of ATP to the NBD triggers the attachment of the hydrophobic interdomain linker and the α -helical lid of the SBD to the NBD, which opens the peptide binding pocket (101, 111), as initially revealed in structures of the Hsp70 homolog Hsp110 (**Figure 5a**) (103–105). Hydrolysis of ATP to ADP triggers the detachment of the lid from the NBD and the closing of the SBD; NBD and SBD are loosely held together by the linker in a dynamic random coil conformation (106, 107).




Figure 5

Structure and functional cycle of Hsp70. (a) Structure of Hsp70. Hsp70 consists of two domains, the nucleotide-binding domain (NBD) and the substrate-binding domain (SBD), connected by a conserved linker. The closed state of *Escherichia coli* DnaK (left; PDB 2KHO) was solved using a combination of solution nuclear magnetic resonance and crystal structures of the individual domains (106). The structure illustrates the ADP-bound NBD separated by a linker from the SBD. The α -helical lid of the SBD is closed over the substrate peptide (NRLLLTG) bound in the pocket of the β -sandwich domain. The open state is illustrated by the crystal structure of ATP-bound Sse1 (yeast Hsp110; right; PDB 2QXL). The β -sandwich domain contacts subdomain IA of the NBD, whereas the α -helical lid contacts subdomains IA and IB. (b) Reaction cycle. ATP binding to the NBD stabilizes the open state of Hsp70, facilitating the binding of substrate protein recruited to Hsp70 by Hsp40 cochaperone. The open state has fast on and off rates for substrate peptide. Hsp40 stimulates ATP hydrolysis on Hsp70, resulting in the closing of the SBD α -helical lid over the bound substrate peptide. The closed state has slow on and off rates for substrate peptide. NEFs stimulate the release of ADP from the NBD, and ATP binding causes substrate release.

Details of this allosteric mechanism are under intense investigation (108). The ATP-bound open state has high on and off rates for peptide substrate, whereas the ADP-bound closed state has low on and off rates (**Figure 5b**) (5, 100). In turn, substrate binding increases the rate of ATP hydrolysis (109–111). Substrate release allows folding (i.e., the burial of hydrophobic residues) to proceed. Proteins unable to fold rapidly upon dissociation from Hsp70 may rebind, transfer to downstream chaperones, or be transferred to the degradation machinery.

Cochaperones. Hsp40 (J protein) and NEF cochaperones regulate the Hsp70 reaction cycle (38, 100). The Hsp40 proteins constitute a large family with more than 40 members in humans (75). All of them contain a J domain, which binds to the N-terminal ATPase domain of Hsp70 and the adjacent linker region (112, 113). Canonical Hsp40s (members of classes I and II) function as chaperones independently and recruit Hsp70 to nonnative substrate proteins. Other Hsp40s (class III) are more diverse and combine the J domain with a variety of functional modules (75, 114, 115). The interaction with Hsp70 strongly stimulates (>1,000-fold) the hydrolysis of Hsp70-bound ATP to ADP, resulting in stable substrate binding by Hsp70 in the closed conformation (38, 100). Subsequent binding of an NEF to the NBD of Hsp70 catalyzes the exchange of ADP to ATP, opening the SBD and thereby triggering substrate release (**Figure 5b**).

Bacteria have one Hsp70 NEF, GrpE, whereas eukaryotic cells contain several structurally unrelated families of NEFs, including the Bcl-2-associated athanogene (BAG) domain proteins, HspBP1 and Hsp110 proteins (116–119). Crystal structures of Hsp70-NEF complexes suggest that all NEFs, except HspBP1, stabilize the Hsp70 NBD in a conformation with an open nucleotide-binding cleft (104, 105, 120–123). The most abundant and ubiquitous eukaryotic NEFs are the Hsp110 proteins, which are structurally related to Hsp70 (103–105). Hsp110 may function as holdases for nonnative proteins and cooperate with Hsp70

and Hsp40 in protein disaggregation (104, 124, 125).

Proper regulation of the ATPase cycle is crucial for efficient Hsp70 function. Hsp40 and NEF proteins are present at lower levels relative to their partner Hsp70. They provide a means of diversifying Hsp70 function and establish substrate specificities for the Hsp70 machinery. For example, the human genome encodes 41 different J-domain proteins compared with 11 Hsp70s and 13 NEFs (75). Interestingly, inactivating mutations in the NEF Sil1 cause the neurodegenerative disease Marinesco-Sjögren syndrome, which highlights the critical importance of the nucleotide exchange function (126, 127).

The Chaperonin System

Chaperonins are large double-ring complexes of 7–9 ~60 kDa subunits per ring. They are unique among molecular chaperones in that they encapsulate their substrate proteins, one molecule at a time, in a central cavity for folding unimpaired by aggregation (4, 5, 17). Two groups of distantly related chaperonins can be distinguished (6, 76, 77).

Group I chaperonin GroEL. Group I chaperonins are present in the bacterial cytosol (GroEL), the mitochondrial matrix (Hsp60), and the stroma of chloroplasts (Cpn60). They have seven subunits per ring and are defined by their functional requirement for lid-shaped cochaperones (GroES in bacteria, Hsp10 in mitochondria, and Cpn10/Cpn20 in chloroplasts). The GroEL-GroES system of *E. coli* has been studied most extensively. The subunits of GroEL contain an equatorial ATPase domain, an intermediate hinge domain, and an apical domain (**Figure 6a**) (77, 100). The apical domains form the entrance to the GroEL cavity and expose hydrophobic amino acid residues, which mediate substrate binding. GroEL-bound substrates typically interact with multiple apical domains (77, 128) and populate an ensemble of compact and locally expanded states that lack stable tertiary interactions, similar to a molten globule (4, 129–131). Binding to GroEL prevents aggregation of these flexible

folding intermediates, and subsequent folding depends critically on the global encapsulation of the substrate in the chaperonin cavity by the cochaperone GroES (7, 132–135). GroES is a heptameric ring of ~10 kDa subunits that binds to the apical GroEL domains, capping the GroEL cylinder (**Figure 6a,b**) (4, 77, 100).


GroEL and GroES undergo a complex binding-and-release cycle that is allosterically regulated by ATP binding and hydrolysis in the GroEL subunits (**Figure 6b**) (4, 77, 100, 136). Cooperative binding of ATP to GroEL initiates a series of conformational changes that trigger the association of GroES, followed by substrate release from hydrophobic binding sites into a GroES-capped, hydrophilic folding chamber (137). Proteins up to ~60 kDa can be encapsulated and are free to fold in the cage for ~10 s (at 25°C) (132), the time needed for ATP hydrolysis in the GroES-capped ring (*cis* ring). The protein substrate leaves GroEL upon GroES dissociation, which is induced by ATP binding in the opposite ring (*trans* ring) (**Figure 6b**) (77). Folding intermediates still exposing hydrophobic regions rapidly rebind to GroEL for repeated folding cycles. The exact coordination of the two GroEL rings in the folding cycle is still under investigation (138). Proteins that exceed the size limit of the GroEL-GroES cage may utilize the Hsp70 system for folding (15, 37, 56) or undergo cycles of binding and release from GroEL without GroES encapsulation (77).

In addition to providing an isolated folding environment, other mechanistic elements of the chaperonin cycle contribute to optimizing the rate and yield of the folding process. Repeated events of substrate unfolding in successive binding and release cycles may reverse kinetically trapped states (iterative annealing) (139, 140). Studies have documented unfolding upon binding with Förster resonance energy transfer (FRET) using fluorescence-labeled substrate proteins (130, 140). Additionally, some active unfolding may occur as a result of ATP-dependent movement of the GroEL apical domains (130, 140). However, the significance of iterative annealing is unclear


considering that a single round of substrate binding and encapsulation (using a single-ring mutant of GroEL that binds GroES stably) results in substrate protein folding with equal efficiency and kinetics as achieved through multiple cycles of binding and release (7, 132, 141). Whereas partial unfolding upon initial binding may dissociate nonnative interactions in kinetically trapped folding intermediates, the release of protein from the GroEL apical domains may follow a stepwise mechanism in which less tightly bound hydrophobic regions are released first (130). This stepwise release may contribute to avoiding nonnative interactions during protein collapse.

In addition (or as an alternative) to the mechanisms discussed above, growing experimental and theoretical evidence suggests that the GroEL-GroES cage promotes folding by sterically confining folding intermediates (7, 132, 141–145). This model assigns an active role to the chaperonin cage, as opposed to the view that it functions solely as a passive aggregation-prevention device (146). In support of the confinement model, electron microscopy demonstrated that substrates fully occupy the limited volume of the GroEL-GroES cage (134). The resulting constraints on the substrate protein will inevitably affect its folding energy landscape. Indeed, evidence shows that encapsulation in the GroEL-GroES cage accelerates folding up to tenfold compared with the rate of spontaneous folding (measured without interference of aggregation) (7, 132, 141, 147). In addition to steric confinement, mutational analysis demonstrated that the charged residues of the GroEL cavity wall are critical for the observed acceleration (132, 133, 148). According to molecular dynamics simulations, these polar residues accumulate ordered water molecules in their vicinity, thereby generating a local environment in which a substrate protein may bury exposed hydrophobic residues more effectively (144). This would result in the entropic destabilization of flexible folding intermediates in a manner similar to the role of intramolecular disulfide bonds in promoting the folding of secretory proteins. In this context, it is interesting


a Group I chaperonins



b



c Group II chaperonins



to note that oxidizing folding compartments supporting disulfide bond formation (e.g., the bacterial periplasm or ER) do not contain chaperonins (141).

Obligate GroEL substrates typically have complex α/β or $\alpha + \beta$ domain topologies, such as the $(\beta/\alpha)_8$ TIM barrel fold (15, 46, 53), which are stabilized by many long-range interactions. Such proteins must navigate complex folding energy landscapes and overcome entropic barriers to reach their native states (141); hence, they may benefit from steric confinement during folding. Moreover, obligate GroEL substrates often have relatively low sequence hydrophobicity, consistent with inefficient hydrophobic collapse and population of aggregation-prone intermediates that are recognized by GroEL (46, 141, 149). Recent evidence indicates that GroEL-GroES can also accelerate the folding of proteins containing trefoil-knotted structures, a complex topological arrangement that would plausibly form more efficiently in a confining environment (150). Future studies are needed to define precisely what structural properties distinguish GroEL-dependent from GroEL-independent proteins.

Group II chaperonin TRiC/CCT. Although all chaperonins share a common cylindrical architecture, substantial structural differences exist between group I and group II chaperonins (Figure 6a,c). For example, group II chaperonins have apical protrusions that function as a built-in lid in place of a separate GroES-like cochaperone. Also, group II chaperonins contain eight or nine subunits per ring, which are stacked directly opposite one another in the two rings (77, 151–158). In contrast, each of the seven subunits of group I chaperonins interdigitates between two subunits of the opposite ring. In many cases, group II chaperonins are hetero-oligomeric, containing up to eight paralogous subunits per ring in a defined order in the case of the eukaryotic chaperonin TRiC (159, 160). Similar to GroEL, group II chaperonins also cycle between open and closed states, and substrate encapsulation is essential for folding (161). But in contrast to GroEL, ATP hydrolysis, not ATP binding, triggers cavity closure, and ATP hydrolysis transition state analogs (e.g., ADP-aluminum fluoride) stabilize the closed state (136, 161, 162). During the transition to the closed state, the apical domains of adjacent subunits undergo pairwise

Figure 6

Structure and function of the chaperonin systems. (a) Structure of group I chaperonin. The crystal structures of GroEL (PDB 1SS8) (left) and the asymmetrical GroEL (7 ADP)-GroES complex (PDB 1PF9) (right) appear with GroES shown in green and one subunit of GroEL colored to indicate its domain structure (equatorial nucleotide-binding domain in red; intermediate hinge domain in blue; and the apical substrate and GroES-binding domain in yellow). The conformational differences between the GroEL subunits in the open state (GroEL and *trans* ring of GroEL-GroES complex) and in the closed state (*cis* ring of GroEL-GroES complex) appear in ribbon representations of single subunits (middle). The green spheres represent hydrophobic residues on helices 8 and 9 of the apical domain that are involved in substrate binding in the open conformation and in GroES binding in the closed state. (b) GroEL-GroES reaction cycle. Substrate protein as a collapsed folding intermediate is bound by the open GroEL ring of the asymmetrical GroEL-GroES complex, shown in panel a. Binding of ATP to each of the seven GroEL subunits causes a conformational change in the apical domains, which results in the exposure of the GroES binding residues, allowing substrate encapsulation in the *cis* complex. ADP and GroES dissociate from the opposite ring (*trans* ring) together with the previously bound substrate. The newly encapsulated substrate is free to fold in the GroEL cavity during the time needed to hydrolyze the seven ATP molecules bound to the *cis* ring (~10 s). ATP binding followed by GroES binding to the *trans* ring triggers GroES dissociation from the *cis* ring, releasing the substrate protein. (c) Structure of group II chaperonin. The crystal structures of the open form of the homo-oligomeric *Methanococcus maripaludis* thermosome (PDB 3KFK) (left) and the closed form of TRiC/CCT (PDB 4D8Q) (right) are shown. The eight paralogous TRiC/CCT subunits appear in different colors, showing both homotypic (blue subunits) and heterotypic contacts (green-beige, blue-beige) between the top and bottom rings. One subunit (CCT3 in the case of TRiC) is colored to indicate its domain structure (equatorial nucleotide-binding domain in red; intermediate hinge domain in green; and the apical substrate-binding domain in yellow). The conformational differences between the open and the closed state, taken from the thermosome crystal structures (PDB 3KFK for the open and PDB 3KFB for the closed state) (middle), appear in ribbon representation. In place of a GroES-like cochaperone, the group II chaperonins have an extended apical domain that functions as a built-in lid.

association, resulting in intermediate structures with pseudo-fourfold symmetry (158). Closure is completed when the tips of the helical protrusions form a mixed eight-stranded β -barrel structure around the apical pore.

TRiC interacts with a wide range of cytosolic proteins (49, 50). Prominent substrates include actin and tubulin, which are strictly dependent on the chaperonin for folding (4, 77). A recent crystal structure of bovine TRiC in the open conformation bound to a folding intermediate of tubulin suggests that the substrate initially interacts with loops in the apical and equatorial domains of the TRiC subunits exposed toward the central cavity (157). Helical motifs at the interface between adjacent apical domains have also been implicated in substrate interactions (163). Differences in binding specificities among the different subunits may be important in binding and folding a range of structurally diverse proteins. Accordingly, all eight TRiC subunits are essential in *S. cerevisiae*. Interestingly, the cavity wall exhibits a pronounced segregation of positive and negative surface charges into opposing halves, a feature that might be functionally important (160). The reaction cycle of TRiC is slower than that of GroEL, providing a substantially longer period of protein encapsulation and folding in the cage (164). Also, the iris-like closing mechanism enables the encapsulation of TRiC-dependent domains in the context of large multidomain proteins that cannot be encapsulated in their entirety (164a). Such a mechanism would circumvent the size con-

straints of chaperonin-assisted folding and may have facilitated the evolution of eukaryotic multidomain proteins with complex architectures. TRiC also interacts with N-terminal fragments of mutant huntingtin that contain an expanded polyglutamine repeat sequence (165–168). Binding to TRiC modulates the aggregation properties of this protein and reduces its cytotoxicity.

The Hsp90 System

The Hsp90 chaperone system has a central role in cell regulation. Among its substrates are multiple signaling molecules, which are delivered to Hsp90 by Hsp70 chaperones and other cofactors.


Structure and reaction cycle. The Hsp90 chaperones structurally belong to the gyrase, histidine kinase, and MutL superfamily of ATPases (169). Crystal structures of homodimeric Hsp90 molecules from bacteria (170), yeast (171), and mammals (172) illustrate both open and closed functional states of the chaperone (Figure 7a). In addition, complete or partial structures of Hsp90 in complex with different cochaperones or small molecule inhibitors were obtained by X-ray crystallography and cryoelectron microscopy (81, 169, 171, 173–176). Collectively, these structures reveal a high degree of flexibility in Hsp90 conformations (177), consistent with the diversity of Hsp90 client proteins (45).

Figure 7

Structure and functional cycle of the Hsp90 system. (a) Structure of Hsp90. Crystal structures of the Hsp90 dimer in the ATP-bound closed state (*Saccharomyces cerevisiae*; PDB 2CG9) (left) and the nucleotide-free open state (*E. coli*; PDB 2IOQ) (right) are shown with the nucleotide-binding N-terminal domain in orange, middle domain in green, and C-terminal domain in yellow. (b) Hsp90 reaction cycle. Inactive substrate protein binds to ATP-bound Hsp90. In this state the ATP lids are closed and the N-terminal domains are separated. In the next step, the N-terminal domains dimerize, forming the closed Hsp90 dimer (referred to as a molecular clamp) with twisted subunits. This metastable state is committed to ATP hydrolysis, upon which the N-terminal domains dissociate. The bound substrate protein is conformationally activated as Hsp90 proceeds through the cycle. Cofactors such as Cdc37 and the Hsp90 organizing protein (HOP) slow the ATP hydrolysis step of the cycle, whereas the activator of Hsp90 ATPase (Aha1) enhances ATP hydrolysis. The cofactor p23 stabilizes the closed dimer to slow the release of substrate protein from Hsp90.

Hsp90 consists of three domains: the highly conserved N-terminal domain, the middle domain, and the C-terminal domain (**Figure 7a**) (169). The N-terminal domain contains the ATP binding site, which comprises a two-layer

α/β -sandwich structure. Inhibition of ATP binding and hydrolysis either by mutagenesis of the N domain or by specific inhibitors, such as geldanamycin and radicicol, demonstrated the functional requirement of the Hsp90



Proteostasis network (PN): the collection of cellular components involved in proteostasis maintenance

ATPase (81, 169). The N-terminal domain contains a long, highly charged loop segment with regulatory function located in proximity to the middle domain (178, 179). The middle domain, composed of α - β - α motifs, is essential for interactions with substrate proteins and regulation of ATP hydrolysis (179a, 180, 181). The C-terminal domain of Hsp90 contains the interaction site for the assembly of the functional Hsp90 dimer, a mixed four-helix bundle (173). The extreme C terminus of Hsp90 contains the MEEVD sequence motif, which mediates interactions with numerous cochaperones containing TPR domains (81).

Despite an abundance of structural information, the detailed mechanism by which Hsp90 recognizes kinases and other substrate proteins and facilitates their folding or conformational regulation is not well understood. Similar to other chaperones, the Hsp90 dimer undergoes an ATP-regulated reaction cycle accompanied by extensive structural rearrangements (**Figure 7b**) (100, 182). In this process, the open, V-shaped state of Hsp90 receives inactive client protein and then converts into the closed form, often described as a molecular clamp formed by the N domains (176, 182). This reaction is driven by the combined effects of ATP binding, ATP hydrolysis, posttranslational modifications (183, 184), and interactions with multiple cochaperones (**Figure 7b**). Formation of the closed conformation results in a compaction of the Hsp90 dimer in which the individual subunits twist around each other (171). After hydrolysis the Hsp90 N termini separate, releasing the client protein in an active state (**Figure 7b**). Recent electron microscopy and FRET studies using ATP and nonhydrolyzable ATP analogs suggest that the N-terminally clamped, closed state and the open conformation are in a more dynamic equilibrium than previously believed (185–187).

Cochaperones. Hsp90 intimately cooperates with multiple cochaperones that regulate different steps of the reaction cycle (**Figure 7b**). Cochaperones HOP (St1) and

Cdc37 (p50) stabilize the open conformation of the Hsp90 dimer (81, 174, 175, 188), inhibit ATP hydrolysis, and facilitate substrate protein binding. HOP mediates client transfer from the Hsp70-Hsp40 system, whereas Cdc37 functions as an adapter for kinases. Conversely, p23 (Sba1) couples the Hsp90 ATPase activity to efficient polypeptide dissociation. Two molecules of p23 bind to the N domains of closed Hsp90 and presumably stabilize the ATP-bound state (81, 171, 189). The activator of Hsp90 ATPase (Aha1) binds asymmetrically to the Hsp90 middle domain, stimulating ATP hydrolysis and inducing transition to the closed state (180, 181, 185, 190). In addition, Hsp90 interacts via its C terminus with a range of TPR domain-containing cochaperones. These factors often contain PPIase domains (Cyp40, FKBP51, and FKBP52) and participate in Hsp90-mediated client protein folding (45).

PROTEOSTASIS NETWORK

The successful folding of newly synthesized proteins and their conformational maintenance are essential in sustaining a functional proteome. In addition, the cellular concentration, localization, and activity of each protein must be carefully controlled in response to intrinsic and environmental stimuli. Although research has made major advances in elucidating the mechanics of individual chaperones, we are far from understanding how the various chaperone systems cooperate as a network and function in conjunction with the protein transport and degradation machineries to ensure proteome integrity. The term proteostasis describes this state of healthy proteome balance, whereas proteostasis network (PN) refers to the collection of cellular components involved in proteostasis maintenance (8). Failure of proteostasis is implicated in disease and the deleterious effects of aging (10). Molecular chaperones, through their ability to recognize incorrectly folded proteins, have multiple key roles in the PN (**Figure 8a**).

The PN is regulated by interconnected pathways that respond to specific forms of

cellular stress, including the cytosolic heat shock response (HSR) (191), the unfolded protein response (UPR) in the endoplasmic reticulum (192), and the mitochondrial UPR (193). Additionally, PN regulation is integrated with pathways involved in inflammation, response to oxidative stress, caloric restriction/starvation, and longevity. The PN of mammalian cells consists of ~1,300 different proteins involved in protein biogenesis (~400), conformational maintenance (~300), and degradation (~700), with many proteins being part of more than one pathway (**Figure 8b**). Different cell types vary in their proteostasis capacity and thus in their stress sensitivity and vulnerability to protein aggregation (194, 195).

Significance of Conformational Maintenance

After their initial folding and assembly, many proteins remain reliant on molecular chaperones throughout their cellular lifetime to maintain their functionally active conformations. This is consistent with the notion that proteins with key cellular functions are often structurally dynamic and may be expressed at levels at which they are poorly soluble (14, 46). Many of the chaperone systems discussed in the previous sections function not only in de novo folding but also in conformational maintenance, i.e., they prevent aggregation of misfolded proteins and mediate their refolding. Specific proteins may interact with as many as 25 different types of chaperones throughout their lifetime, as shown in yeast (51). Pulse-chase labeling and quantitative proteomics have described the contributions of the bacterial Hsp70 and chaperonin systems to conformational maintenance (15, 56). Upregulation of chaperones under conditions of conformational stress, such as heat shock or oxidative stress, expands the cellular capacity for the prevention of aggregation. Failure of conformational maintenance is particularly relevant to the onset of age-related degenerative disorders, which typically involve protein aggregation (9).

Degradation

A central feature in the organization of the PN is the tight interconnection of molecular chaperone functions with the pathways of protein degradation, which serve to remove nonfunctional, misfolded, or aggregated proteins that may otherwise disrupt proteostasis. The PN branch of degradation includes the ubiquitin-proteasome system (UPS) and machinery of autophagy (23, 196–200). Approximately 700 proteins are implicated in protein degradation, reflecting the fundamental importance of these pathways in cell regulation and protein homeostasis.

Degradation via the UPS depends on protein unfolding by the 26S proteasome (201) and generally requires that chaperones maintain target proteins in a nonaggregated state. Chaperones cooperate with various E3 ubiquitin ligases in recognizing and targeting misfolded proteins for proteasomal degradation. For example, Hsp70 and Hsp90 cooperate with the U-box-dependent ubiquitin ligase CHIP (C terminus of Hsc70-interacting protein) and a variety of other cofactors (such as BAG1 and BAG3) to ubiquitinate client proteins (23, 86, 198). CHIP interacts with either Hsc70 or Hsp90 via its TPR domain (86, 198, 202). CHIP also cooperates with the E2 enzyme Ubc13-Uev1a to form noncanonical Lys63-linked polyubiquitin chains, which suggests an additional role in targeting proteins for destruction via autophagy (202, 203). BAG1 and BAG3 associate (via their BAG domain) with the NBD of Hsp70 and also interact with CHIP to promote the ubiquitination of Hsp70-bound client proteins (86, 198, 204). BAG1 targets proteins for degradation by the UPS, whereas BAG3 mediates degradation by macroautophagy.


Aggregated proteins that cannot be unfolded for proteasomal degradation may be removed by autophagy and lysosomal/vacuolar degradation. Loss of autophagy causes inclusion body formation and neurodegeneration, even in the absence of additional stress, demonstrating the importance of this pathway for proteostasis (205). Aggregate removal by

UPR: unfolded protein response

UPS:
ubiquitin-proteasome system

autophagy entails the sorting and concentration of small protein aggregates to specific sites in the cytosol (206–209) with the participation of chaperones such as Hsp42 (small Hsp) (209). These deposition sites include the aggresome, to which components mediating autophagic vacuole formation are recruited (206, 210, 211). Additional pathways of autophagy in-

clude chaperone-mediated autophagy (CMA) and chaperone-assisted selective autophagy (CASA) (198, 200, 212). In CMA, Hsc70 and certain cochaperones bind to a KFERQ sequence motif present in approximately 30% of all cytoplasmic proteins. This complex binds the lysosomal membrane protein LAMP2, followed by translocation of the substrate



protein across the lysosomal membrane for degradation (200). Whereas CMA is ubiquitin-independent, CASA uses ubiquitination as a signal for degradation in a pathway that involves the ubiquitin ligase CHIP, Hsc70, and BAG3, as well as the autophagic ubiquitin adaptor p62 (213). CASA is reportedly required for the removal of damaged skeletal muscle proteins such as filamin (213). Notably, chaperones such as Hsp70/Hsc70 act in parallel in the different branches of the PN, and our understanding of how they switch their function between initial folding and conformational maintenance to degradation is still rudimentary.

Proteostasis Collapse in Aging and Disease

As shown in *Caenorhabditis elegans*, *Drosophila*, and the mouse, the ability of cells and tissues to maintain proteostasis declines during aging, concurrent with the capacity to respond to conformational stresses (214–220). Why this is the case is still unclear, but one proposed explanation is that multicellular organisms place less value on protecting the somatic proteome against internal and external stress once propagation of the germ line is certain. The gradual decline in proteostasis capacity would then result in the accumulation of misfolded (or oxidized) proteins, leading to the deposition of aggregates, cellular toxicity, and cell death (214, 219, 221). Accordingly, age is a

universal risk factor for a range of degenerative diseases associated with protein misfolding and aggregate deposition.

The diseases of aberrant protein folding associated with aging are usually categorized as toxic gain-of-function disorders and include type 2 diabetes and the major neurodegenerative diseases (Parkinson's, Huntington's, and Alzheimer's disease, as well as amyotrophic lateral sclerosis) (17, 222). They either occur sporadically or are dominantly inherited. Classical examples of the latter type are Huntington's disease and related disorders, in which the age of onset is inversely correlated with the length of an expanded polyglutamine tract in the disease protein (214). As aggregation propensity increases with polyglutamine length, manifestation of neuronal degeneration may occur when available proteostasis capacity is no longer sufficient to prevent the formation of toxic aggregates. This is supported by experiments demonstrating that the onset of polyglutamine toxicity in *C. elegans* correlates with age-dependent proteostasis decline (214, 215, 217, 220). The accumulation of protein aggregates in turn exerts pressure on the PN, further accelerating its decline. This is exemplified by the observation that toxic polyglutamine repeat proteins interfere with normal protein clearance by the UPS and the conformational maintenance of metastable proteins by the chaperone network (17, 195, 223, 224). Furthermore, toxic protein aggregation also

Figure 8

The proteostasis network. (a) Protein fates in the proteostasis network (PN). The PN integrates chaperone pathways for the folding of newly synthesized proteins, the remodeling of misfolded states, and disaggregation with protein degradation. Panel a adapted and modified from Reference 16. (b) Central role of molecular chaperones. The three branches of the proteostasis network are interconnected by the functions of molecular chaperones. The approximate number of proteins in each branch as well as the number of chaperone components (including cofactors) is indicated. Activation of the cytosolic heat shock response (HSR) and the unfolded protein response (UPR) of the endoplasmic reticulum generally increases proteostasis capacity in all three branches. Various heat shock factor 1 activators can pharmacologically induce the HSR (38, 229, 230). Treatment with guanabenz results in attenuation of translation and increases proteostasis capacity by reducing the load of potentially misfolding proteins (226). The small molecule compound IU-1 inhibits protein deubiquitination and increases degradation by the ubiquitin-proteasome system (UPS) (236). The drug rapamycin activates autophagy by inhibiting the kinase mammalian target of rapamycin (mTOR) (199).

compromises the cellular response to stress stimuli (221).

The Proteostasis Network as a Drug Target

Considering the impact of proteostasis imbalance in age-related degenerative diseases, returning the cell to a more youthful state by pharmacologically boosting proteostasis capacity is a promising therapeutic strategy. Whereas ligand compounds can stabilize specific disease proteins against aggregation (225), activating proteostasis could benefit a wide variety of diseases and might also delay the deleterious effects of aging (8). In principle, we might achieve this by manipulating the three branches of the PN: biogenesis, conformational maintenance, and degradation—either individually or in combination (**Figure 8b**). In the biogenesis branch, attenuation of translation may be beneficial by reducing the load of misfolding proteins. The antihypertensive drug guanabenz demonstrates this; besides being an α_2 receptor antagonist, it also stabilizes the translation initiation factor 2 (eIF2) in its inactive, phosphorylated state (226). Furthermore, boosting cellular chaperone capacity can increase the efficiency of folding or degradation of proteins carrying destabilizing mutations and inhibit their aggregation (227). For example, small molecules (e.g., geldanamycin) that activate heat shock factor 1, the main transcriptional regulator of the cytosolic stress response, increase the effective concentration of cytosolic chaperones and suppress the aggregation of various disease proteins (8, 38, 228–230). This approach is based on multiple lines of evidence demonstrating that overexpression of chaperones such as Hsp70 and Hsp40 prevents the aggregation and toxicity of huntingtin and α -synuclein (38, 231–234). The Hsp70 system acts synergistically with the cytosolic chaperonin TRiC to prevent aggregation of proteins with expanded polyglutamine tracts (165–168). Finally, activating the UPS or

inducing autophagy can increase the clearance of potentially toxic proteins (**Figure 8b**) (212, 235, 236).

OUTLOOK

Studies over the past decade have revealed fascinating insights into the structures of a variety of chaperone systems and the mechanisms by which they assist in protein folding. However, most of these advances are derived from analyses *in vitro*, and consequently, our understanding of how the pathways of folding in the cell differ from those studied in the test tube is still incomplete. Moreover, for most newly synthesized proteins, the relevant quantitative parameters of folding (rate, yield, and overall efficiency) are unknown. Likewise, we are just beginning to understand how the cellular environment influences protein folding and stability and how translation affects the folding process. For example, what is the role of translational pausing in protein folding and trafficking? Much future work will also be directed toward developing an integrated view of the different aspects of the PN, with particular regard to the cooperation between folding and degradation machineries. Solving this problem will require a broad systems biology approach relying on a combination of ribosome profiling, quantitative proteomics, and computational modeling. How do cells react to conformational stresses or proteostasis deficiencies at the proteome level? Which proteins are prone to misfolding, and why do certain proteins aggregate into toxic species whereas others get degraded? How does the composition of the proteome change during aging, what are the signatures of a youthful proteome, and can we find ways to preserve it longer as we age? Addressing these and related questions offers not only a deeper understanding of cell biology but also the prospect of great medical benefits should we be able to intervene in the numerous, presently incurable diseases of protein aggregation and proteostasis deficiency.

SUMMARY POINTS

1. The efficient biogenesis of proteins in the densely crowded cellular environment depends on molecular chaperones to avoid protein misfolding and aggregation.
2. Chaperones promote folding and inhibit aggregation through cycles of binding and release of nonnative proteins (often ATP regulated) that allow kinetic partitioning.
3. Different classes of molecular chaperones may cooperate in sequential pathways.
4. Nascent-chain-binding chaperones prevent misfolding during translation; folding occurs either immediately upon completion of synthesis or after transfer to downstream chaperones, such as the chaperonins, which complete the folding process.
5. Major ATP-dependent chaperone paradigms in the cytosol include the Hsp70 and Hsp90 systems as well as the chaperonins.
6. The chaperonins are cylindrical, ATP-dependent folding machines that encapsulate a single protein chain, allowing it to overcome kinetic folding barriers while being protected against aggregation.
7. Molecular chaperones function as central elements of the large cellular network of proteostasis control, which comprises the protein biogenesis machinery as well as the ubiquitin proteasome and autophagy systems for protein degradation.
8. Understanding the organization of this network and its regulation during stress and aging will help in developing new strategies for the treatment of a range of age-related degenerative diseases associated with protein aggregation.

FUTURE ISSUES

1. How do the pathways of protein folding in the cell differ from those studied in vitro and how does translation affect the folding process?
2. Can we determine the rates, yields, and overall efficiencies of protein folding at the proteome level?
3. How do molecular chaperones of the various branches of the PN cooperate in maintaining proteome integrity?
4. How does the proteome composition change during proteostasis, stress, and aging?
5. What are the mechanisms underlying the cellular toxicity of protein misfolding and aggregation?
6. Can pharmacological chaperone activation serve as a strategy to combat diseases associated with protein misfolding and aggregation?

DISCLOSURE STATEMENT

F.U.H. and M.S.H. are paid consultants of Proteostasis Therapeutics, Inc.

ACKNOWLEDGMENTS

We thank S. Pinkert for providing the bioinformatic analysis of the PN. We apologize that references to original work older than ten years (prior to 2002) could not be cited owing to space limitations. For a description of these contributions, the reader is referred to relevant review articles.

LITERATURE CITED

1. Ellis RJ, Minton AP. 2006. Protein aggregation in crowded environments. *Biol. Chem.* 387:485–97
2. Dobson CM, Sali A, Karplus M. 1998. Protein folding: a perspective from theory and experiment. *Angew. Chem. Int. Ed.* 37:868–93
3. Bartlett AJ, Radford SE. 2009. An expanding arsenal of experimental methods yields an explosion of insights into protein folding mechanisms. *Nat. Struct. Mol. Biol.* 16:582–88
4. Hartl FU. 1996. Molecular chaperones in cellular protein folding. *Nature* 381:571–79
5. Bukau B, Horwich AL. 1998. The Hsp70 and Hsp60 chaperone machines. *Cell* 92:351–66
6. Frydman J. 2001. Folding of newly translated proteins *in vivo*: the role of molecular chaperones. *Annu. Rev. Biochem.* 70:603–47
7. Hartl FU, Hayer-Hartl M. 2002. Molecular chaperones in the cytosol: from nascent chain to folded protein. *Science* 295:1852–58
8. Balch WE, Morimoto RI, Dillin A, Kelly JW. 2008. Adapting proteostasis for disease intervention. *Science* 319:916–19
9. Chiti F, Dobson CM. 2006. Protein misfolding, functional amyloid, and human disease. *Annu. Rev. Biochem.* 75:333–66
10. Morimoto RI. 2008. Proteotoxic stress and inducible chaperone networks in neurodegenerative disease and aging. *Genes Dev.* 22:1427–38
11. Skach WR. 2009. Cellular mechanisms of membrane protein folding. *Nat. Struct. Mol. Biol.* 16:606–12
12. Braakman I, Bulleid NJ. 2011. Protein folding and modification in the mammalian endoplasmic reticulum. *Annu. Rev. Biochem.* 80:71–99
13. Araki K, Nagata K. 2011. Protein folding and quality control in the ER. *Cold Spring Harb. Perspect. Biol.* 3:a007526
14. Dunker AK, Silman I, Uversky VN, Sussman JL. 2008. Function and structure of inherently disordered proteins. *Curr. Opin. Struct. Biol.* 18:756–64
15. Kerner MJ, Naylor DJ, Ishihama Y, Maier T, Chang HC, et al. 2005. Proteome-wide analysis of chaperonin-dependent protein folding in *Escherichia coli*. *Cell* 122:209–20
16. Eichner T, Radford SE. 2011. A diversity of assembly mechanisms of a generic amyloid fold. *Mol. Cell* 43:8–18
17. Hartl FU, Bracher A, Hayer-Hartl M. 2011. Molecular chaperones in protein folding and proteostasis. *Nature* 475:324–32
18. Bolognesi B, Kumita JR, Barros TP, Esbjorner EK, Luheshi LM, et al. 2010. ANS binding reveals common features of cytotoxic amyloid species. *ACS Chem. Biol.* 5:735–40
19. Ignatova Z, Giersch LM. 2004. Monitoring protein stability and aggregation *in vivo* by real-time fluorescent labeling. *Proc. Natl. Acad. Sci. USA* 101:523–28
20. Gershenson A, Giersch LM. 2011. Protein folding in the cell: challenges and progress. *Curr. Opin. Struct. Biol.* 21:32–41
21. McGuffee SR, Elcock AH. 2006. Atomically detailed simulations of concentrated protein solutions: the effects of salt, pH, point mutations, and protein concentration in simulations of 1000-molecule systems. *J. Am. Chem. Soc.* 128:12098–110
22. Hong J, Giersch LM. 2010. Macromolecular crowding remodels the energy landscape of a protein by favoring a more compact unfolded state. *J. Am. Chem. Soc.* 132:10445–52
23. Young JC, Barral JM, Hartl FU. 2003. More than folding: localized functions of cytosolic chaperones. *Trends Biochem. Sci.* 28:541–47

24. Ellis JP, Culviner PH, Cavagnero S. 2009. Confined dynamics of a ribosome-bound nascent globin: cone angle analysis of fluorescence depolarization decays in the presence of two local motions. *Protein Sci.* 18:2003–15
25. Fedyukina DV, Cavagnero S. 2011. Protein folding at the exit tunnel. *Annu. Rev. Biophys.* 40:337–59
26. Woolhead CA, McCormick PJ, Johnson AE. 2004. Nascent membrane and secretory proteins differ in FRET-detected folding far inside the ribosome and in their exposure to ribosomal proteins. *Cell* 116:725–36
27. Lu JL, Deutsch C. 2005. Folding zones inside the ribosomal exit tunnel. *Nat. Struct. Mol. Biol.* 12: 1123–29
28. O'Brien EP, Hsu S-TD, Christodoulou J, Vendruscolo M, Dobson CM. 2010. Transient tertiary structure formation within the ribosome exit port. *J. Am. Chem. Soc.* 132:16928–37
29. Wilson DN, Beckmann R. 2011. The ribosomal tunnel as a functional environment for nascent polypeptide folding and translational stalling. *Curr. Opin. Struct. Biol.* 21:274–82
30. Elcock AH. 2006. Molecular simulations of cotranslational protein folding: fragment stabilities, folding cooperativity, and trapping in the ribosome. *PLoS Comput. Biol.* 2:e98
31. Cabrita LD, Hsu ST, Launay H, Dobson CM, Christodoulou J. 2009. Probing ribosome-nascent chain complexes produced in vivo by NMR spectroscopy. *Proc. Natl. Acad. Sci. USA* 106:22239–44
32. Eichmann C, Preissler S, Riek R, Deuerling E. 2010. Cotranslational structure acquisition of nascent polypeptides monitored by NMR spectroscopy. *Proc. Natl. Acad. Sci. USA* 107:9111–16
33. Ugrinov KG, Clark PL. 2010. Cotranslational folding increases GFP folding yield. *Biophys. J.* 98:1312–20
34. Zhang G, Ignatova Z. 2011. Folding at the birth of the nascent chain: coordinating translation with co-translational folding. *Curr. Opin. Struct. Biol.* 21:25–31
35. Kaiser CM, Goldman DH, Chodera JD, Tinoco I Jr, Bustamante C. 2011. The ribosome modulates nascent protein folding. *Science* 334:1723–27
36. Netzer WJ, Hartl FU. 1998. Protein folding in the cytosol: chaperonin-dependent and -independent mechanisms. *Trends Biochem. Sci.* 23:68–73
37. Agashe VR, Guha S, Chang HC, Genevaux P, Hayer-Hartl M, et al. 2004. Function of trigger factor and DnaK in multidomain protein folding: increase in yield at the expense of folding speed. *Cell* 117:199–209
38. Hartl FU, Hayer-Hartl M. 2009. Converging concepts of protein folding *in vitro* and *in vivo*. *Nat. Struct. Mol. Biol.* 16:574–81
39. Clarke TF IV, Clark PL. 2008. Rare codons cluster. *PLoS ONE* 3:e3412
40. Li G-W, Oh E, Weissman JS. 2012. The anti-Shine-Dalgarno sequence drives translational pausing and codon choice in bacteria. *Nature* 484:538–41
41. O'Brien EP, Vendruscolo M, Dobson CM. 2012. Prediction of variable translation rate effects on cotranslational protein folding. *Nat. Commun.* 3:868
42. Spencer PS, Siller E, Anderson JF, Barral JM. 2012. Silent substitutions predictably alter translation elongation rates and protein folding efficiencies. *J. Mol. Biol.* 422:328–35
43. Brandt F, Etchells SA, Ortiz JO, Elcock AH, Hartl FU, Baumeister W. 2009. The native 3D organization of bacterial polysomes. *Cell* 136:261–71
44. Tokuriki N, Tawfik DS. 2009. Chaperonin overexpression promotes genetic variation and enzyme evolution. *Nature* 459:668–73
45. Taipale M, Jarosz DF, Lindquist S. 2010. Hsp90 at the hub of protein homeostasis: emerging mechanistic insights. *Nat. Rev. Mol. Cell Biol.* 11:515–28
46. Tartaglia GG, Dobson CM, Hartl FU, Vendruscolo M. 2010. Physicochemical determinants of chaperone requirements. *J. Mol. Biol.* 400:579–88
47. Albanese V, Yam AYW, Baughman J, Parnot C, Frydman J. 2006. Systems analyses reveal two chaperone networks with distinct functions in eukaryotic cells. *Cell* 124:75–88
48. McClellan AJ, Xia Y, Deutscher AM, Davis RW, Gerstein M, Frydman J. 2007. Diverse cellular functions of the Hsp90 molecular chaperone uncovered using systems approaches. *Cell* 131:121–35
49. Dekker C, Stirling PC, McCormick EA, Filmore H, Paul A, et al. 2008. The interaction network of the chaperonin CCT. *EMBO J.* 27:1827–39

50. Yam AY, Xia Y, Lin HT, Burlingame A, Gerstein M, Frydman J. 2008. Defining the TRiC/CCT interactome links chaperonin function to stabilization of newly made proteins with complex topologies. *Nat. Struct. Mol. Biol.* 15:1255–62
51. Gong Y, Kakihara Y, Krogan N, Greenblatt J, Emili A, et al. 2009. An atlas of chaperone-protein interactions in *Saccharomyces cerevisiae*: implications to protein folding pathways in the cell. *Mol. Syst. Biol.* 5:275
52. Hirtreiter AM, Calloni G, Forner F, Scheibe B, Puype M, et al. 2009. Differential substrate specificity of group I and group II chaperonins in the archaeon *Methanobarcina mazei*. *Mol. Microbiol.* 74:1152–68
53. Fujiwara K, Ishihama Y, Nakahigashi K, Soga T, Taguchi H. 2010. A systematic survey of *in vivo* obligate chaperonin-dependent substrates. *EMBO J.* 29:1552–64
54. Del Alamo M, Hogan DJ, Pechmann S, Albanese V, Brown PO, Frydman J. 2011. Defining the specificity of cotranslationally acting chaperones by systematic analysis of mRNAs associated with ribosome-nascent chain complexes. *PLoS Biol.* 9:e1001100
55. Oh E, Becker AH, Sandikci A, Huber D, Chaba R, et al. 2011. Selective ribosome profiling reveals the cotranslational chaperone action of trigger factor *in vivo*. *Cell* 147:1295–308
56. Calloni G, Chen T, Schermann SM, Chang H-C, Genevaux P, et al. 2012. DnaK functions as a central hub in the *E. coli* chaperone network. *Cell Rep.* 1:251–64
57. Powers ET, Powers DL, Giersch LM. 2012. FoldEco: a model for proteostasis in *E. coli*. *Cell Rep.* 1:265–76
58. Bukau B, Deuerling E, Pfund C, Craig EA. 2000. Getting newly synthesized proteins into shape. *Cell* 101:119–22
59. Preissler S, Deuerling E. 2012. Ribosome-associated chaperones as key players in proteostasis. *Trends Biochem. Sci.* 37:274–83
60. Kramer G, Rauch T, Rist W, Vorderwülbecke S, Patzelt H, et al. 2002. L23 protein functions as a chaperone docking site on the ribosome. *Nature* 419:171–74
61. Ferbitz L, Maier T, Patzelt H, Bukau B, Deuerling E, Ban N. 2004. Trigger factor in complex with the ribosome forms a molecular cradle for nascent proteins. *Nature* 431:590–96
62. Merz F, Boehringer D, Schaffitzel C, Preissler S, Hoffmann A, et al. 2008. Molecular mechanism and structure of trigger factor bound to the translating ribosome. *EMBO J.* 27:1622–32
63. Kaiser CM, Chang H-C, Agashe VR, Lakshmiapathy SK, Etchells SA, et al. 2006. Real-time observation of trigger factor function on translating ribosomes. *Nature* 444:455–60
64. Eisner G, Koch HG, Beck K, Brunner J, Müller M. 2003. Ligand crowding at a nascent signal sequence. *J. Cell Biol.* 163:35–44
65. Ullers RS, Houben EN, Raine A, ten Hagen-Jongman CM, Ehrenberg M, et al. 2003. Interplay of signal recognition particle and trigger factor at L23 near the nascent chain exit site on the *Escherichia coli* ribosome. *J. Cell Biol.* 161:679–84
66. Bingel-Erlenmeyer R, Kohler R, Kramer G, Sandikci A, Antolić S, et al. 2008. A peptide deformylase-ribosome complex reveals mechanism of nascent chain processing. *Nature* 452:108–13
67. Genevaux P, Keppel F, Schwager F, Langendijk-Genevaux PS, Hartl FU, Georgopoulos C. 2004. *In vivo* analysis of the overlapping functions of DnaK and trigger factor. *EMBO Rep.* 5:195–200
68. Gautschi M, Mun A, Ross S, Rospert S. 2002. A functional chaperone triad on the yeast ribosome. *Proc. Natl. Acad. Sci. USA* 99:4209–14
69. Raue U, Oellerer S, Rospert S. 2007. Association of protein biogenesis factors at the yeast ribosomal tunnel exit is affected by the translational status and nascent polypeptide sequence. *J. Biol. Chem.* 282:7809–16
70. Peisker K, Braun D, Wölfle T, Hentschel J, Fünfschilling U, et al. 2008. Ribosome-associated complex binds to ribosomes in close proximity of Rpl31 at the exit of the polypeptide tunnel in yeast. *Mol. Biol. Cell* 19:5279–88
71. Koplin A, Preissler S, Ilina Y, Koch M, Scior A, et al. 2010. A dual function for chaperones SSB-RAC and the NAC nascent polypeptide-associated complex on ribosomes. *J. Cell Biol.* 189:57–68
72. Wegrzyn RD, Hofmann D, Merz F, Nikolay R, Rauch T, et al. 2006. A conserved motif is prerequisite for the interaction of NAC with ribosomal protein L23 and nascent chains. *J. Biol. Chem.* 281:2847–57
73. Pech M, Spreter T, Beckmann R, Beatrix B. 2010. Dual binding mode of the nascent polypeptide-associated complex reveals a novel universal adapter site on the ribosome. *J. Biol. Chem.* 285:19679–87

74. Niwa T, Kanamori T, Ueda T, Taguchi H. 2012. Global analysis of chaperone effects using a reconstituted cell-free translation system. *Proc. Natl. Acad. Sci. USA* 109:8937–42
75. Kampinga HH, Craig EA. 2010. The Hsp70 chaperone machinery: J proteins as drivers of functional specificity. *Nat. Rev. Mol. Cell Biol.* 11:579–92
76. Tang YC, Chang HC, Hayer-Hartl M, Hartl FU. 2007. Snapshot: molecular chaperones, part II. *Cell* 128:412
77. Horwich AL, Fenton WA, Chapman E, Farr GW. 2007. Two families of chaperonin: physiology and mechanism. *Annu. Rev. Cell Dev. Biol.* 23:115–45
78. Etchells SA, Meyer AS, Yam AY, Roobol A, Miao Y, et al. 2005. The cotranslational contacts between ribosome-bound nascent polypeptides and the subunits of the hetero-oligomeric chaperonin TRiC probed by photocross-linking. *J. Biol. Chem.* 280:28118–26
79. Cuellar J, Martin-Benito J, Scheres SH, Sousa R, Moro F, et al. 2008. The structure of CCT-Hsc70NBD suggests a mechanism for Hsp70 delivery of substrates to the chaperonin. *Nat. Struct. Mol. Biol.* 15:858–64
80. Zhao R, Houry WA. 2007. Molecular interaction network of the Hsp90 chaperone system. *Adv. Exp. Med. Biol.* 594:27–36
81. Young JC, Moarefi I, Hartl FU. 2001. Hsp90: a specialized but essential protein-folding tool. *J. Cell Biol.* 154:267–73
82. Young JC, Hoogenraad NJ, Hartl FU. 2003. Molecular chaperones Hsp90 and Hsp70 deliver preproteins to the mitochondrial import receptor Tom70. *Cell* 112:41–50
83. Sharma K, Vabulas RM, Macek B, Pinkert S, Cox J, et al. 2012. Quantitative proteomics reveals that Hsp90 inhibition preferentially targets kinases and the DNA damage response. *Mol. Cell. Proteomics* 11:M111.014654
84. Neckers L. 2007. Heat shock protein 90: the cancer chaperone. *J. Biosci.* 32:517–30
85. Haslberger T, Bukau B, Mogk A. 2010. Towards a unifying mechanism for ClpB/Hsp104-mediated protein disaggregation and prion propagation. *Biochem. Cell Biol.* 88:63–75
86. Arndt V, Rogon C, Höhfeld J. 2007. To be, or not to be—molecular chaperones in protein degradation. *Cell. Mol. Life Sci.* 64:2525–41
87. Lakshmiapthy SK, Tomic S, Kaiser CM, Chang HC, Genevaux P, et al. 2007. Identification of nascent chain interaction sites on trigger factor. *J. Biol. Chem.* 282:12186–93
88. Lakshmiapthy SK, Gupta R, Pinkert S, Etchells SA, Hartl FU. 2010. Versatility of trigger factor interactions with ribosome-nascent chain complexes. *J. Biol. Chem.* 285:27911–23
89. Baram D, Pyetan E, Sittner A, Auerbach-Nevo T, Bashan A, Yonath A. 2005. Structure of trigger factor binding domain in biologically homologous complex with eubacterial ribosome reveals its chaperone action. *Proc. Natl. Acad. Sci. USA* 102:12017–22
90. Schlünzen F, Wilson DN, Tian P, Harms JM, McInnes SJ, et al. 2005. The binding mode of the trigger factor on the ribosome: implications for protein folding and SRP interaction. *Structure* 13:1685–94
91. Kramer G, Rutkowska A, Wegrzyn RD, Patzelt H, Kurz TA, et al. 2004. Functional dissection of *Escherichia coli* trigger factor: unraveling the function of individual domains. *J. Bacteriol.* 186:3777–84
92. Jakob RP, Zoldak G, Aumüller T, Schmid FX. 2009. Chaperone domains convert prolyl isomerases into generic catalysts of protein folding. *Proc. Natl. Acad. Sci. USA* 106:20282–87
93. Merz F, Hoffmann A, Rutkowska A, Zachmann-Brand B, Bukau B, Deuerling E. 2006. The C-terminal domain of *Escherichia coli* trigger factor represents the central module of its chaperone activity. *J. Biol. Chem.* 281:31963–71
94. Patzelt H, Kramer G, Rauch T, Schönfeld HJ, Bukau B, Deuerling E. 2002. Three-state equilibrium of *Escherichia coli* trigger factor. *Biol. Chem.* 383:1611–19
95. Gupta R, Lakshmiapthy SK, Chang HC, Etchells SA, Hartl FU. 2010. Trigger factor lacking the PPIase domain can enhance the folding of eukaryotic multi-domain proteins in *Escherichia coli*. *FEBS Lett.* 584:3620–24
96. Hoffmann A, Becker AH, Zachmann-Brand B, Deuerling E, Bukau B, Kramer G. 2012. Concerted action of the ribosome and the associated chaperone trigger factor confines nascent polypeptide folding. *Mol. Cell* 48:63–74

97. Liu C-P, Perrett S, Zhou J-M. 2005. Dimeric trigger factor stably binds folding-competent intermediates and cooperates with the DnaK-DnaJ-GrpE chaperone system to allow refolding. *J. Biol. Chem.* 280:13315–20
98. Martinez-Hackert E, Hendrickson WA. 2009. Promiscuous substrate recognition in folding and assembly activities of the trigger factor chaperone. *Cell* 138:923–34
99. Zuiderweg ER, Bertelsen EB, Rousaki A, Mayer MP, Gestwicki JE, Ahmad A. 2013. Allostery in the Hsp70 chaperone proteins. *Top. Curr. Chem.* 328:99–152
100. Mayer MP. 2010. Gymnastics of molecular chaperones. *Mol. Cell* 39:321–31
101. Kityk R, Kopp J, Sinning I, Mayer MP. 2012. Structure and dynamics of the ATP-bound open conformation of Hsp70 chaperones. *Mol. Cell* 48:863–74
102. Deleted in proof
103. Liu QL, Hendrickson WA. 2007. Insights into Hsp70 chaperone activity from a crystal structure of the yeast Hsp110 Sse1. *Cell* 131:106–20
104. Polier S, Dragovic Z, Hartl FU, Bracher A. 2008. Structural basis for the cooperation of Hsp70 and Hsp110 chaperones in protein folding. *Cell* 133:1068–79
105. Schuermann JP, Jiang JW, Cuellar J, Llorea O, Wang L, et al. 2008. Structure of the Hsp110:Hsc70 nucleotide exchange machine. *Mol. Cell* 31:232–43
106. Bertelsen EB, Chang L, Gestwicki JE, Zuiderweg ER. 2009. Solution conformation of wild-type *E. coli* Hsp70 (DnaK) chaperone complexed with ADP and substrate. *Proc. Natl. Acad. Sci. USA* 106:8471–76
107. Mapa K, Sikor M, Kudryavtsev V, Waegemann K, Kalinin S, et al. 2010. The conformational dynamics of the mitochondrial Hsp70 chaperone. *Mol. Cell* 38:89–100
108. Schweizer RS, Aponte RA, Zimmermann S, Weber A, Reinstein J. 2011. Fine tuning of a biological machine: DnaK gains improved chaperone activity by altered allosteric communication and substrate binding. *ChemBioChem* 12:1559–73
109. Swain JF, Dinler G, Sivendran R, Montgomery DL, Stotz M, Gerasch LM. 2007. Hsp70 chaperone ligands control domain association via an allosteric mechanism mediated by the interdomain linker. *Mol. Cell* 26:27–39
110. Smock RG, Rivoire O, Russ WP, Swain JF, Leibler S, et al. 2010. An interdomain sector mediating allostery in Hsp70 molecular chaperones. *Mol. Syst. Biol.* 6:414
111. Zhuravleva A, Gerasch LM. 2011. Allosteric signal transmission in the nucleotide-binding domain of 70-kDa heat shock protein (Hsp70) molecular chaperones. *Proc. Natl. Acad. Sci. USA* 108:6987–92
112. Jiang J, Maes EG, Taylor AB, Wang L, Hinck AP, et al. 2007. Structural basis of J cochaperone binding and regulation of Hsp70. *Mol. Cell* 28:422–33
113. Ahmad A, Bhattacharya A, McDonald RA, Cordes M, Ellington B, et al. 2011. Heat shock protein 70 kDa chaperone/DnaJ cochaperone complex employs an unusual dynamic interface. *Proc. Natl. Acad. Sci. USA* 108:18966–71
114. Svärd M, Biterova EI, Bourhis JM, Guy JE. 2011. The crystal structure of the human co-chaperone p58^{IPK}. *PLoS ONE* 6:e22337
115. Hagiwara M, Maegawa K-I, Suzuki M, Ushioda R, Araki K, et al. 2011. Structural basis of an ERAD pathway mediated by the ER-resident protein disulfide reductase ERdj5. *Mol. Cell* 41:432–44
116. Kabani M, Beckerich JM, Brodsky JL. 2002. Nucleotide exchange factor for the yeast Hsp70 molecular chaperone Ssa1p. *Mol. Cell Biol.* 22:4677–89
117. Steel GJ, Fullerton DM, Tyson JR, Stirling CJ. 2004. Coordinated activation of Hsp70 chaperones. *Science* 303:98–101
118. Dragovic Z, Broadley SA, Shomura Y, Bracher A, Hartl FU. 2006. Molecular chaperones of the Hsp110 family act as nucleotide exchange factors of Hsp70s. *EMBO J.* 25:2519–28
119. Raviol H, Sadlish H, Rodriguez F, Mayer MP, Bukau B. 2006. Chaperone network in the yeast cytosol: Hsp110 is revealed as an Hsp70 nucleotide exchange factor. *EMBO J.* 25:2510–18
120. Shomura Y, Dragovic Z, Chang HC, Tzvetkov N, Young JC, et al. 2005. Regulation of Hsp70 function by HspBP1: Structural analysis reveals an alternate mechanism for Hsp70 nucleotide exchange. *Mol. Cell* 17:367–79
121. Xu Z, Page RC, Gomes MM, Kohli E, Nix JC, et al. 2008. Structural basis of nucleotide exchange and client binding by the Hsp70 cochaperone Bag2. *Nat. Struct. Mol. Biol.* 15:1309–17

122. Arakawa A, Handa N, Ohsawa N, Shida M, Kigawa T, et al. 2010. The C-terminal BAG domain of BAG5 induces conformational changes of the Hsp70 nucleotide-binding domain for ADP-ATP exchange. *Structure* 18:309–19
123. Yan M, Li J, Sha B. 2011. Structural analysis of the Sil1-Bip complex reveals the mechanism for Sil1 to function as a nucleotide-exchange factor. *Biochem. J.* 438:447–55
124. Shorter J. 2011. The mammalian disaggregase machinery: Hsp110 synergizes with Hsp70 and Hsp40 to catalyze protein disaggregation and reactivation in a cell-free system. *PLoS ONE* 6:e26319
125. Rampelt H, Kirstein-Miles J, Nillegoda NB, Chi K, Scholz SR, et al. 2012. Metazoan Hsp70 machines use Hsp110 to power protein disaggregation. *EMBO J.* 31:4221–35
126. Anttonen AK, Mahjneh I, Hämäläinen RH, Lagier-Tourenne C, Kopra O, et al. 2005. The gene disrupted in Marinesco-Sjögren syndrome encodes SIL1, an HSPA5 cochaperone. *Nat. Genet.* 37:1309–11
127. Senderek J, Krieger M, Stendel C, Bergmann C, Moser M, et al. 2005. Mutations in SIL1 cause Marinesco-Sjögren syndrome, a cerebellar ataxia with cataract and myopathy. *Nat. Genet.* 37:1312–14
128. Elad N, Farr GW, Clare DK, Orlova EV, Horwich AL, Saibil HR. 2007. Topologies of a substrate protein bound to the chaperonin GroEL. *Mol. Cell* 26:415–26
129. Horst R, Bertelsen EB, Fiaux J, Wider G, Horwich AL, Wüthrich K. 2005. Direct NMR observation of a substrate protein bound to the chaperonin GroEL. *Proc. Natl. Acad. Sci. USA* 102:12748–53
130. Sharma S, Chakraborty K, Müller BK, Astola N, Tang Y-C, et al. 2008. Monitoring protein conformation along the pathway of chaperonin-assisted folding. *Cell* 133:142–53
131. Hillger F, Hanni D, Nettels D, Geister S, Grandin M, et al. 2008. Probing protein-chaperone interactions with single-molecule fluorescence spectroscopy. *Angew. Chem. Int. Ed.* 47:6184–88
132. Tang Y-C, Chang H-C, Roeben A, Wischnewski D, Wischnewski N, et al. 2006. Structural features of the GroEL-GroES nano-cage required for rapid folding of encapsulated protein. *Cell* 125:903–14
133. Tang YC, Chang HC, Chakraborty K, Hartl FU, Hayer-Hartl M. 2008. Essential role of the chaperonin folding compartment *in vivo*. *EMBO J.* 27:1458–68
134. Clare DK, Bakkes PJ, van Heerikhuizen H, van der Vies SM, Saibil HR. 2009. Chaperonin complex with a newly folded protein encapsulated in the folding chamber. *Nature* 457:107–10
135. Motojima F, Yoshida M. 2010. Polypeptide in the chaperonin cage partly protrudes out and then folds inside or escapes outside. *EMBO J.* 29:4008–19
136. Horovitz A, Willison KR. 2005. Allosteric regulation of chaperonins. *Curr. Opin. Struct. Biol.* 15:646–51
137. Clare DK, Vasishtan D, Stagg S, Quispe J, Farr GW, et al. 2012. ATP-triggered conformational changes delineate substrate-binding and -folding mechanics of the GroEL chaperonin. *Cell* 149:113–23
138. Koike-Takeshita A, Yoshida M, Taguchi H. 2008. Revisiting the GroEL-GroES reaction cycle via the symmetric intermediate implied by novel aspects of the GroEL(D398A) mutant. *J. Biol. Chem.* 283:23774–81
139. Thirumalai D, Lorimer GH. 2001. Chaperonin-mediated protein folding. *Annu. Rev. Biophys. Biomol. Struct.* 30:245–69
140. Lin Z, Madan D, Rye HS. 2008. GroEL stimulates protein folding through forced unfolding. *Nat. Struct. Mol. Biol.* 15:303–11
141. Chakraborty K, Chatila M, Sinha J, Shi Q, Poschner BC, et al. 2010. Chaperonin-catalyzed rescue of kinetically trapped states in protein folding. *Cell* 142:112–22
142. Baumketner A, Jewett A, Shea JE. 2003. Effects of confinement in chaperonin assisted protein folding: rate enhancement by decreasing the roughness of the folding energy landscape. *J. Mol. Biol.* 332:701–13
143. Hayer-Hartl M, Minton AP. 2006. A simple semiempirical model for the effect of molecular confinement upon the rate of protein folding. *Biochemistry* 45:13356–60
144. England J, Lucent D, Pande V. 2008. Rattling the cage: computational models of chaperonin-mediated protein folding. *Curr. Opin. Struct. Biol.* 18:163–69
145. Tian J, Garcia AE. 2011. Simulation studies of protein folding/unfolding equilibrium under polar and nonpolar confinement. *J. Am. Chem. Soc.* 133:15157–64
146. Apetri AC, Horwich AL. 2008. Chaperonin chamber accelerates protein folding through passive action of preventing aggregation. *Proc. Natl. Acad. Sci. USA* 105:17351–55
147. Lin Z, Rye HS. 2004. Expansion and compression of a protein folding intermediate by GroEL. *Mol. Cell* 16:23–34

148. Wang JD, Herman C, Tipton KA, Gross CA, Weissman JS. 2002. Directed evolution of substrate-optimized GroEL/S chaperonins. *Cell* 111:1027–39
149. Raineri E, Ribeca P, Serrano L, Maier T. 2010. A more precise characterization of chaperonin substrates. *Bioinformatics* 26:1685–89
150. Mallam AL, Jackson SE. 2012. Knot formation in newly translated proteins is spontaneous and accelerated by chaperonins. *Nat. Chem. Biol.* 8:147–53
151. Shomura Y, Yoshida T, Iizuka R, Maruyama T, Yohda M, Miki K. 2004. Crystal structures of the group II chaperonin from *Thermococcus* strain KS-1: steric hindrance by the substituted amino acid, and inter-subunit rearrangement between two crystal forms. *J. Mol. Biol.* 335:1265–78
152. Huo Y, Hu Z, Zhang K, Wang L, Zhai Y, et al. 2010. Crystal structure of group II chaperonin in the open state. *Structure* 18:1270–79
153. Cong Y, Baker ML, Jakana J, Woolford D, Miller EJ, et al. 2010. 4.0-Å resolution cryo-EM structure of the mammalian chaperonin TRiC/CCT reveals its unique subunit arrangement. *Proc. Natl. Acad. Sci. USA* 107:4967–72
154. Pereira JH, Ralston CY, Douglas NR, Meyer D, Knee KM, et al. 2010. Crystal structures of a group II chaperonin reveal the open and closed states associated with the protein folding cycle. *J. Biol. Chem.* 285:27958–66
155. Dekker C, Roe SM, McCormack EA, Beuron F, Pearl LH, Willison KR. 2011. The crystal structure of yeast CCT reveals intrinsic asymmetry of eukaryotic cytosolic chaperonins. *EMBO J.* 30:3078–90
156. Zhang J, Ma B, DiMaio F, Douglas NR, Joachimiak LA, et al. 2011. Cryo-EM structure of a group II chaperonin in the prehydrolysis ATP-bound state leading to lid closure. *Structure* 19:633–39
157. Munoz IG, Yebenes H, Zhou M, Mesa P, Serna M, et al. 2011. Crystal structure of the open conformation of the mammalian chaperonin CCT in complex with tubulin. *Nat. Struct. Mol. Biol.* 18:14–19
158. Cong Y, Schroder GF, Meyer AS, Jakana J, Ma B, et al. 2012. Symmetry-free cryo-EM structures of the chaperonin TRiC along its ATPase-driven conformational cycle. *EMBO J.* 31:720–30
159. Kalisman N, Adams CM, Levitt M. 2012. Subunit order of eukaryotic TRiC/CCT chaperonin by cross-linking, mass spectrometry, and combinatorial homology modeling. *Proc. Natl. Acad. Sci. USA* 109:2884–89
160. Leitner A, Joachimiak LA, Bracher A, Mönkemeyer L, Walzthoeni T, et al. 2012. The molecular architecture of the eukaryotic chaperonin TRiC/CCT. *Structure* 20:814–25
161. Douglas NR, Reissmann S, Zhang J, Chen B, Jakana J, et al. 2011. Dual action of ATP hydrolysis couples lid closure to substrate release into the group II chaperonin chamber. *Cell* 144:240–52
162. Meyer AS, Gillespie JR, Walther D, Millet IS, Doniach S, Frydman J. 2003. Closing the folding chamber of the eukaryotic chaperonin requires the transition state of ATP hydrolysis. *Cell* 113:369–81
163. Spiess C, Miller EJ, McClellan AJ, Frydman J. 2006. Identification of the TRiC/CCT substrate binding sites uncovers the function of subunit diversity in eukaryotic chaperonins. *Mol. Cell* 24:25–37
164. Reissmann S, Parnot C, Booth CR, Chiu W, Frydman J. 2007. Essential function of the built-in lid in the allosteric regulation of eukaryotic and archaeal chaperonins. *Nat. Struct. Mol. Biol.* 14:432–40
- 164a. Rübbmann F, Stemp MJ, Mönkemeyer L, Etchells SA, Bracher A, Hartl FU. 2012. Folding of large multidomain proteins by partial encapsulation in the chaperonin TRiC/CCT. *Proc. Natl. Acad. Sci. USA* 109:21208–15
165. Kitamura A, Kubota H, Pack CG, Matsumoto G, Hirayama S, et al. 2006. Cytosolic chaperonin prevents polyglutamine toxicity with altering the aggregation state. *Nat. Cell Biol.* 8:1163–70
166. Behrends C, Langer CA, Boteva R, Böttcher UM, Stemp MJ, et al. 2006. Chaperonin TRiC promotes the assembly of polyQ expansion proteins into nontoxic oligomers. *Mol. Cell* 23:887–97
167. Tam S, Geller R, Spiess C, Frydman J. 2006. The chaperonin TRiC controls polyglutamine aggregation and toxicity through subunit-specific interactions. *Nat. Cell Biol.* 8:1155–62
168. Tam S, Spiess C, Auyeung W, Joachimiak L, Chen B, et al. 2009. The chaperonin TRiC blocks a huntingtin sequence element that promotes the conformational switch to aggregation. *Nat. Struct. Mol. Biol.* 16:1279–85
169. Pearl LH, Prodromou C. 2006. Structure and mechanism of the Hsp90 molecular chaperone machinery. *Annu. Rev. Biochem.* 75:271–94

170. Shiu AK, Harris SF, Southworth DR, Agard DA. 2006. Structural analysis of *E. coli* Hsp90 reveals dramatic nucleotide-dependent conformational rearrangements. *Cell* 127:329–40
171. Ali MM, Roe SM, Vaughan CK, Meyer P, Panaretou B, et al. 2006. Crystal structure of an Hsp90-nucleotide-p23/Sba1 closed chaperone complex. *Nature* 440:1013–17
172. Dollins DE, Warren JJ, Immormino RM, Gewirth DT. 2007. Structures of Grp94-nucleotide complexes reveal mechanistic differences between the Hsp90 chaperones. *Mol. Cell* 28:41–56
173. Harris SF, Shiu AK, Agard DA. 2004. The crystal structure of the carboxy-terminal dimerization domain of htpG, the *Escherichia coli* Hsp90, reveals a potential substrate binding site. *Structure* 12:1087–97
174. Roe SM, Ali MM, Meyer P, Vaughan CK, Panaretou B, et al. 2004. The mechanism of Hsp90 regulation by the protein kinase-specific cochaperone p50^{cdc37}. *Cell* 116:87–98
175. Vaughan CK, Gohlke U, Sobott F, Good VM, Ali MMU, et al. 2006. Structure of an Hsp90-Cdc37-Cdk4 complex. *Mol. Cell* 23:697–707
176. Southworth DR, Agard DA. 2011. Client-loading conformation of the Hsp90 molecular chaperone revealed in the cryo-EM structure of the human Hsp90:HOP complex. *Mol. Cell* 42:771–81
177. Krukenberg KA, Forster F, Rice LM, Sali A, Agard DA. 2008. Multiple conformations of *E. coli* Hsp90 in solution: insights into the conformational dynamics of Hsp90. *Structure* 16:755–65
178. Hainzl O, Lapina MC, Buchner J, Richter K. 2009. The charged linker region is an important regulator of Hsp90 function. *J. Biol. Chem.* 284:22559–67
179. Tsutsumi S, Mollapour M, Graf C, Lee C-T, Scroggins BT, et al. 2009. Hsp90 charged-linker truncation reverses the functional consequences of weakened hydrophobic contacts in the N domain. *Nat. Struct. Mol. Biol.* 16:1141–47
- 179a. Meyer P, Prodromou C, Hu B, Vaughan C, Roe SM, et al. Structural and functional analysis of the middle segment of hsp90: implications for ATP hydrolysis and client protein and cochaperone interactions. *Mol. Cell* 11:647–58
180. Retzlaff M, Hagn F, Mitschke L, Hessling M, Gugel F, et al. 2010. Asymmetric activation of the Hsp90 dimer by its cochaperone Aha1. *Mol. Cell* 37:344–54
181. Koulov AV, Lapointe P, Lu B, Razvi A, Coppinger J, et al. 2010. Biological and structural basis for Aha1 regulation of Hsp90 ATPase activity in maintaining proteostasis in the human disease cystic fibrosis. *Mol. Biol. Cell* 21:871–84
182. Li J, Soroka J, Buchner J. 2012. The Hsp90 chaperone machinery: conformational dynamics and regulation by co-chaperones. *Biochim. Biophys. Acta* 1823:624–35
183. Mollapour M, Tsutsumi S, Truman AW, Xu W, Vaughan CK, et al. 2011. Threonine 22 phosphorylation attenuates Hsp90 interaction with cochaperones and affects its chaperone activity. *Mol. Cell* 41:672–81
184. Soroka J, Wandinger SK, Mäusbacher N, Schreiber T, Richter K, et al. 2012. Conformational switching of the molecular chaperone Hsp90 via regulated phosphorylation. *Mol. Cell* 45:517–28
185. Hessling M, Richter K, Buchner J. 2009. Dissection of the ATP-induced conformational cycle of the molecular chaperone Hsp90. *Nat. Struct. Mol. Biol.* 16:287–93
186. Southworth DR, Agard DA. 2008. Species-dependent ensembles of conserved conformational states define the Hsp90 chaperone ATPase cycle. *Mol. Cell* 32:631–40
187. Mickler M, Hessling M, Ratzke C, Buchner J, Hugel T. 2009. The large conformational changes of Hsp90 are only weakly coupled to ATP hydrolysis. *Nat. Struct. Mol. Biol.* 16:281–86
188. Schmid AB, Lagleder S, Gräwert MA, Röhl A, Hagn F, et al. 2012. The architecture of functional modules in the Hsp90 co-chaperone Sti1/HOP. *EMBO J.* 31:1506–17
189. Richter K, Walter S, Buchner J. 2004. The co-chaperone Sba1 connects the ATPase reaction of Hsp90 to the progression of the chaperone cycle. *J. Mol. Biol.* 342:1403–13
190. Panaretou B, Siligardi G, Meyer P, Maloney A, Sullivan JK, et al. 2002. Activation of the ATPase activity of Hsp90 by the stress-regulated cochaperone Aha1. *Mol. Cell* 10:1307–18
191. Anckar J, Sistonen L. 2011. Regulation of HSF1 function in the heat stress response: implications in aging and disease. *Annu. Rev. Biochem.* 80:1089–115
192. Walter P, Ron D. 2011. The unfolded protein response: from stress pathway to homeostatic regulation. *Science* 334:1081–86

193. Haynes CM, Ron D. 2010. The mitochondrial UPR—protecting organelle protein homeostasis. *J. Cell Sci.* 123:3849–55
194. Kern A, Ackermann B, Clement AM, Duerk H, Behl C. 2010. HSF1-controlled and age-associated chaperone capacity in neurons and muscle cells of *C. elegans*. *PLoS ONE* 5:e8568
195. Gupta R, Kasturi P, Bracher A, Loew C, Zheng M, et al. 2011. Firefly luciferase mutants as sensors of proteome stress. *Nat. Methods* 8:879–84
196. Hershko A, Ciechanover A. 1998. The ubiquitin system. *Annu. Rev. Biochem.* 67:425–79
197. Finley D. 2009. Recognition and processing of ubiquitin-protein conjugates by the proteasome. *Annu. Rev. Biochem.* 78:477–513
198. Kettern N, Dreiseidler M, Tawo R, Höhfeld J. 2010. Chaperone-assisted degradation: multiple paths to destruction. *Biol. Chem.* 391:481–89
199. Rubinsztein DC, Marino G, Kroemer G. 2011. Autophagy and aging. *Cell* 146:682–95
200. Arias E, Cuervo AM. 2011. Chaperone-mediated autophagy in protein quality control. *Curr. Opin. Cell Biol.* 23:184–89
201. Prakash S, Matouschek A. 2004. Protein unfolding in the cell. *Trends Biochem. Sci.* 29:593–600
202. Zhang MH, Windheim M, Roe SM, Peggy M, Cohen P, et al. 2005. Chaperoned ubiquitylation—crystal structures of the CHIP U box E3 ubiquitin ligase and a CHIP-Ubc13-Uev1a complex. *Mol. Cell* 20:525–38
203. Xu Z, Kohli E, Devlin KI, Bold M, Nix JC, Misra S. 2008. Interactions between the quality control ubiquitin ligase CHIP and ubiquitin conjugating enzymes. *BMC Struct. Biol.* 8:26
204. Gamerdinger M, Hajieva P, Kaya AM, Wolfrum U, Hartl FU, Behl C. 2009. Protein quality control during aging involves recruitment of the macroautophagy pathway by BAG3. *EMBO J.* 28:889–901
205. Komatsu M, Waguri S, Chiba T, Murata S, Iwata J, et al. 2006. Loss of autophagy in the central nervous system causes neurodegeneration in mice. *Nature* 441:880–84
206. Kopito RR. 2000. Aggresomes, inclusion bodies and protein aggregation. *Trends Cell Biol.* 10:524–30
207. Kaganovich D, Kopito R, Frydman J. 2008. Misfolded proteins partition between two distinct quality control compartments. *Nature* 454:1088–95
208. Tyedmers J, Mogk A, Bukau B. 2010. Cellular strategies for controlling protein aggregation. *Nat. Rev. Mol. Cell Biol.* 11:777–88
209. Specht S, Miller SBM, Mogk A, Bukau B. 2011. Hsp42 is required for sequestration of protein aggregates into deposition sites in *Saccharomyces cerevisiae*. *J. Cell Biol.* 195:617–29
210. Iwata A, Riley BE, Johnston JA, Kopito RR. 2005. HDAC6 and microtubules are required for autophagic degradation of aggregated huntingtin. *J. Biol. Chem.* 280:40282–92
211. Iwata A, Christianson JC, Bucci M, Ellerby LM, Nukina N, et al. 2005. Increased susceptibility of cytoplasmic over nuclear polyglutamine aggregates to autophagic degradation. *Proc. Natl. Acad. Sci. USA* 102:13135–40
212. Kon M, Cuervo AM. 2010. Chaperone-mediated autophagy in health and disease. *FEBS Lett.* 584:1399–404
213. Arndt V, Dick N, Tawo R, Dreiseidler M, Wenzel D, et al. 2010. Chaperone-assisted selective autophagy is essential for muscle maintenance. *Curr. Biol.* 20:143–48
214. Morley JF, Brignull HR, Weyers JJ, Morimoto RI. 2002. The threshold for polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and influenced by aging in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* 99:10417–22
215. Morley JF, Morimoto RI. 2004. Regulation of longevity in *Caenorhabditis elegans* by heat shock factor and molecular chaperones. *Mol. Biol. Cell* 15:657–64
216. Cohen E, Bieschke J, Perciavalle RM, Kelly JW, Dillin A. 2006. Opposing activities protect against age-onset proteotoxicity. *Science* 313:1604–10
217. Ben-Zvi A, Miller EA, Morimoto RI. 2009. Collapse of proteostasis represents an early molecular event in *Caenorhabditis elegans* aging. *Proc. Natl. Acad. Sci. USA* 106:14914–19
218. Cohen E, Paulsson JF, Blinder P, Burstyn-Cohen T, Du D, et al. 2009. Reduced IGF-1 signaling delays age-associated proteotoxicity in mice. *Cell* 139:1157–69
219. David DC, Ollikainen N, Trinidad JC, Cary MP, Burlingame AL, Kenyon C. 2010. Widespread protein aggregation as an inherent part of aging in *C. elegans*. *PLoS Biol.* 8:e1000450

220. Demontis F, Perrimon N. 2010. FOXO/4E-BP signaling in *Drosophila* muscles regulates organism-wide proteostasis during aging. *Cell* 143:813–25
221. Olzscha H, Schermann SM, Woerner AC, Pinkert S, Hecht MH, et al. 2011. Amyloid-like aggregates sequester numerous metastable proteins with essential cellular functions. *Cell* 144:67–78
222. Orr HT, Zoghbi HY. 2007. Trinucleotide repeat disorders. *Annu. Rev. Neurosci.* 30:575–621
223. Gidalevitz T, Ben-Zvi A, Ho KH, Brignull H, Morimoto RI. 2006. Progressive disruption of cellular protein folding in models of polyglutamine diseases. *Science* 311:1471–74
224. Hipp MS, Patel CN, Bersuker K, Riley BE, Kaiser SE, et al. 2012. Indirect inhibition of 26S proteasome activity in a cellular model of Huntington’s disease. *J. Cell Biol.* 196:573–87
225. Bulawa CE, Connelly S, Devit M, Wang L, Weigel C, et al. 2012. Tafamidis, a potent and selective transthyretin kinetic stabilizer that inhibits the amyloid cascade. *Proc. Natl. Acad. Sci. USA* 109:9629–34
226. Tsayler P, Harding HP, Ron D, Bertolotti A. 2011. Selective inhibition of a regulatory subunit of protein phosphatase 1 restores proteostasis. *Science* 332:91–94
227. Nagai Y, Fujikake N, Popiel HA, Wada K. 2010. Induction of molecular chaperones as a therapeutic strategy for the polyglutamine diseases. *Curr. Pharm. Biotechnol.* 11:188–97
228. Westerheide SD, Morimoto RI. 2005. Heat shock response modulators as therapeutic tools for diseases of protein conformation. *J. Biol. Chem.* 280:33097–100
229. Mu TW, Ong DS, Wang YJ, Balch WE, Yates JR, et al. 2008. Chemical and biological approaches synergize to ameliorate protein-folding diseases. *Cell* 134:769–81
230. Calamini B, Silva MC, Madoux F, Hutt DM, Khanna S, et al. 2012. Small-molecule proteostasis regulators for protein conformational diseases. *Nat. Chem. Biol.* 8:185–96
231. Schaffar G, Breuer P, Boteva R, Behrends C, Tzvetkov N, et al. 2004. Cellular toxicity of polyglutamine expansion proteins: mechanism of transcription factor deactivation. *Mol. Cell* 15:95–105
232. Lotz GP, Legleiter J, Aron R, Mitchell EJ, Huang SY, et al. 2010. Hsp70 and Hsp40 functionally interact with soluble mutant huntingtin oligomers in a classic ATP-dependent reaction cycle. *J. Biol. Chem.* 285:38183–93
233. Auluck PK, Chan HY, Trojanowski JQ, Lee VM, Bonini NM. 2002. Chaperone suppression of α -synuclein toxicity in a *Drosophila* model for Parkinson’s disease. *Science* 295:865–68
234. Hageman J, Rujano MA, van Waarde MAWH, Kakkar V, Dirks RP, et al. 2010. A DNAJB chaperone subfamily with HDAC-dependent activities suppresses toxic protein aggregation. *Mol. Cell* 37:355–69
235. Sarkar S, Ravikumar B, Floto RA, Rubinsztein DC. 2009. Rapamycin and mTOR-independent autophagy inducers ameliorate toxicity of polyglutamine-expanded huntingtin and related proteinopathies. *Cell Death Differ.* 16:46–56
236. Lee B-H, Lee MJ, Park S, Oh DC, Elsasser S, et al. 2010. Enhancement of proteasome activity by a small-molecule inhibitor of USP14. *Nature* 467:179–84
237. Jahn TR, Radford SE. 2005. The yin and yang of protein folding. *FEBS J.* 272:5962–70

Contents

Prefatory

Christian Raetz: Scientist and Friend Extraordinaire

- William Dowhan, Hiroshi Nikaido, JoAnne Stubbe, John W. Kozarich,
William T. Wickner, David W. Russell, Teresa A. Garrett, Kathryn Brozek,
and Paul Modrich* 1

Recent Advances in Biochemistry

Mechanisms for Initiating Cellular DNA Replication

- Alessandro Costa, Iris V. Hood, and James M. Berger* 25

The Chromatin Response to DNA Breaks: Leaving a Mark on Genome Integrity

- Godelieve Smeenk and Haico van Attikum* 55

Readout of Epigenetic Modifications

- Dinshaw J. Patel and Zhanxin Wang* 81

Flap Endonuclease 1

- Lata Balakrishnan and Robert A. Bambara* 119

New Mechanistic and Functional Insights into DNA Topoisomerases

- Stefanie Hartman Chen, Nei-Li Chan, and Tao-shih Hsieh* 139

Arrest Peptides: *Cis*-Acting Modulators of Translation

- Koreaki Ito and Shinobu Chiba* 171

Structural Basis of the Translational Elongation Cycle

- Rebecca M. Voorhees and V. Ramakrishnan* 203

CRISPR-Mediated Adaptive Immune Systems in Bacteria and Archaea

- Rotem Sorek, C. Martin Lawrence, and Blake Wiedenheft* 237

Correlating Structure and Energetics in Protein-Ligand Interactions:

Paradigms and Paradoxes

- Stephen F. Martin and John H. Clements* 267

Extracellular Chaperones and Proteostasis

- Amy R. Wyatt, Justin J. Yerbury, Heath Ecroyd, and Mark R. Wilson* 295

Molecular Chaperone Functions in Protein Folding and Proteostasis <i>Yujin E. Kim, Mark S. Hipp, Andreas Bracher, Manajit Hayer-Hartl, and F. Ulrich Hartl</i>	323
Sumoylation: A Regulatory Protein Modification in Health and Disease <i>Annette Flotho and Frauke Melchior</i>	357
Ubiquitin Ligases and Cell Cycle Control <i>Leonardo K. Teixeira and Steven I. Reed</i>	387
Molecular Architecture and Assembly of the Eukaryotic Proteasome <i>Robert J. Tomko Jr. and Mark Hochstrasser</i>	415
Design of Protein Catalysts <i>Donald Hilvert</i>	447
Hydrogen Tunneling Links Protein Dynamics to Enzyme Catalysis <i>Judith P. Klinman and Amnon Kohen</i>	471
Methylerythritol Phosphate Pathway of Isoprenoid Biosynthesis <i>Lishan Zhao, Wei-chen Chang, Youli Xiao, Hung-wen Liu, and Pinghua Liu</i>	497
Posttranslational Biosynthesis of the Protein-Derived Cofactor Tryptophan Tryptophylquinone <i>Victor L. Davidson and Carrie M. Wilmot</i>	531
Mitochondrial Complex I <i>Judy Hirst</i>	551
Photosystem II: The Reaction Center of Oxygenic Photosynthesis <i>David J. Vinyard, Gennady M. Ananyev, and G. Charles Dismukes</i>	577
The Voltage-Gated Calcium Channel Functions as the Molecular Switch of Synaptic Transmission <i>Daphne Atlas</i>	607
Sphingosine-1-Phosphate and Its Receptors: Structure, Signaling, and Influence <i>Hugh Rosen, Raymond C. Stevens, Michael Hanson, Edward Roberts, Michael B.A. Oldstone</i>	637
Membrane Fission Reactions of the Mammalian ESCRT Pathway <i>John McCullough, Jeremy A. Colf, and Wesley I. Sundquist</i>	663
Signal Recognition Particle: An Essential Protein-Targeting Machine <i>David Akopian, Kuang Shen, Xin Zhang, and Shu-ou Shan</i>	693
Peroxisome Formation and Maintenance Are Dependent on the Endoplasmic Reticulum <i>Henk F. Tabak, Ineke Braakman, and Adabella van der Zand</i>	723

Systemic Amyloidoses <i>Luis M. Blancas-Mejía and Marina Ramirez-Alvarado</i>	745
Nanobodies: Natural Single-Domain Antibodies <i>Serge Muyldermans</i>	775

Indexes

Cumulative Index of Contributing Authors, Volumes 78–82	799
Cumulative Index of Article Titles, Volumes 78–82	803

Errata

An online log of corrections to *Annual Review of Biochemistry* articles may be found at
<http://biochem.annualreviews.org/errata.shtml>