
Sum Sensitivity Proofs

May 21, 2020

Definition 1. A sum query on database x of size n is defined to be

$$s(x) = \sum_{i=1}^n x_i.$$

1 NEIGHBORING DEFINITION: CHANGE ONE

1.1 ℓ_1 -sensitivity

Theorem 1. Say the space of datapoints \mathcal{X} is bounded above by M and bounded below by m . Then s over \mathcal{X}^n has ℓ_1 -sensitivity in the change-one model bounded above by $M - m$.

Proof. Say X and X' are neighboring databases which differ at data-point x_j . Then

$$\begin{aligned} \Delta s &= \max_{X, X'} |s(X) - s(X')| \\ &= \max_{X, X'} \left| \left(\sum_{\{i \in [n] | i \neq j\}} x_i \right) + x_j - \left(\sum_{\{i \in [n] | i \neq j\}} x'_i \right) + x'_j \right| \\ &= \max_{X, X'} |x_j - x'_j| \\ &\leq M - m. \end{aligned}$$

□

1.2 ℓ_2 -sensitivity

Theorem 2. Say the space of datapoints \mathcal{X} is bounded above by M and bounded below by m . Then s over \mathcal{X}^n has ℓ_1 -sensitivity in the change-one model bounded above by $M - m$.

Proof. This follows the logic of the above proof. □

2 NEIGHBORING DEFINITION: ADD/DROP ONE

2.1 ℓ_1 -sensitivity

Theorem 3. *Say the space of datapoints \mathcal{X} is bounded above by M and bounded below by m . Then s has ℓ_1 -sensitivity in the add/drop-one model bounded above by $\max(|m|, |M|)$.*

Proof. For notational ease, let n always refer to the size of database x . We must consider both adding and removing an element from x . First, consider adding a point:

Let $X' = X \cup x$. Without loss of generality, assume the point added is the $(n+1)^{\text{th}}$ element of database x' . Note that

$$\begin{aligned} |s(X) - s(X')| &= \left| \sum_{i=1}^n x_i - \sum_{i=1}^{n+1} x_i \right| \\ &= \left| \sum_{i=1}^n x_i - \left(\sum_{i=1}^n x_i \right) - x \right| \\ &= x \\ &\leq \max(|m|, |M|). \end{aligned}$$

Second, consider removing a point:

Let $X' = X \setminus \{x\}$. Without loss of generality assume that the point subtracted is the n^{th} element of database x .

$$\begin{aligned} |s(X) - s(X')| &= \left| \sum_{i=1}^n x_i - \sum_{i=1}^{n-1} x_i \right| \\ &= \left| \left(\sum_{i=1}^{n-1} x_i \right) + x - \sum_{i=1}^{n-1} x_i \right| \\ &= x \\ &\leq \max(|m|, |M|). \end{aligned}$$

□

2.2 ℓ_2 -sensitivity

Theorem 4. *Say the space of datapoints \mathcal{X} is bounded above by M and bounded below by m . Then s has ℓ_2 -sensitivity in the add/drop-one model bounded above by $\max(|m|, |M|)$.*

Proof. This follows the same logic as the above proof. □