

fn integrate_discrete_laplacian_tail

Michael Shoemate

August 13, 2024

This proof resides in “**contrib**” because it has not completed the vetting process.

Proof for `integrate_discrete_laplace_tail`.

Definition 0.1. Define $Y \sim \mathcal{L}_{\mathbb{Z}}(0, s)$, a random variable following the discrete laplace distribution:

$$\forall x \in \mathbb{Z} \quad \Pr[X = x] = \frac{e^{1/s} - 1}{e^{1/s} + 1} e^{-|x|/s} \quad (1)$$

Theorem 0.2. Assume $X \sim \mathcal{L}_{\mathbb{Z}}(0, s)$, and $t > 0$.

$$\alpha = P[X > t] = \frac{e^{-t/s}}{e^{-1/s} + 1} \quad (2)$$

Proof.

$$\begin{aligned} \alpha &= P[X > t] \\ &= \sum_{x=t+1}^{\infty} \frac{e^{1/s} - 1}{e^{1/s} + 1} e^{-|x|/s} \\ &= \frac{e^{1/s} - 1}{e^{1/s} + 1} \sum_{x=t+1}^{\infty} e^{-x/s} && \text{since } t > 0 \\ &= \frac{e^{1/s} - 1}{e^{1/s} + 1} \frac{e^{(1-(t+1))/s}}{e^{1/s} - 1} && \text{since } \sum_{x=t}^{\infty} p^x = \frac{p^t}{1-p} \text{ if } |p| < 1, \text{ let } p = e^{1/s} \\ &= \frac{e^{-t/s}}{e^{1/s} + 1} \end{aligned}$$

□