# fn sample_geometric_exp_slow

Michael Shoemate

May 5, 2024

This proof resides in **"contrib"** because it has not completed the vetting process.

Proves soundness of `fn sample_geometric_exp_slow` in `mod.rs` at commit 0be3ab3e6 (outdated[1]). This proof is an adaptation of subsection 5.2 of [CKS20].

### Vetting history

- Pull Request #519

## 1 Hoare Triple

### Precondition

$\texttt{x} \in \mathbb{Q} \wedge \texttt{x} > 0$

### Pseudocode

```
def sample_geometric_exp_slow(x) -> int:
    k = 0
    while True:
        if sample_bernoulli_exp(x):
            k += 1
        else:
            return k
```

### Postcondition

For any setting of the input parameter `x` such that the given preconditions hold, `sample_geometric_exp_slow` either returns `Err(e)` due to a lack of system entropy, or `Ok(out)`, where `out` is distributed as $Geometric(1 - exp(-x))$.

## 2 Proof

Assume the preconditions are met.

**Lemma 2.1.** `sample_geometric_exp_slow` only returns `Err(e)` when there is a lack of system entropy.

[1] See new changes with `git diff 0be3ab3e6..8241be0 rust/src/traits/samplers/cks20/mod.rs`

*Proof.* The preconditions on `x` satisfy the preconditions on `sample_bernoulli_exp`, so by its definition, it only returns an error if there is a lack of system entropy. The only source of errors is from this function, therefore `sample_geometric_exp_slow` only returns `Err(e)` when there is a lack of system entropy. □

**Theorem 2.2.** [CKS20] If the outcome of `sample_geometric_exp_slow` is `Ok(out)`, then `out` is distributed as $Geometric(1 - exp(-x))$. That is, $P[\texttt{out} = k] = exp(-x)(1 - exp(-x))^k$

*Proof.* The distribution of the $i^{th}$ boolean returned on line 4 is $B_i \sim Bernoulli(exp(x))$, because the preconditions on `x` satisfy the preconditions for `sample_bernoulli_exp`.

$$\begin{aligned} P[\texttt{out} = k] &= P[B_1 = B_2 = ... = B_k = \bot \wedge B_{k+1} = \top] \\ &= P[B_{k+1} = \top] \prod_{i=1}^{k} P[B_i = \bot] && \text{All } B_i \text{ are independent.} \\ &= exp(-x)(1 - exp(-x))^k \end{aligned}$$

□

*Proof.* 1 holds by 2.1 and 2.2. □

# References


[CKS20] Clément L. Canonne, Gautam Kamath, and Thomas Steinke. The discrete gaussian for differential privacy. *CoRR*, abs/2004.00010, 2020.