fn make_sized_quantile_score_candidates

Michael Shoemate

March 10, 2023

This proof resides in **"contrib"** because it has not completed the vetting process.

Vetting History

- Pull Request #456

Proves soundness of `make_sized_quantile_score_candidates` in `mod.rs` at commit f5bb719 (outdated[1]). `make_sized_quantile_score_candidates` returns a Transformation that takes a numeric vector database and a vector of numeric quantile candidates, and returns a vector of scores, where higher scores correspond to more accurate candidates.

It is the same as the unsized version, `make_quantile_score_candidates`, but with a sized input domain, and different stability map. The unsized version has a more thorough explanation of the utility function.

1 Hoare Triple

Precondition

- `TIA` (input atom type) is a type with trait `Number`.
- `TOA` (output atom type) is a type with trait `Float`.

Function

```python
def make_sized_quantile_score_candidates(size: usize, candidates: List[TIA], alpha: TOA):
    for i in range(len(candidates) - 1):
        assert candidates[i] > candidates[i + 1]

    def function(arg: List[TIA]):
        return score(arg, candidates, alpha)

    def stability_map(d_in: IntDistance):
        return TOA.inf_cast(d_in // 2).inf_mul(4)

    return Transformation(
        input_domain=SizedDomain(VectorDomain(AllDomain(TIA)), size),
        output_domain=VectorDomain(AllDomain(TOA)),
        function=function,
        input_metric=SymmetricDistance(),
        output_metric=InfDifferenceDistance(),
        stability_map=stability_map,
    )
```

[1] See new changes with `git diff f5bb719..4e842c74 rust/src/transformations/quantile_score_candidates/mod.rs`

1.1 Appropriate Output Domain

The raw type and domain are equivalent, save for potential nullity in the atomic type. The scalar scorer structurally cannot emit null, because the input argument is non-null.

1.2 Domain-metric compatibility

On the input side, SymmetricDistance is well-defined on sized VectorDomains. On the output side, InfDifferenceDistance is well-defined on VectorDomains.

1.3 Stability Guarantee

Lemma 1.1. If $d_{CO}(X, X') \leq 1$, then $d_\infty(\texttt{function}(X), \texttt{function}(X')) \leq 2$.

Proof. Assume $d_{CO}(X, X') \leq 1$. For convenience, let $s = \texttt{function}(X)$. Consider each of the four cases of changing a row in X.

$$d_\infty(s_i, s_i') = \max_i |s_i - s_i'| \qquad \text{by definition of } d_\infty$$

Case 1. Assume X' is equal to X, but with some $X_j < C_i$ replaced with $X_j' > C_i$.

$$\begin{aligned}
&= 2 \max_{X \sim X'} \max_i ||(1-\alpha)\#(X < C_i) - \alpha\#(X > C_i)| \\
&\qquad - (1-\alpha)(\#(X < C_i) - 1) - \alpha(\#(X > C_i) + 1)|| && \text{by definition of } \texttt{function} \\
&\leq 2 \max_{X \sim X'} \max_i ||(1-\alpha)\#(X < C_i) - \alpha\#(X > C_i)| \\
&\qquad - (|(1-\alpha)\#(X < C_i) - \alpha\#(X > C_i)| + |1|)| && \text{by triangle inequality} \\
&= 2 \max_{X \sim X'} \max_i |1| && \text{scores cancel} \\
&= 2
\end{aligned}$$

Case 2. Assume X' is equal to X, but with some $X_j > C_i$ replaced with $X_j' < C_i$.

$$= 2 \qquad \text{by symmetry, follows from Case 1.}$$

Case 3. Assume X' is equal to X, but with some $X_j \neq C_i$ replaced with C_i.

$$\leq 2max(\alpha, 1-\alpha) \qquad \text{equivalent to one removal (see } \texttt{make_quantile_score_c}$$

Case 4. Assume X' is equal to X, but with some $X_j = C_i$ replaced with $X_j' \neq C_i$.

$$\leq 2max(\alpha, 1-\alpha) \qquad \text{equivalent to one addition (see } \texttt{make_quantile_score_c}$$

Take the union bound over all cases.

$$d_\infty(s_i, s_i') \leq max(2, 2max(\alpha, 1-\alpha)) = 2 \qquad \text{since } max(\alpha, 1-\alpha) \leq 1$$

□

Take any two elements X, X' in the `input_domain` and any pair (`d_in`, `d_out`), where `d_in` has the associated type for `input_metric` and `d_out` has the associated type for `output_metric`. Assume X, X' are `d_in`-close under `input_metric` and that `stability_map`(`d_in`) $\leq$ `d_out`.

$$
\begin{aligned}
\texttt{d_out} &= \max_{X \sim X'} d_{IDD}(s, s') \\
&= \max_{X \sim X'} \max_{ij} |(s_i - s'_i) - (s_j - s'_j)| && \text{by definition of } \texttt{InfDifferenceDistance} \\
&\leq 2 \max_{X \sim X'} \max_{i} |s_i - s'_i| && s \text{ is not monotonic; take looser bound} \\
&\leq 2 \sum_{j}^{\texttt{d_in}//2} \max_{Z_j \sim Z_{j+1}} \max_{i} |s_{i,j} - s_{i,j+1}| && \text{by path property where } d_{CO}(Z_i, Z_{i+1}) = 1, X = Z_0 \text{ and } Z_{\texttt{d_in}} = X' \\
&\leq 2 \sum_{j}^{\texttt{d_in}//2} 2 && \text{by 1.1} \\
&\leq 4(\texttt{d_in}//2)
\end{aligned}
$$

It is shown that `function`(X), `function`(X') are `d_out`-close under `output_metric`.