fn sample_bernoulli_exp

Michael Shoemate (@Shoeboxam)

October 18, 2026

Proves soundness of sample_bernoulli_exp in mod.rs at commit 0be3ab3e6 (outdated[1]).
fn sample_bernoulli_exp returns a sample from the $Bernoulli(exp(-x))$ distribution for some rational, non-negative x. This proof is an adaptation of subsection 5.1 of [CKS20].

1 Hoare Triple

Precondition

Compiler-verified

- Argument x is of type RBig, a bignum rational

Caller-verified

- $x > 0$.

Pseudocode

```
def sample_bernoulli_exp(x) -> bool:
    while x > 1:
        if sample_bernoulli_exp1(1):  #
            x -= 1
        else:
            return False
    return sample_bernoulli_exp1(x)  #
```

Postcondition

Theorem 1.1. For any setting of the input parameters x such that the given preconditions hold, sample_bernoulli_exp either returns Err(e) due to a lack of system entropy, or Ok(out), where out is distributed as $Bernoulli(exp(-x))$.

Assume the preconditions are met.

Lemma 1.2. sample_bernoulli_exp only returns Err(e) when there is a lack of system entropy.

Proof. In all invocations of sample_bernoulli_exp1, the argument passed satisfies its definition preconditions, by the preconditions on x and function logic. Thus, by its definition, sample_bernoulli_exp1 only returns an error when there is a lack of system entropy. The only source of errors in sample_bernoulli_exp is from the invocation of sample_bernoulli_exp1. Therefore sample_bernoulli_exp only returns Err(e) when there is a lack of system entropy. □

Lemma 1.3. out is distributed as $Bernoulli(exp(-x))$.

[1] See new changes with git diff 0be3ab3e6..5c6c303 rust/src/traits/samplers/cks20/mod.rs

Proof. For $1 \leq i \leq \lfloor x \rfloor$, let b_i denote the ith outcome of `sample_bernoulli_exp1` on line 3. By the definition of `sample_bernoulli_exp1`, under the established conditions and preconditions, each B_i is distributed as $Bernoulli(exp(-1))$. Similarly as before, C is distributed $Bernoulli(exp(-(x - \lfloor x \rfloor)))$.

$$
\begin{aligned}
P[\texttt{out} = \top] &= P[B_1 = B_2 = ... = B_{\lfloor x \rfloor} = C = \top] && \texttt{out} \text{ is only } \top \text{ if } \forall i,\ B_i = \top \text{ and } C = \top \\
&= \prod_{i=1}^{\lfloor x \rfloor} P[B_i = \top] P[C = \top] && \text{all } B_i \text{ and } C \text{ are independent} \\
&= exp(-1)^{\lfloor x \rfloor} exp(\lfloor x \rfloor - x) \\
&= exp(-x)
\end{aligned}
$$

Therefore, `out` is distributed as $Bernoulli(exp(-x))$. □

Proof. 1.1 holds by 1.2 and 1.3. □

References

[CKS20] Clément L. Canonne, Gautam Kamath, and Thomas Steinke. The discrete gaussian for differential privacy. *CoRR*, abs/2004.00010, 2020.