# fn sample_logarithmic_exp

Michael Shoemate

April 22, 2026

This proof resides in **"contrib"** because it has not completed the vetting process.

Proves soundness of `fn sample_logarithmic_exp` in `mod.rs`. This proof uses the exact geometric-exp sampler and exact rational Bernoulli sampler to realize the logarithmic distribution.

## 1 Hoare Triple

### Precondition

$\mathtt{x} \in \mathbb{Q} \wedge \mathtt{x} > 0$

### Pseudocode

```
1
2 def sample_logarithmic_exp(x):
3     while True:
4         k = sample_geometric_exp_fast(x)
5         if k == 0:
6             continue
7
8         if sample_bernoulli_rational(1 / k):
9             return k
```

### Postcondition

**Theorem 1.1.** For any setting of the input parameter `x` such that the given preconditions hold, `sample_logarithmic_exp` either returns `Err(e)` due to a lack of system entropy, or `Ok(out)`, where `out` is distributed on $\mathbb{N}_{>0}$ with

$$\forall k \in \mathbb{N}_{>0}, \qquad \Pr[\mathtt{out} = k] = \frac{e^{-\mathtt{x}k}}{k \log(1/(1 - e^{-\mathtt{x}}))}.$$

## 2 Proof

Assume the preconditions are met and write $\lambda = e^{-x} \in (0, 1)$.

**Lemma 2.1.** `sample_logarithmic_exp` only returns `Err(e)` when there is a lack of system entropy.

*Proof.* Since $\mathtt{x} > 0$, the precondition on `sample_geometric_exp_fast` is met. By its definition, it can only return an error due to a lack of system entropy.

If `k` on line **??** is nonzero, then `k` is a positive integer, so the preconditions on `sample_bernoulli_rational` are met when called with probability $1/\mathtt{k}$ on line **??**. By its definition, it can only return an error due to a lack of system entropy.

The only source of errors in `sample_logarithmic_exp` is from the invocation of these functions, therefore `sample_logarithmic_exp` only returns `Err(e)` when there is a lack of system entropy. □

We now condition on not returning an error and fix any iteration of the loop.

- Let $g$ be a realization of a random variable $G \sim \mathit{Geometric}(1 - e^{-x})$, supported on $\mathbb{N}_0$.
- If $g > 0$, let $b$ be a realization of a random variable $B \sim \mathit{Bernoulli}(1/g)$.

**Lemma 2.2.** For any $k \in \mathbb{N}_{>0}$, the probability that a single iteration of the loop returns $k$ is

$$(1 - e^{-x})\frac{e^{-xk}}{k}.$$

*Proof.* A single iteration returns $k$ exactly when:

1. the geometric draw equals $k$, and
2. the Bernoulli draw with success probability $1/k$ succeeds.

Therefore,

$$\begin{aligned}\Pr[\text{single iteration returns } k] &= \Pr[G = k]\Pr[B = \top \mid G = k] \\ &= (1 - e^{-x})e^{-xk} \cdot \frac{1}{k}.\end{aligned}$$

□

**Lemma 2.3.** The probability that a single iteration of the loop returns some value in $\mathbb{N}_{>0}$ is

$$(1 - e^{-x}) \log\left(\frac{1}{1 - e^{-x}}\right).$$

*Proof.* By Lemma 2.2,

$$\begin{aligned}\Pr[\text{single iteration returns a value}] &= \sum_{k=1}^{\infty}(1 - e^{-x})\frac{e^{-xk}}{k} \\ &= (1 - e^{-x})\sum_{k=1}^{\infty}\frac{(e^{-x})^k}{k}.\end{aligned}$$

Since $e^{-x} \in (0, 1)$, the power-series identity

$$\sum_{k=1}^{\infty}\frac{z^k}{k} = \log\left(\frac{1}{1 - z}\right) \quad \text{for } z \in (0, 1)$$

applies with $z = e^{-x}$, giving

$$\Pr[\text{single iteration returns a value}] = (1 - e^{-x}) \log\left(\frac{1}{1 - e^{-x}}\right).$$

□

**Lemma 2.4.** If the outcome of `sample_logarithmic_exp` is `Ok(out)`, then `out` is distributed on $\mathbb{N}_{>0}$ with

$$\forall k \in \mathbb{N}_{>0}, \qquad \Pr[\texttt{out} = k] = \frac{e^{-xk}}{k \log(1/(1 - e^{-x}))}.$$

*Proof.* By Lemma 2.2, the probability that a fixed loop iteration returns $k$ is

$$r_k := (1 - e^{-x}) \frac{e^{-xk}}{k}.$$

By Lemma 2.3, the probability that a fixed loop iteration returns any value is

$$a := (1 - e^{-x}) \log\left(\frac{1}{1 - e^{-x}}\right).$$

Since loop iterations are independent, the probability that the algorithm returns $k$ on the $t$-th iteration is

$$(1 - a)^{t-1} r_k.$$

Summing over all $t \geq 1$,

$$\begin{aligned}
\Pr[\texttt{out} = k] &= \sum_{t=1}^{\infty} (1 - a)^{t-1} r_k \\
&= \frac{r_k}{a} \\
&= \frac{(1 - e^{-x}) e^{-xk}/k}{(1 - e^{-x}) \log(1/(1 - e^{-x}))} \\
&= \frac{e^{-xk}}{k \log(1/(1 - e^{-x}))}.
\end{aligned}$$

Thus `out` has the claimed distribution. □

*Proof of Theorem 1.1.* Holds by Lemma 2.1 and Lemma 2.4. □