





pyTMD: Python-based tidal prediction software

Tyler C. Sutterley¹, Susan L. Howard^{2*}, Laurie Padman^{2*}, and
Matthew R. Siegfried^{3*}

¹ Polar Science Center, Applied Physics Laboratory, University of Washington, United States ² Earth &
Space Research, United States ³ Department of Geophysics, Colorado School of Mines, United States ¶
Corresponding author * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Taher Chagini](#) 

Reviewers:

- [@castelao](#)
- [@rcaneill](#)

Submitted: 05 June 2025

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#)).

Summary

pyTMD is an open-source tidal prediction software that aims to simplify the calculation of ocean and Earth tides. It is not an ocean or load tide model, but a tool for using tidal constants provided by tide models to predict the height deflections or currents at particular locations and times. It is designed to be able to handle a wide range of model formats and can incorporate different physics schemes. This flexibility allows pyTMD to be tailored to fit specific use cases, while still allowing ease of use.

Statement of need

There are several ocean tide prediction software options available. The OSU Tidal Inversion Software (OTIS) and OSU Tidal Prediction Software (OTPS) are Fortran programs developed by Lana Erofeeva and Gary Egbert for the [TPXO](#) family of models. The Tidal Model Driver (TMD) is a MATLAB Toolbox developed by Laurie Padman, Lana Erofeeva and Susan Howard for the same family of models. An updated version of the MATLAB Toolbox (TMD3) was developed by Chad Greene. [pyFES](#) is a Python library produced by LEGOS, NOVELTIS and CLS Ocean and Climate Division, and funded by CNES ([Lyard et al., in review](#)) for the Finite Element Solution (FES) family of models. The NASA GSFC PREDict Tidal Heights (PERTH3) software is a Fortran program developed by Richard Ray and Remko Scharroo for the Goddard Ocean Tide (GOT) family of models. An updated and more versatile version of the NASA GSFC Fortran software ([PERTH5](#)) was developed by Richard Ray. PERTH5 is a more generalized program, and can read from multiple different tide model formats. These software options are typically created by or for the model providers, and, with the exception of PERTH5, singly support their specific model formats.

pyTMD is a generalized tide program that allows users to calculate both tide deflections and currents. As a tide model driver, pyTMD can read from a broad suite of models, and use different physics schemes in the internal calculations. Over 50 different models are presently supported within pyTMD, and additional models can be defined with a JSON file.

pyTMD was designed to be used by beginners and scientific researchers alike. The online documentation contains background information for both tidal modeling and prediction. The software has been used in a number of scientific publications for modeling regional tides ([Freer et al., 2023](#); [Millan et al., 2023](#); [T. C. Sutterley et al., 2019](#)), modeling global tides ([Gregg et al., 2024](#); [Paprotny et al., 2024](#)), and creating several Earth observation datasets ([ENVEO et al., 2021](#); [Smith et al., 2024a, 2024b](#)). It has also been leveraged within larger earth-observation software packages ([Bishop-Taylor et al., 2025](#); [Fitzpatrick et al., 2024](#)).

Functionality

Ocean and Load Tides

With the harmonic method, tides are decomposed into harmonic constants, or constituents, associated with the relative positions of the sun, moon and Earth (Padman et al., 2018). These constituents are typically classified into different “species” based on their approximate period: short-period, semi-diurnal, diurnal, and long-period. pyTMD.io contains routines for reading major constituent values (amplitude and phase) from commonly available tide models, which typically fall within a few general formats: OTIS-binary (Egbert & Erofeeva, 2002; Padman et al., 2008), OTIS-compact, OTIS-netcdf, TMD3-netcdf (Greene et al., 2024), GOT-ascii (Ray, 1999), GOT-netcdf, FES-ascii (Le Provost et al., 1994) and FES-netcdf (Lyard et al., in review). Information for each of the supported tide models is stored within a JSON database. For tidal predictions, pyTMD.io interpolates the amplitude and phase lag of tide model constituents to sets of spatial coordinates.

pyTMD uses the astronomical argument formalism outlined in Doodson & Lamb (1921) to compute the temporal elements. For a set of temporal values, pyTMD 1) calculates the astronomical angles (S , H , P , N , P_s) (Meeus, 1991; Simon et al., 1994), 2) combines these angles with the “Doodson numbers” in a Fourier series to compute the equilibrium tide phase (G) of each constituent, and 3) computes the 18.6-year nodal amplitude and phase corrections (f and u) of each constituent (Dietrich, 1980; Doodson & Lamb, 1921; Pugh & Woodworth, 2014). The spatial and temporal components are then combined, and the output tidal time series is calculated through a summation over all constituents (Egbert & Erofeeva, 2002). To include more of the tidal spectrum, the contributions of “minor” constituents can additionally be estimated using inference methods (Ray, 2017; Schureman, 1958).

Long-period ocean tides can independently be predicted assuming an “equilibrium response” (Cartwright & Edden, 1973; Cartwright & Tayler, 1971; Doodson & Lamb, 1921). Here, the oceanic surface is estimated to respond instantaneously to the tide-producing forces of the moon and sun, and is not influenced by inertia, currents or the irregular distribution of land (Proudman, 1960; Ray & Erofeeva, 2014; Schureman, 1958).

Pole Tides

The Earth’s rotation axis is inclined at an angle of 23.5 degrees to the celestial pole, of which it rotates about every 26,000 years (Kantha & Clayson, 2000). Superimposed on this long-term precession, the rotation axis shifts with respect to its mean pole position due to nutation, Chandler wobble, annual variations, and other processes (Desai, 2002; Wahr, 1985). Load and ocean pole tides are driven by variations in the Earth’s rotation axis with respect to its mean position, along with corresponding elastic responses and secondary effects (Desai, 2002; Desai et al., 2015; Wahr, 1985). pyTMD.predict estimates pole tide variations following IERS Conventions by differencing the daily IERS polar motion “finals” from the reference “secular” pole positions (Petit & Luzum, 2010).

Solid Earth Tides

The tidal deformation of the solid Earth can be modeled in pyTMD using one of the following two methods: 1) the ephemerides formalism from Wahr (1981) and Mathews et al. (1997) as described in Petit & Luzum (2010), and 2) the tide catalog formalism outlined in Cartwright & Tayler (1971). For the ephemerides method, pyTMD.astro has options for calculating approximate ephemerides following Meeus (1991) and Montenbruck (1989) or using high-resolution JPL ephemerides from Park et al. (2021) with the jplephem package (Rhodes, 2011). The calculation for solid Earth tides includes multiple Love and Shida number corrections including the frequency-dependent and dissipative mantle anelasticity corrections from Mathews et al. (1997).

Time

pyTMD uses the timescale library (T. Sutterley et al., 2025) to manage temporal conversions, calculate “dynamical” time scales, and estimate Earth Orientation Parameters (EOPs).

Acknowledgements

pyTMD was first supported through an appointment to the NASA Postdoctoral Program (NPP) at NASA Goddard Space Flight Center (GSFC), and currently supported under the NASA Cryospheric Sciences Program (Grant Number 80NSSC22K0379). The software was initially developed with the goal of supporting science applications for airborne and satellite altimetry in preparation for the launch of the NASA ICESat-2 mission. It was designed for scientific and technical purposes, and not for coastal navigation or applications risking life or property.

We wish to acknowledge the invaluable comments, contributions, and support from Karen Alley (University of Manitoba), Robbi Bishop-Taylor (Geoscience Australia), Kelly Brunt (NSF) and Richard Ray (NASA GSFC) towards the development of pyTMD. We additionally wish to acknowledge the comments, issues and discussions of all contributors to the pyTMD GitHub repository.

References

- Bishop-Taylor, R., Phillips, C., Sagar, S., Newey, V., & Sutterley, T. (2025). eo-tides: Tide modelling tools for large-scale satellite Earth observation analysis. *Journal of Open Source Software*, 10(109), 7786. <https://doi.org/10.21105/joss.07786>
- Cartwright, D. E., & Edden, A. C. (1973). Corrected Tables of Tidal Harmonics. *Geophysical Journal International*, 33(3), 253–264. <https://doi.org/10.1111/j.1365-246X.1973.tb03420.x>
- Cartwright, D. E., & Tayler, R. J. (1971). New Computations of the Tide-generating Potential. *Geophysical Journal of the Royal Astronomical Society*, 23(1), 45–73. <https://doi.org/10.1111/j.1365-246X.1971.tb01803.x>
- Desai, S. (2002). Observing the pole tide with satellite altimetry. *Journal of Geophysical Research: Oceans*, 107(C11). <https://doi.org/10.1029/2001JC001224>
- Desai, S., Wahr, J., & Beckley, B. (2015). Revisiting the pole tide for and from satellite altimetry. *Journal of Geodesy*, 89(12), 1233–1243. <https://doi.org/10.1007/s00190-015-0848-7>
- Dietrich, G. (1980). *General oceanography: an introduction* (2nd ed.). John Wiley & Sons, Inc. ISBN: 0471021024
- Doodson, A. T., & Lamb, H. (1921). The harmonic development of the tide-generating potential. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 100(704), 305–329. <https://doi.org/10.1098/rspa.1921.0088>
- Egbert, G. D., & Erofeeva, S. Y. (2002). Efficient Inverse Modeling of Barotropic Ocean Tides. *Journal of Atmospheric and Oceanic Technology*, 19(2), 183–204. [https://doi.org/10.1175/1520-0426\(2002\)019%3C0183:EIMOBO%3E2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019%3C0183:EIMOBO%3E2.0.CO;2)
- ENVEO, Wuite, J., Hetzenecker, M., Nagler, T., & Scheiblaue, S. (2021). *ESA Antarctic Ice Sheet Climate Change Initiative (Antarctic_Ice_Sheet_cci): Antarctic Ice Sheet monthly velocity from 2017 to 2020, derived from Sentinel-1*. NERC EDS Centre for Environmental Data Analysis. <https://doi.org/10.5285/00fe090efc58446e8980992a617f632f>
- Fitzpatrick, S., Buscombe, D., Warrick, J. A., Lundine, M. A., & Vos, K. (2024). CoastSeg: an accessible and extendable hub for satellite-derived-shoreline (SDS) detection and mapping.

- 132 *Journal of Open Source Software*, 9(99), 6683. <https://doi.org/10.21105/joss.06683>
- 133 Freer, B. I. D., Marsh, O. J., Hogg, A. E., Fricker, H. A., & Padman, L. (2023). Modes
134 of Antarctic tidal grounding line migration revealed by Ice, Cloud, and land Elevation
135 Satellite-2 (ICESat-2) laser altimetry. *The Cryosphere*, 17(9), 4079–4101. <https://doi.org/10.5194/tc-17-4079-2023>
- 136
- 137 Greene, C. A., Erofeeva, S., Padman, L., Howard, S. L., Sutterley, T., & Egbert, G. (2024).
138 Tide Model Driver for MATLAB. *Journal of Open Source Software*, 9(95), 6018. <https://doi.org/10.21105/joss.06018>
- 139
- 140 Gregg, D. E., Penna, N. T., Jones, C., & Maqueda, M. A. (2024). Accuracy assessment of
141 recent global ocean tide models in coastal waters of the European North West Shelf. *Ocean*
142 *Modelling*, 192, 102448. <https://doi.org/10.1016/j.ocemod.2024.102448>
- 143 Kantha, L. H., & Clayson, C. A. (2000). *Numerical models of oceans and oceanic processes*
144 (Vol. 66). Academic Press. [https://doi.org/10.1016/s0074-6142\(00\)x8001-1](https://doi.org/10.1016/s0074-6142(00)x8001-1)
- 145 Le Provost, C., Genco, M. L., Lyard, F., Vincent, P., & Cancail, P. (1994). Spectroscopy of
146 the world ocean tides from a finite element hydrodynamic model. *Journal of Geophysical*
147 *Research: Oceans*, 99(C12), 24777–24797. <https://doi.org/10.1029/94JC01381>
- 148 Lyard, F., Carrere, L., Fouchet, E., Cancet, M., Greenberg, D., Dibarbour, G., & Picot, N.
149 (in review). FES2022: a step towards a SWOT-compliant tidal correction. *Journal of*
150 *Geophysical Research: Oceans*.
- 151 Mathews, P. M., Dehant, V., & Gipson, J. M. (1997). Tidal station displacements. *Journal*
152 *of Geophysical Research: Solid Earth*, 102(B9), 20469–20477. <https://doi.org/10.1029/97JB01515>
- 153
- 154 Meeus, J. H. (1991). *Astronomical Algorithms*. Willmann-Bell, Inc. ISBN: 0943396352
- 155 Millan, R., Jager, E., Mouginot, J., Wood, M. H., Larsen, S. H., Mathiot, P., Jourdain, N. C.,
156 & Bjørk, A. (2023). Rapid disintegration and weakening of ice shelves in North Greenland.
157 *Nature Communications*, 14(1). <https://doi.org/10.1038/s41467-023-42198-2>
- 158 Montenbruck, O. (1989). *Practical Ephemeris Calculations*. Springer-Verlag. <https://ui.adsabs.harvard.edu/abs/1989pec..book.....M>
- 159
- 160 Padman, L., Erofeeva, S. Y., & Fricker, H. A. (2008). Improving Antarctic tide models by
161 assimilation of ICESat laser altimetry over ice shelves. *Geophysical Research Letters*, 35(22).
162 <https://doi.org/10.1029/2008GL035592>
- 163 Padman, L., Siegfried, M. R., & Fricker, H. A. (2018). Ocean Tide Influences on the
164 Antarctic and Greenland Ice Sheets. *Reviews of Geophysics*, 56(1), 142–184. <https://doi.org/10.1002/2016RG000546>
- 165
- 166 Paprotny, D., Rhein, B., Voutsoukas, M. I., Terefenko, P., Dottori, F., Treu, S., Śledziowski,
167 J., Feyen, L., & Kreibich, H. (2024). Merging modelled and reported flood impacts in
168 Europe in a combined flood event catalogue for 1950–2020. *Hydrology and Earth System*
169 *Sciences*, 28(17), 3983–4010. <https://doi.org/10.5194/hess-28-3983-2024>
- 170 Park, R. S., Folkner, W. M., Williams, J. G., & Boggs, D. H. (2021). The JPL Planetary
171 and Lunar Ephemerides DE440 and DE441. *The Astronomical Journal*, 161(3), 105.
172 <https://doi.org/10.3847/1538-3881/abd414>
- 173 Petit, G., & Luzum, B. (2010). *IERS Conventions (2010)* (No. 36). Bureau International des
174 Poids et Mesures (BIPM), US Naval Observatory (USNO); International Earth Rotation;
175 Reference Systems Service (IERS). http://www.iers.org/nn_11216/IERS/EN/Publications/TechnicalNotes/tn36.html
- 176
- 177 Proudman, J. (1960). The Condition that a Long-Period Tide shall follow the Equilibrium-Law.
178 *Geophysical Journal International*, 3(2), 244–249. <https://doi.org/10.1111/j.1365-246X.1960.tb00000.x>

- 179 [1960.tb00392.x](#)
- 180 Pugh, D., & Woodworth, P. (2014). *Sea-Level Science: Understanding Tides, Surges,*
181 *Tsunamis and Mean Sea-Level Changes*. Cambridge University Press. [https://doi.org/10.](https://doi.org/10.1017/CBO9781139235778)
182 [1017/CBO9781139235778](https://doi.org/10.1017/CBO9781139235778)
- 183 Ray, R. D. (1999). *A global ocean tide model from Topex/Poseidon altimetry: GOT99.2*
184 *(TM-1999-209478)*. NASA Goddard Space Flight Center; NASA Goddard Space Flight
185 Center. <https://ntrs.nasa.gov/citations/19990089548>
- 186 Ray, R. D. (2017). On Tidal Inference in the Diurnal Band. *Journal of Atmospheric and*
187 *Oceanic Technology*, 34(2), 437–446. <https://doi.org/10.1175/jtech-d-16-0142.1>
- 188 Ray, R. D., & Erofeeva, S. Y. (2014). Long-period tidal variations in the length of day.
189 *Journal of Geophysical Research: Solid Earth*, 119(2), 1498–1509. [https://doi.org/10.](https://doi.org/10.1002/2013JB010830)
190 [1002/2013JB010830](https://doi.org/10.1002/2013JB010830)
- 191 Rhodes, B. C. (2011). *PyEphem: Astronomical Ephemeris for Python [Computer software]*.
192 Astrophysics Source Code Library. <https://ui.adsabs.harvard.edu/abs/2011ascl.soft12014R>
- 193 Schureman, P. (1958). *Manual of Harmonic Analysis and Prediction of Tides* (Special
194 Edition No. 98). US Coast; Geodetic Survey; United States Government Printing Office.
195 <https://tidesandcurrents.noaa.gov/publications/SpecialPubNo98.pdf>
- 196 Simon, J. L., Bretagnon, P., Chapront, J., Chapront-Touzé, M., Francou, G., & Laskar, J.
197 (1994). Numerical expressions for precession formulae and mean elements for the Moon
198 and the planets. *Astronomy and Astrophysics*, 282, 663–683. [https://ui.adsabs.harvard.](https://ui.adsabs.harvard.edu/abs/1994A&A...282..663S)
199 [edu/abs/1994A&A...282..663S](https://ui.adsabs.harvard.edu/abs/1994A&A...282..663S)
- 200 Smith, B., Sutterley, T. C., Dickinson, S., Jelley, B. P., Felikson, D., Neumann, T. A.,
201 Fricker, H. A., Gardner, A., Padman, L., Markus, T., Kurtz, N., Bhardwaj, S., Hancock,
202 D., & Lee, J. (2024a). *ATLAS/ICESat-2 L3B Gridded Antarctic and Arctic Land Ice*
203 *Height*. NASA National Snow; Ice Data Center Distributed Active Archive Center. [https:](https://doi.org/10.5067/ATLAS/ATL14.004)
204 [//doi.org/10.5067/ATLAS/ATL14.004](https://doi.org/10.5067/ATLAS/ATL14.004)
- 205 Smith, B., Sutterley, T. C., Dickinson, S., Jelley, B. P., Felikson, D., Neumann, T. A.,
206 Fricker, H. A., Gardner, A., Padman, L., Markus, T., Kurtz, N., Bhardwaj, S., Hancock,
207 D., & Lee, J. (2024b). *ATLAS/ICESat-2 L3B Gridded Antarctic and Arctic Land Ice*
208 *Height Change*. NASA National Snow; Ice Data Center Distributed Active Archive Center.
209 <https://doi.org/10.5067/ATLAS/ATL15.004>
- 210 Sutterley, T. C., Markus, T., Neumann, T. A., Broeke, M. van den, Wessem, J. van, &
211 Ligtenberg, S. R. (2019). Antarctic ice shelf thickness change from multimission lidar
212 mapping. *The Cryosphere*, 13(7), 1801–1817. <https://doi.org/10.5194/tc-13-1801-2019>
- 213 Sutterley, T., Alley, K., Brunt, K., Howard, S., Padman, L., & Siegfried, M. (2025). *timescale*
214 *[Computer software]* (Version 0.0.8). Zenodo. <https://doi.org/10.5281/zenodo.8284224>
- 215 Wahr, J. M. (1981). Body tides on an elliptical, rotating, elastic and oceanless Earth.
216 *Geophysical Journal of the Royal Astronomical Society*, 64(3), 677–703. [https://doi.org/](https://doi.org/10.1111/j.1365-246X.1981.tb02690.x)
217 [10.1111/j.1365-246X.1981.tb02690.x](https://doi.org/10.1111/j.1365-246X.1981.tb02690.x)
- 218 Wahr, J. M. (1985). Deformation induced by polar motion. *Journal of Geophysical Research:*
219 *Solid Earth*, 90(B11), 9363–9368. <https://doi.org/10.1029/JB090iB11p09363>