

pyMannKendall: a python package for non parametric Mann Kendall family of trend tests.

Md. Manjurul Hussain¹ and Ishtiak Mahmud²

¹ Institute of Water and Flood Management, Bangladesh University of Engineering and Technology, Dhaka, Bangladesh ² Shahjalal University of Science and Technology, Sylhet, Bangladesh

DOI: [10.21105/joss.01556](https://doi.org/10.21105/joss.01556)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Submitted: 30 June 2019

Published: 25 July 2019

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

Summary

Trend analysis is one of the most important measurements in studying time series data. Both parametric and non-parametric tests are commonly used in trend analysis. Parametric tests require data to be independent and normally distributed. On the other hand, non-parametric trend tests require only that the data be independent and can tolerate outliers in the data (Hamed & Rao, 1998). However, parametric tests are more powerful than nonparametric ones.

The Mann–Kendall trend test (Kendall, 1975; Mann, 1945) is a widely used non-parametric tests to detect significant trends in time series. However, the original Mann-Kendall test didn't consider serial correlation or seasonality effects (Bari, Rahman, Hoque, & Hussain, 2016; Hirsch, Slack, & Smith, 1982). But, in many real situations, the observed data are autocorrelated and this autocorrelation will result in misinterpretation of trend tests results (Cox & Stuart, 1955; Hamed & Rao, 1998). Contrariwise, water quality, hydrologic, as well as climatic and other natural time series also have seasonality. To overcome those limitations of original Mann-Kendall test, various modified Mann-Kendall test have been developed.

Again, Python is one of the widely used tools for data analysis. A large number of data analysis and research tools are also developed using Python. But, till now, there is no Mann-Kendall trend relation Python package available. `pyMannKendall` package fills this gap.

`pyMannKendall` is written in pure Python and uses a vectorization approach to increase its performance. Currently, this package has 11 Mann-Kendall Tests and 2 Sen's slope estimator functions. Brief description of the functions are below:

1. **Original Mann-Kendall test (`original_test`):** Original Mann-Kendall test (Kendall, 1975; Mann, 1945) is a nonparametric test, which does not consider serial correlation or seasonal effects.
2. **Hamed and Rao Modified MK Test (`hamed_rao_modification_test`):** This modified MK test was proposed by Hamed & Rao (1998) to address serial autocorrelation issues. They suggested a variance correction approach to improve trend analysis. Users can consider first n significant lag by insert lag number in this function. By default, it considered all significant lags.
3. **Yue and Wang Modified MK Test (`yue_wang_modification_test`):** This is also a variance correction method for considered serial autocorrelation proposed by Yue & Wang (2004). Users can also set their desired significant number of lags for the calculation.
4. **Modified MK test using Pre-Whitening method (`pre_whitening_modification_test`):** This test was suggested by Yue & Wang (2002) to use Pre-Whitening the time series before the application of trend test.

5. **Modified MK test using Trend free Pre-Whitening method (*trend_free_pre_whitening_m*)**: This test was also proposed by Yue, Pilon, Phinney, & Cavadias (2002) to remove trend components and then Pre-Whitening the time series before application of trend test.
6. **Multivariate MK Test (*multivariate_test*)**: This is an MK test for multiple parameters proposed by Hirsch et al. (1982). They used this method for seasonal MK tests, where they considered every month as a parameter.
7. **Seasonal MK Test (*seasonal_test*)**: For seasonal time series data, Hirsch et al. (1982) proposed this test to calculate the seasonal trend.
8. **Regional MK Test (*regional_test*)**: Based on the proposed seasonal MK test of Hirsch et al. (1982), Helsel & Frans (2006) suggest a regional MK test to calculate the overall trend on a regional scale.
9. **Correlated Multivariate MK Test (*correlated_multivariate_test*)**: This multivariate MK test was proposed by Hipel & McLeod (1994) for where the parameters are correlated.
10. **Correlated Seasonal MK Test (*correlated_seasonal_test*)**: This method was proposed by Hipel & McLeod (1994), for when time series significantly correlate with the preceding one or more months/seasons.
11. **Partial MK Test (*partial_test*)**: In a real event, many factors affect the main studied response parameter, which can bias the trend results. To overcome this problem, Libiseller & Grimvall (2002) proposed this partial mk test. It required two parameters as input, where one is the response parameter and other is an independent parameter.
12. **Theil-sen's Slope Estimator (*sens_slope*)**: This method was proposed by Theil (1950) and Sen (1968) to estimate the magnitude of the monotonic trend.
13. **Seasonal sen's Slope Estimator (*seasonal_sens_slope*)**: This method was proposed by Hipel & McLeod (1994) to estimate the magnitude of the monotonic trend, when data has seasonal effects.

pyMannKendall is a non-parametric Mann-Kendall trend analysis package implemented in pure Python, which brings together almost all types of Mann-Kendall tests, which might help researchers to check Mann-Kendall trends in Python.

References

- Bari, S. H., Rahman, M. T. U., Hoque, M. A., & Hussain, M. M. (2016). Analysis of seasonal and annual rainfall trends in the northern region of Bangladesh. *Atmospheric Research*, 176, 148–158. doi:[10.1016/j.atmosres.2016.02.008](https://doi.org/10.1016/j.atmosres.2016.02.008)
- Cox, D. R., & Stuart, A. (1955). Some quick sign tests for trend in location and dispersion. *Biometrika*, 42(1/2), 80–95. doi:[10.2307/2333424](https://doi.org/10.2307/2333424)
- Hamed, K. H., & Rao, A. R. (1998). A modified Mann–Kendall trend test for autocorrelated data. *Journal of hydrology*, 204(1-4), 182–196. doi:[10.1016/S0022-1694\(97\)00125-X](https://doi.org/10.1016/S0022-1694(97)00125-X)
- Helsel, D. R., & Frans, L. M. (2006). Regional Kendall test for trend. *Environmental science & technology*, 40(13), 4066–4073. doi:[10.1021/es051650b](https://doi.org/10.1021/es051650b)
- Hipel, K. W., & McLeod, A. I. (1994). *Time series modelling of water resources and environmental systems* (Vol. 45). Elsevier.

- Hirsch, R. M., Slack, J. R., & Smith, R. A. (1982). Techniques of trend analysis for monthly water quality data. *Water resources research*, 18(1), 107–121. doi:[10.1029/WR018i001p00107](https://doi.org/10.1029/WR018i001p00107)
- Kendall, M. (1975). Rank correlation measures. *Charles Griffin, London*, 202, 15.
- Libiseller, C., & Grimvall, A. (2002). Performance of partial Mann–Kendall tests for trend detection in the presence of covariates. *Environmetrics: The official journal of the International Environmetrics Society*, 13(1), 71–84. doi:[10.1002/env.507](https://doi.org/10.1002/env.507)
- Mann, H. B. (1945). Nonparametric tests against trend. *Econometrica: Journal of the Econometric Society*, 245–259. doi:[10.2307/1907187](https://doi.org/10.2307/1907187)
- Sen, P. K. (1968). Estimates of the regression coefficient based on Kendall's tau. *Journal of the American statistical association*, 63(324), 1379–1389. doi:[10.1080/01621459.1968.10480934](https://doi.org/10.1080/01621459.1968.10480934)
- Theil, H. (1950). A rank-invariant method of linear and polynomial regression analysis (parts 1-3). In *Ned. Akad. Wetensch. Proc. Ser. A* (Vol. 53, pp. 1397–1412).
- Yue, S., & Wang, C. (2004). The Mann–Kendall test modified by effective sample size to detect trend in serially correlated hydrological series. *Water resources management*, 18(3), 201–218. doi:[10.1023/B:WARM.0000043140.61082.60](https://doi.org/10.1023/B:WARM.0000043140.61082.60)
- Yue, S., & Wang, C. Y. (2002). Applicability of prewhitening to eliminate the influence of serial correlation on the Mann–Kendall test. *Water resources research*, 38(6), 4–1. doi:[10.1029/2001WR000861](https://doi.org/10.1029/2001WR000861)
- Yue, S., Pilon, P., Phinney, B., & Cavadias, G. (2002). The influence of autocorrelation on the ability to detect trend in hydrological series. *Hydrological processes*, 16(9), 1807–1829. doi:[10.1002/hyp.1095](https://doi.org/10.1002/hyp.1095)