






















# DVMDOSTEM v0.8.2: a terrestrial ecosystem model designed to represent arctic, boreal and permafrost ecosystem dynamics

Tobey B Carman <sup>1\*</sup>, Hélène Genet <sup>1\*</sup>, Ruth A Rutter <sup>1</sup>, Elchin Jafarov <sup>2</sup>, Eugénie Euskirchen <sup>1</sup>, Ben Maglio <sup>1</sup>, Joy Clein <sup>1</sup>, Valeria Briones <sup>2</sup>, Andrew L Mullen <sup>2</sup>, Heather Greaves <sup>1</sup>, A. D. McGuire <sup>1</sup>, Fengming Yuan <sup>3</sup>, Chu-Chun Chang <sup>2</sup>, Joshua M. Rady <sup>2</sup>, Doğukan Teber<sup>2</sup>, Brendan M Rogers <sup>2</sup>, Trevor Smith<sup>2</sup>, Hannah Mevenkamp <sup>1</sup>, Mark J Lara <sup>6</sup>, Qianlai Zhuang <sup>4</sup>, and Shuhua Yi<sup>5</sup>

<sup>1</sup> Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks 99775, Alaska, USA <sup>2</sup> Woodwell Climate Research Center, Falmouth 02540, Massachusetts, USA <sup>3</sup> Environmental Science Division and Climate Change Science Institute, Oak Ridge National Laboratory, Oak Ridge, TN 37831 <sup>4</sup> Department of Earth, Atmospheric, and Planetary Science, Purdue University, West Lafayette IN 47907 USA <sup>5</sup> School of Geographic Sciences, Nantong University, Nantong Jiangsu, 226019 P.R. China <sup>6</sup> Department(s) of Plant Biology and Geography, University of Illinois Urbana-Champaign, Urbana, IL 61801 ¶ Corresponding author \* These authors contributed equally.

DOI: [10.21105/joss.07667](https://doi.org/10.21105/joss.07667)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Mengqi Zhao](#) 

## Reviewers:

- [@kdoorheim](#)
- [@arbennett](#)

Submitted: 21 August 2024

Published: 10 June 2025

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Summary

The impacts of climate change on natural ecosystems are the result of complex physical and ecological processes operating and interacting at a variety of spatio-temporal scales, that can be represented in process-based ecosystem models.

DVMDOSTEM is an advanced process-based terrestrial ecosystem model (TEM) designed to study ecosystem responses to climate changes and disturbances. It has a particular focus on permafrost regions (i.e. regions characterized by soils that stay partially frozen all year round for at least two consecutive years), encompassing boreal, arctic, and alpine landscapes. The model couples two previous versions of the Terrestrial Ecosystem Model (TEM), ([McGuire et al., 1992](#)): DVMTEM that includes a dynamic vegetation module (DVM) ([E. S. Euskirchen et al., 2009](#)), and DOSTEM that includes a dynamic organic soil module (DOS) ([H. Genet et al., 2013](#); [Yi et al., 2010](#)). DVMDOSTEM simulates processes at yearly and monthly scales, with some physical processes operating at an even finer temporal resolution. Its versatility allows for site-specific to regional simulations, making it valuable for predicting shifts in permafrost, vegetation, and carbon (C) and nitrogen (N) dynamics. While DVMDOSTEM has been described in the methods sections of many manuscripts, this paper is the first stand alone description of DVMDOSTEM, independent of a particular scientific investigation.




Figure 1: Logo for DVMDOSTEM

## Statement of need

Arctic and boreal regions underlain by permafrost store nearly half of the world's soil organic C - approximately 1,440-1,600 Pg (Hugelius et al., 2014; Edward A. G. Schuur et al., 2022). These regions are warming four times faster than the rest of the globe, driving widespread and rapid permafrost thaw (Rantanen et al., 2022; Smith et al., 2022). As permafrost thaws, soil organic C becomes available for decomposition and release as greenhouse gasses (GHGs) to the atmosphere. Climate-driven permafrost thaw and the associated release of GHGs can influence the global climate system, a phenomenon called the permafrost carbon-climate feedback or PCCF (Koven et al., 2011; E. A. G. Schuur et al., 2015). The PCCF has been identified as one of the largest sources of uncertainty in future climate projections and therefore needs to be accurately represented in global earth system models (Schädel et al., 2024). DVMDOSTEM has been developed with special emphasis on physical and biological processes driving permafrost and carbon cycling in high latitude ecosystems. DVMDOSTEM is therefore well suited to assessing and informing our understanding of the PCCF.

## Model Design

DVMDOSTEM is designed to simulate the key biophysical and biogeochemical processes between the soil, the vegetation and the atmosphere. The evolution and refinement of DVMDOSTEM have been shaped by extensive research programs and applications both in permafrost and non-permafrost regions (Briones et al., 2024; Eugénie S. Euskirchen et al., 2022; H. Genet et al., 2013; Hélène Genet et al., 2018; Jafarov et al., 2013; Yi et al., 2009, 2010). The model is spatially explicit and represents ecosystem response to climate and disturbances at seasonal (i.e. monthly) to centennial scales.



**Figure 2:** Overview of DVMDOSTEM soil and vegetation structure. On the left is the soil structure showing the layers and different properties that are tracked (purple bubble: carbon (C), nitrogen (N), temperature (T), volumetric water content (VWC), ice). Each of the layers with properties described above is also categorized as organic (fibric or humic) or mineral. Additionally, the model simulates snow layers and the removal of soil organic layers due to fire. On the right is the vegetation structure showing plant functional types (PFTs) within a community type (CMT) and the associated pools and fluxes of C and N. Each PFT is split into compartments (leaf, stem and root) which track their own C and N content and associated fluxes. The fluxes are represented with red text while the pools are black. In addition, there is competition among the PFTs for light, water, and available N, shown with the purple arrow in the top center.

The snow and soil columns are split into a dynamic number of layers to represent their impact on thermal and hydrological dynamics and the consequences for soil C and N dynamics. Vegetation composition is modeled using community types (CMTs), each of which consists of multiple plant functional types (PFTs - groups of species sharing similar ecological traits). This structure allows the model to represent the effect of competition for light, water and nutrients on vegetation composition (E. S. Euskirchen et al., 2009), as well as the role of nutrient limitation on permafrost ecosystem dynamics, with coupling between C and N cycles (E. S. Euskirchen et al., 2009; McGuire et al., 1992). Finally, the model represents the effects of wildfire in order to evaluate the role of climate-driven fire intensification on ecosystem structure and function (H. Genet et al., 2013; Yi et al., 2010). The structure of DVMDOSTEM is represented visually in Figure 2.

## State of the Field

In the field of ecosystem models, several prominent models such as CLM5 (Lawrence et al., 2019), ELM-FATES (Fisher et al., 2015), LANDIS (Scheller et al., 2007), and iLand (Seidl et al., 2012) have been developed to simulate ecological processes at various scales, resolutions and ecotypes. Like DVMDOSTEM, ELM-FATES, and early versions of CLM, are “offline” land

models that do not include feedback with atmospheric or oceanic models, focusing instead on land surface processes. LANDIS and iLand emphasize forest dynamics with some interaction between grid cells, while DVMDOSTEM does not model interaction between grid cells. While DVMDOSTEM contains detailed representations of vegetation - multiple Plant Functional Types (PFTs) and individual compartments within PFTs - it also has the concept of community types (collections of PFTs and soil properties) and is designed to run at the landscape scale by representing more than a single stand of trees or a single forest type. DVMDOSTEM is unique in its detailed representation of high latitude processes, particularly the dynamic organic soils in regions with frozen ground coupled with dynamic vegetation and high latitude specific parameterizations. This focus allows DVMDOSTEM to simulate the complex interactions between soil, vegetation, and climate in permafrost ecosystems, providing valuable insights into the PCCF. For a more detailed assessment and comparison of high latitude vegetation modeling, see (Heffernan et al., 2024).

## Acknowledgements

The current version of DVMDOSTEM is the result of decades of model developments from the original TEM code developed at the Marine Biological Laboratory (Raich, 1991). Following is the list of recent and current programs and projects that supported model developments over the past ten years: (1) the Integrated Ecosystem Model for Alaska and Northwest Canada project supported by the US Geological Survey and the Arctic, Western Alaska, and Northwest Boreal Landscape Conservation Cooperatives in Alaska, (2) the Bonanza Creek Long Term Ecological Research Program funded by the National Science Foundation and the USDA Forest Service, (3) the Permafrost Pathways project through the TED Audacious Project and the Quadrature Climate Foundation, (4) the Next Generation Ecosystem Experiment program for the Arctic region supported by the Department of Energy.

## References

- Briones, V., Jafarov, E. E., Genet, H., Rogers, B. M., Rutter, R. M., Carman, T. B., Clein, J., Euskirchen, E. S., Schuur, E. A., Watts, J. D., & Natali, S. M. (2024). Exploring the interplay between soil thermal and hydrological changes and their impact on carbon fluxes in permafrost ecosystems. *Environmental Research Letters*, 19, 074003. <https://doi.org/10.1088/1748-9326/ad50ed>
- Euskirchen, E. S., McGuire, A. D., Chapin, F. S., Yi, S., & Thompson, C. C. (2009). Changes in vegetation in northern alaska under scenarios of climate change, 2003–2100: Implications for climate feedbacks. *Ecological Applications*, 19, 1022–1043. <https://doi.org/10.1890/08-0806.1>
- Euskirchen, Eugénie S., Serbin, S. P., Carman, T. B., Fraterrigo, J. M., Genet, H., Iversen, C. M., Salmon, V., & McGuire, A. D. (2022). Assessing dynamic vegetation model parameter uncertainty across alaskan arctic tundra plant communities. *Ecological Applications*, 32. <https://doi.org/10.1002/eap.2499>
- Fisher, R., Muszala, S., Versteinsten, M., Lawrence, P., Xu, C., McDowell, N., Knox, R., Koven, C., Holm, J., Rogers, B., & others. (2015). Taking off the training wheels: The properties of a dynamic vegetation model without climate envelopes. *Geoscientific Model Development Discussions (Online)*, 8(4). <https://doi.org/10.5194/gmdd-8-3293-2015>
- Genet, Hélène, He, Y., Lyu, Z., McGuire, A. D., Zhuang, Q., Clein, J., D'Amore, D., Bennett, A., Breen, A., Biles, F., Euskirchen, E. S., Johnson, K., Kurkowski, T., Schroder, S. (Kushch), Pastick, N., Rupp, T. S., Wylie, B., Zhang, Y., Zhou, X., & Zhu, Z. (2018). The role of driving factors in historical and projected carbon dynamics of upland ecosystems in alaska. *Ecological Applications*, 28, 5–27. <https://doi.org/10.1002/eap.1641>

- Genet, H., McGuire, A. D., Barrett, K., Breen, A., Euskirchen, E. S., Johnstone, J. F., Kasischke, E. S., Melvin, A. M., Bennett, A., Mack, M. C., Rupp, T. S., Schuur, A. E. G., Turetsky, M. R., & Yuan, F. (2013). Modeling the effects of fire severity and climate warming on active layer thickness and soil carbon storage of black spruce forests across the landscape in interior alaska. *Environmental Research Letters*, 8, 045016. <https://doi.org/10.1088/1748-9326/8/4/045016>
- Heffernan, E., Epstein, H., McQuinn, T. D., Rogers, B. M., Virkkala, A.-M., Lutz, D., & Armstrong, A. (2024). Comparing assumptions and applications of dynamic vegetation models used in the arctic-boreal zone of alaska and canada. *Environmental Research Letters*, 19, 093003. <https://doi.org/10.1088/1748-9326/ad6619>
- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L., Schirrmeister, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., & Kuhry, P. (2014). Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps. *Biogeosciences*, 11, 6573–6593. <https://doi.org/10.5194/bg-11-6573-2014>
- Jafarov, E. E., Romanovsky, V. E., Genet, H., McGuire, A. D., & Marchenko, S. S. (2013). The effects of fire on the thermal stability of permafrost in lowland and upland black spruce forests of interior alaska in a changing climate. *Environmental Research Letters*, 8. <https://doi.org/10.1088/1748-9326/8/3/035030>
- Koven, C. D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G., & Tarnocai, C. (2011). Permafrost carbon-climate feedbacks accelerate global warming. *Proceedings of the National Academy of Sciences of the United States of America*, 108. <https://doi.org/10.1073/pnas.1103910108>
- Lawrence, D. M., Fisher, R. A., Koven, C. D., Oleson, K. W., Swenson, S. C., Bonan, G., Collier, N., Ghimire, B., Kampenhout, L. van, Kennedy, D., Kluzek, E., Lawrence, P. J., Li, F., Li, H., Lombardozzi, D., Riley, W. J., Sacks, W. J., Shi, M., Vertenstein, M., ... Zeng, X. (2019). The community land model version 5: Description of new features, benchmarking, and impact of forcing uncertainty. *Journal of Advances in Modeling Earth Systems*, 11, 4245–4287. <https://doi.org/10.1029/2018MS001583>
- McGuire, A. D., Melillo, J. M., Joyce, L. A., Kicklighter, D. W., Grace, A. L., Moore, B., & Vorosmarty, C. J. (1992). Interactions between carbon and nitrogen dynamics in estimating net primary productivity for potential vegetation in north america. *Global Biogeochemical Cycles*, 6. <https://doi.org/10.1029/92GB00219>
- Raich, J. W. (1991). Potential net primary productivity in south america: Application of a global model. *Ecological Applications*, 1. <https://doi.org/10.2307/1941899>
- Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., & Laaksonen, A. (2022). The arctic has warmed nearly four times faster than the globe since 1979. *Communications Earth and Environment*, 3. <https://doi.org/10.1038/s43247-022-00498-3>
- Schädel, C., Rogers, B. M., Lawrence, D. M., Koven, C. D., Brovkin, V., Burke, E. J., Genet, H., Huntzinger, D. N., Jafarov, E., McGuire, A. D., Riley, W. J., & Natali, S. M. (2024). Earth system models must include permafrost carbon processes. In *Nature Climate Change* (Vol. 14). <https://doi.org/10.1038/s41558-023-01909-9>
- Scheller, R. M., Domingo, J. B., Sturtevant, B. R., Williams, J. S., Rudy, A., Gustafson, E. J., & Mladenoff, D. J. (2007). Design, development, and application of LANDIS-II, a spatial landscape simulation model with flexible temporal and spatial resolution. *Ecological Modelling*, 201(3-4), 409–419. <https://doi.org/10.1016/j.ecolmodel.2006.10.009>
- Schuur, Edward A. G., Abbott, B. W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G., Grosse, G., Jones, M., Koven, C., Leshyk, V., Lawrence, D., Loranty, M. M., Mauritz,

- M., Olefeldt, D., Natali, S., Rodenhizer, H., Salmon, V., Schädel, C., Strauss, J., ... Turetsky, M. (2022). Permafrost and climate change: Carbon cycle feedbacks from the warming arctic. *Annual Review of Environment and Resources*, 47, 343–371. <https://doi.org/10.1146/annurev-environ-012220-011847>
- Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., & Vonk, J. E. (2015). Climate change and the permafrost carbon feedback. *Nature*, 520, 171–179. <https://doi.org/10.1038/nature14338>
- Seidl, R., Rammer, W., Scheller, R. M., & Spies, T. A. (2012). An individual-based process model to simulate landscape-scale forest ecosystem dynamics. *Ecological Modelling*, 231, 87–100. <https://doi.org/10.1016/j.ecolmodel.2012.02.015>
- Smith, S. L., O'Neill, H. B., Isaksen, K., Noetzli, J., & Romanovsky, V. E. (2022). The changing thermal state of permafrost. In *Nature Reviews Earth and Environment* (Vol. 3). <https://doi.org/10.1038/s43017-021-00240-1>
- Yi, S., McGuire, A. D., Harden, J., Kasischke, E., Manies, K., Hinzman, L., Liljedahl, A., Randerson, J., Liu, H., Romanovsky, V., Marchenko, S., & Kim, Y. (2009). Interactions between soil thermal and hydrological dynamics in the response of alaska ecosystems to fire disturbance. *Journal of Geophysical Research: Biogeosciences*, 114. <https://doi.org/10.1029/2008JG000841>
- Yi, S., McGuire, A. D., Kasischke, E., Harden, J., Manies, K., MacK, M., & Turetsky, M. (2010). A dynamic organic soil biogeochemical model for simulating the effects of wildfire on soil environmental conditions and carbon dynamics of black spruce forests. *Journal of Geophysical Research: Biogeosciences*, 115. <https://doi.org/10.1029/2010JG001302>