




fireexposuR: An R package for computing and visualizing wildfire exposure

Air M. Forbes ¹ and Jennifer L. Beverly ¹

¹ Department of Renewable Resources, University of Alberta, Edmonton, AB T6G 2H1, Canada

DOI: [10.21105/joss.08479](https://doi.org/10.21105/joss.08479)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Mengqi Zhao](#)  

Reviewers:

- [@thomasarsouze](#)
- [@mstorey87](#)

Submitted: 21 May 2025

Published: 30 October 2025

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

`fireexposuR` (v1.2.0) is an R package that automates wildfire exposure methodologies presented in several scientific publications ([Beverly et al., 2010, 2021](#); [Beverly & Forbes, 2023](#)), providing an accessible and adaptable platform to a broad user base. Wildfire exposure is a numeric rating of potential wildfire transmission to, or from, a location based on the composition and configuration of the surrounding wildland fuels (i.e., flammable vegetation). Exposure assessments can be applied at multiple spatial scales (e.g., neighbourhood, region, country) to inform a wide range of management objectives and decisions (e.g., fuel management priorities, evacuation planning, land-use planning, habitat conservation, firefighting resource optimization); and are particularly well-suited for scenario and contingency planning. `fireexposuR` has functions for computational analysis and generating visualizations of results in summary tables, plots, and maps.

Background

Wildfire exposure assessments were originally introduced to assess fire entry points to the built environment ([Beverly et al., 2010](#)). The method is promoted nationally by FireSmart Canada as a tool for community protection planning ([Firesmart Canada, 2018](#)). The approach was later adapted for landscape scale assessments by Beverly et al. (2021), which included a performance validation confirming real wildfires burn preferentially in high exposure areas. Exposure assessments are omnidirectional but can be summarized locally to assess the directional vulnerability of a given location to potential fire encroachment using radial graphs developed by Beverly & Forbes (2023). Outputs of wildfire exposure and directional vulnerability assessments are typically well received and easily understood by diverse audiences due to the intuitive visualizations. The methods and data requirements are also comparatively simpler than alternative wildfire risk models, making them an ideal option when constrained by budget or time. These methods were developed in Alberta, Canada; but have since been adapted successfully to other geographic areas. Schmidt et al. (2024) customized exposure assessments for use in Alaska, USA, Khan et al. (2025) have applied them in Portugal, and a publication is currently in preparation that applies and validates the exposure metric across the entire Canadian land base ([Beverly & Mahler, 2025](#)). Research is also underway to adapt the methods for use in the United Kingdom. The exposure metric has been used as an input to evacuation planning ([Kim et al., 2024](#)), fuel treatment planning ([Karimi et al., 2024](#)), and to assess telecommunication vulnerability ([Kuiper & Beverly, 2025](#)).

Statement of need

Broad distribution and uptake of exposure and directional vulnerability assessments has been constrained by the technical nature of the original documentation within scientific publications,

some with access fees; and the need for proficiency in, and access to, geographic information system (GIS) software. The published methods were described in relation to Esri Inc. ArcGIS software and geoprocessing tools which has a subscription-based model that can be financially prohibitive to some users.

Conceptually, the methodologies are relatively simple, but they involve a detailed sequence of data processing steps that can intimidate new users, especially those lacking foundational knowledge in GIS and spatial analysis. New users may introduce unintentional errors, by misinterpreting processing steps or input data requirements and can lack confidence in their understanding of the methods and their process for executing the analysis independently. The objectives of `fireexposuR` are to reduce barriers to access and give users structured support to independently apply the methods with confidence. The analysis functions were developed and vetted in collaboration with the primary author of the wildfire exposure publications upon which `fireexposuR` is based. Custom parameterization options have been designed and described for adaptability within the suggested scope. In addition, a validation function has been included to confirm any custom decisions in alternative use cases or geographic areas. The package was developed in the R environment (R Core Team, 2024), an open-source platform with no financial fees. Package documentation is presented in plain language to ensure access to a non-technical audience. Access to the package requires basic proficiency in the R environment, which could present a barrier to some users; however, abundant online resources and tutorials are available to anyone willing to learn.

State of the field

At the time of writing, `fireexposuR` is the only fully open-source software option for computing and visualizing wildfire exposure. A custom Python toolbox has also been developed to replicate the wildfire exposure methods in Schmidt et al. (2024), and is freely available on [GitHub](#) (v1.0.5; GINA/ISER (2024)) but requires an ArcGIS license. The toolbox guides users through the steps of defining a hazardous fuel raster, computing ember exposure for fixed distances, and other additional steps specific to the methods presented in Schmidt et al. (2024). In comparison, `fireexposuR` includes transmission distance customization, directional vulnerability assessments, and rapid visualization of results with sensible defaults.

Target audience

The target audience for `fireexposuR` includes researchers, agencies, practitioners, and individuals who are interested in conducting wildfire exposure assessments. Package functions are highly customizable, for ease of application in new geographic areas or use cases.

Typical workflow

The workflow for conducting wildfire exposure assessments using the functions within the `fireexposuR` package is demonstrated in [Figure 1](#). Input data requirements are detailed in the function documentation.




Figure 1: Figure 1. Suggested workflow of core fireexposuR functionality. Computational outputs, outlined in navy, can be further manipulated in R, visualized with fireexposuR functions, or exported. Images and tables can be exported with base R and spatial data can be exported with {terra}.

Acknowledgements

The fireexposuR project was funded by the National Research Council of Canada, the Forest Resource Improvement Association of Alberta, and Housing, Infrastructure and Communities Canada. We thank ROpenSci reviewers Ronny Hernandez Mora (@ronnyhdez) and Sherry Zhang (@huizezhang-sherry) for their helpful feedback, which significantly improved the code and documentation in this R package. We also thank the many early users of the package for providing feedback during initial development.

References

Beverly, J. L., Bothwell, P., Conner, J. C. R., & Herd, E. P. K. (2010). Assessing the exposure of the built environment to potential ignition sources generated from vegetative fuel. *International Journal of Wildland Fire*, 19(3), 299. <https://doi.org/10.1071/WF09071>

- Beverly, J. L., & Forbes, A. M. (2023). Assessing directional vulnerability to wildfire. *Natural Hazards*, 117(1), 831–849. <https://doi.org/10.1007/s11069-023-05885-3>
- Beverly, J. L., & Mahler, P. (2025). 2024 *National Landscape Fire Exposure Map of Canada*. <https://doi.org/10.17605/OSF.IO/623JM>
- Beverly, J. L., McLoughlin, N., & Chapman, E. (2021). A simple metric of landscape fire exposure. *Landscape Ecology*, 36(3), 785–801. <https://doi.org/10.1007/s10980-020-01173-8>
- Firesmart Canada. (2018). *Wildfire exposure assessment*. https://wildfireanalytics.org/uploads/1/1/2/4/112421061/fs_exposureassessment_sept2018.pdf
- GINA/ISER. (2024). *Gina-alaska/wildfire-exposure-toolbox: Repository for GINA/ISER ArcGIS Wildfire Exposure Python Toolbox*. <https://github.com/gina-alaska/wildfire-exposure-toolbox/releases/tag/v1.0.5>
- Karimi, N., Mahler, P., & Beverly, J. L. (2024). Optimizing fuel treatments for community wildfire mitigation planning. *Journal of Environmental Management*, 370, 122325. <https://doi.org/10.1016/j.jenvman.2024.122325>
- Khan, S. I., Beverly, J. L., Colaço, M. C., Rego, F. C., & Sequeira, A. C. (2025). Applying a fire exposure metric in the artificial territories of portugal: Mafra municipality case study. *Fire*, 8(5). <https://doi.org/10.3390/fire8050179>
- Kim, A. M., Beverly, J. L., & Al Zahid, A. (2024). Directional analysis of community wildfire evacuation capabilities. *Safety Science*, 171, 106378. <https://doi.org/10.1016/j.ssci.2023.106378>
- Kuiper, C., & Beverly, J. (2025). *Assessing Telecommunication Vulnerabilities to Wildfire in Alberta, Canada*. Social Science Research Network. <https://doi.org/10.2139/ssrn.5119206>
- R Core Team. (2024). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://doi.org/10.32614/r.manuals>
- Schmidt, J. I., Ziel, R. H., Calef, M. P., & Varvak, A. (2024). Spatial distribution of wildfire threat in the far north: Exposure assessment in boreal communities. *Natural Hazards*. <https://doi.org/10.1007/s11069-023-06365-4>