

# Lithology: A Landlab submodule for spatially variable rock properties

Katherine R. Barnhart<sup>1, 2</sup>, Eric Hutton<sup>3, 4</sup>, Nicole M. Gasparini<sup>5</sup>, and Gregory E. Tucker<sup>1, 2, 3</sup>

**1** University of Colorado at Boulder, Department of Geological Sciences **2** University of Colorado at Boulder, Cooperative Institute for Research in Environmental Sciences **3** University of Colorado at Boulder, Community Surface Dynamics Modeling System Integration Facility **4** University of Colorado at Boulder, Institute for Arctic and Alpine Research **5** Tulane University, Department of Earth and Environmental Sciences

DOI: [10.21105/joss.00972](https://doi.org/10.21105/joss.00972)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

**Submitted:** 22 September 2018

**Published:** 23 September 2018

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

## Summary

The surface of the Earth reflects the competing advection of rock by tectonic processes and the erosion of rock by wind, water, and ice. Rock properties influence erosion rates by changing the processes responsible for erosion and the rate at which rock is weathered, detached, and turned into mobile sediment. Variations in the rock properties over space and with depth reflect the legacy of sedimentary deposition and tectonic deformation. Long-term landscape evolution modeling experiments that include the impact of spatially and temporally variable rock characteristics can be used to identify the impact of rock strength patterns on other geologic observables such as topography, erosion rates, and detrital mineral records (e.g. (Forte, Yanites, & Whipple, 2016), (Perne, Covington, Thaler, & Myre, 2017)). Identifying these relationships allows for better interpretations of the geologic record.

Landlab is an Open Source Python package that provides a framework for the development of 2D numerical models, typically in Earth surface dynamics ((Hobley et al., 2017)). Landlab was designed as a modular framework in which different process components can be mixed and matched to construct a model based on a user's needs. Prior work on spatially variable lithology in landscape evolution has been done using a modified version of the channel-hillslope integrated landscape development (CHILD, (G. Tucker, Lancaster, Gasparini, & Bras, 2001)) model (e.g.(???)) and the FastScape V5 model ((Braun & Willett, 2013), (Perne et al., 2017)). To provide these capabilities within the Landlab framework, there is a need for a Landlab submodule that can treat spatial variations in rock materials.

This contribution describes **Lithology**, a Landlab submodule designed to support the representation of 3D variations in rock material properties within the Landlab framework. It includes two classes: **Lithology** is a generic representation of spatially varying rock material, and **LithoLayers** is a derived class that treats parallel layers of material variations. In both classes, each rock type may have multiple attributes. Rock layers may be removed through erosion, or added to through deposition. Two options for the underlying datastructure are supported: “event layers”, in which the data structure stores each time-step as an event even if there is no material in the layer, or “material layers”, in which entries in the datastructure represent contiguous material of the same property, but not necessarily the same age. This second option is more memory efficient but does not record the transient dynamics of erosion and deposition.

Source code for `Lithology` and `Litholayers` is available as part of the [Landlab python package](#). The `Lithology` submodule is documented using Docstrings, and the documentation can be found on the Landlab ReadTheDocs site. One page exists for the [Lithology component](#) and a second for the [LithoLayers component](#). Unit and docstring tests provide 100% coverage of this submodule. [Pull Request # 674](#) brought the `Lithology` submodule into the core Landlab source code (note: as of initial JOSS submission this PR is still active as we anticipate making revisions before merging). A tagged version of Landlab that includes the `Lithology` submodule is archived in Zenodo with the linked DOI: [DOI:10.5281/zenodo.1234567](#) (note: doi not active for initial JOSS submission).

The Landlab project maintains a separate repository containing tutorials that introduce core concepts and the use of individual submodules. In addition to the source code, a Jupyter Notebook introducing the use of `Lithology` and `Litholayers` is now part of the Landlab tutorials repository. This tutorial was brought into the repository with [Pull Request # 19](#) (note: as of initial JOSS submission this PR is still active as we anticipate making revisions before merging). A tagged version of Landlab that includes this functionality has been archived with the following DOI: [DOI:10.5281/zenodo.1234567](#) (note: doi not active for initial JOSS submission).

## Acknowledgements

The authors thank Adam Forte, Matt Rossi, and Brian Yanites for helpful discussions during the development of this code and the accompanying Jupyter notebooks.

## References

- Braun, J., & Willett, S. D. (2013). A very efficient  $O(n)$ , implicit and parallel method to solve the stream power equation governing fluvial incision and landscape evolution. *Geomorphology*, 180, 170–179.
- Forte, A. M., Yanites, B. J., & Whipple, K. X. (2016). Complexities of landscape evolution during incision through layered stratigraphy with contrasts in rock strength. *Earth Surface Processes and Landforms*, 41(12), 1736–1757. doi:[10.1002/esp.3947](#)
- Hobley, D. E. J., Adams, J. M., Nudurupati, S. S., Hutton, E. W. H., Gasparini, N. M., Istanbuluoglu, E., & Tucker, G. E. (2017). Creative computing with landlab: An open-source toolkit for building, coupling, and exploring two-dimensional numerical models of earth-surface dynamics. *Earth Surface Dynamics*, 5(1), 21–46. doi:[10.5194/esurf-5-21-2017](#)
- Perne, M., Covington, M. D., Thaler, E. A., & Myre, J. M. (2017). Steady state, erosional continuity, and the topography of landscapes developed in layered rocks. *Earth Surface Dynamics*, 5(1), 85–100.
- Tucker, G., Lancaster, S., Gasparini, N., & Bras, R. (2001). The channel-hillslope integrated landscape development model (child). In *Landscape erosion and evolution modeling* (pp. 349–388). Springer.
- ( ).