

INS_IEP: A MATLAB package for fitting Inelastic Neutron Scattering data.

Alban Bloor Riley¹, Marcus Webb¹, and Michael L. Baker²

¹ The University of Manchester, Department of Mathematics ² The University of Manchester, Department of Chemistry ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- Review
- Repository
- Archive

Editor: [Mojtaba Barzegari](#)

Reviewers:

- @KedoKudo
- @granrothge

Submitted: 07 July 2025

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Inelastic neutron scattering (INS) is a spectroscopic technique that can be used to measure the magnetic excitations in materials with interacting electron spins. For samples composed of finite-size clusters of magnetic moment-carrying atoms, such as single ions or molecular-based magnets, INS experiments yield inelastic excitation with energies that correspond to the eigenvalues of the spin Hamiltonian of the material being studied. Fitting a model spin system to these experimental eigenvalues can be formulated as an inverse eigenvalue problem (IEP), where the matrix formed is the spin Hamiltonian operator of the sample molecule. INS_IEP is a MATLAB package that uses deflated numerical optimisation methods to systematically find multiple solutions to this IEP. The package requires and is fully compatible with easyspin (Stoll & Schweiger, 2006), a package for solving fitting problems in electron paramagnetic resonance (EPR).

Statement of need

Neutrons are an excellent bulk probe of material properties since they carry no charge and therefore penetrate deeply into matter. Neutrons also carry a quantum spin of a half, making them a sensitive probe of magnetism (Squires, 2012). Reactors and spallation sources with dedicated high-flux neutron sources serve the international research community with neutron scattering experiment capabilities for material research. Inelastic neutron scattering (INS) is one such experimental technique that can be used to study magnetism. In an INS experiment, a sample under investigation is irradiated with a beam of neutrons and the scattered neutron energy and momentum transfer are detected. For samples composed of finite-size clusters of magnetic moment-carrying atoms, such as single ions or molecular-based magnets, the detected neutron energy transfer gives direct access to the quantum spin excitations (M. Baker & Mutka, 2012; Furrer & Waldmann, 2013; "Spectroscopy Methods for Molecular Nanomagnets," 2014).

The energy of such excitations relates to the energy difference between eigenvalues of the Hamiltonian matrix that describes the quantum spin dynamics of the compound in question. Single-ion and molecular-based magnets are studied as prototype components (quantum bits, sensors) for quantum technologies. INS can therefore provide crucial information concerning the precise quantum properties of such systems. However, to relate the INS experimental results to the Hamiltonian that describes quantum spin dynamics requires parameterisation of matrix elements such that a set of eigenvalues and eigenstates matching the experiment are determined. This situation is known as the inverse eigenvalue problem.

To date, this problem is addressed in an iterative process where parameters of the Hamiltonian are varied manually, often one at a time, and the resultant eigenvalues compared to the experimental values - each such iteration requires an eigendecomposition of the Hamiltonian matrix. INS_IEP presents an elegant solution to solving this problem, using algorithms to

calculate multiple parameter sets that minimise the difference in eigenvalues, reducing the number of Hamiltonian matrix diagonalisations, and providing a more robust method to reliably extract an accurate spin Hamiltonian model from INS experimental data.

Key Concepts

The Spin Hamiltonian

The Spin Hamiltonian, H , is an approximation of the Hamiltonian that uses spin coordinates instead of orbital coordinates, and is widely used to model data arising from many spectroscopy techniques (Launay & Verdaguer, 2014). It can be modeled as a linear combination of interaction terms; in this package we will use the zero field interaction, H_{ZFI} , and the electron-electron interaction, H_{EEI} :

$$H = H_{ZFI} + H_{EEI}.$$

Both of these terms can themselves be modelled as the linear sum of other basis matrices. The zero field interaction can be written as:

$$H_{ZFI} = \sum_{-k \leq q \leq k} B_k^q O_k^q$$

where the O_k^q are Stevens Operators (Rudowicz & Chung, 2004), and B_k^q the associated parameter. When there are multiple spin centres it is necessary to take Kronecker products of the operator with identity matrices of the appropriate for each other spin centre.

When there are multiple spin centres it is also necessary to include an electron-electron interaction term, H_{EEI} . This term will be the sum of interaction terms between each pair of spin centres:

$$H_{EEI} = - \sum_{i \neq j} J_{ij} S_i \cdot S_j$$

where S_i is the vector of spin operators $S_i = [S_x, S_y, S_z]$ for the i -th spin centre, and J_{ij} is the parameter to be found that represents the strength of interaction between the two spin centres. Note that in the isotropic case J can be thought of as a scalar value, but in the anisotropic case will be a matrix where the off diagonals are skew symmetric (often the off diagonals are assumed to be zero). While the summation is in theory over all spin centre combinations, in practice many of these contributions will be negligible - often only the nearest neighbour interactions are significant.

It is important to mention that these matrix operators can be very large. The size is defined by the number of spin centres (n) and the spin (S_i) of each spin centre. The dimension of matrices is given by:

$$\prod_i^n (2S_i + 1).$$

The operators are however highly sparse, this means that it is possible to use eigensolvers that can take advantage of this sparsity.

Inverse Eigenvalue Problem

The INS experiments provide eigenvalues of the Spin Hamiltonian matrix of the sample, the task of calculating the matrix from the eigenvalues is an inverse eigenvalue problem:

75 Let $A(x)$ be the affine family of matrices,

$$A(x) = A_0 + \sum_{i=1}^{\ell} x_i A_i,$$

76 where $x \in \mathbb{C}^{\ell}$ and $A_0, \dots, A_{\ell} \in \mathbb{R}^{n \times n}$ are linearly independent Hermitian matrices, and denote
77 the ordered eigenvalues of $A(x)$ as $\lambda_1(x) \leq \dots \leq \lambda_n(x)$. Then the least squares inverse
78 eigenvalue problem (LSIEP) is to find the parameters $x \in \mathbb{R}^{\ell}$ that minimises

$$F(x) = \frac{1}{2} \|r(x)\|_2^2 = \frac{1}{2} \sum_{i=1}^m (\lambda_i(x) - \lambda_i^*)^2$$

79 where $\lambda_1^* \leq \dots \leq \lambda_m^*$ are the experimental eigenvalues (Chu & Golub, 2005). In the case of INS
80 fitting the A_i basis matrices will be a combination of Stevens operators and electron-electron
81 exchange terms. The IEP described above is formulated as an least squares problem because
82 the number of eigenvalues that can be probed by INS experiments is often a small subset of
83 the full spectrum. Due to the low temperatures that these experiments are performed at (can
84 be as low as 1K) it is generally the smallest eigenvalues that are involved. Note also that since
85 it is the energy difference between the eigenvalues that is probed we actually have to modify
86 the IEP - either by adding an additional parameter (an identity matrix) that shifts the values
87 of the eigenvalues, or by changing the above formula for F to directly sum the difference in
88 eigenvalues thereby reducing the number of residual equations in $r(x)$ by one.

89 As far as we are aware this is the first time that the fitting of INS data has been explicitly
90 formulated as an IEP. An advantage of this formulation is that there are explicit formulas for
91 the derivatives of $r(x)$. The first derivative (Jacobian) is:

$$J_r(x) = \begin{pmatrix} q_1(x)^T A_1 q_1(x) & \dots & q_1(x)^T A_{\ell} q_1(x) \\ \vdots & \ddots & \vdots \\ q_m(x)^T A_1 q_m(x) & \dots & q_m(x)^T A_{\ell} q_m(x) \end{pmatrix},$$

92 and the second derivative (Hessian) is:

$$(H_r)_{ij} = 2 \sum_{k=1}^m (\lambda_k - \lambda_k^*) \sum_{\substack{t=1 \\ \lambda_t \neq \lambda_k}}^m \frac{(q_t^T A_i q_k)(q_t^T A_j q_k)}{\lambda_k - \lambda_t}.$$

93 Another advantage is the number of constraints to fit is much smaller than fitting the spectrum
94 itself, as it corresponds to fitting only the locations of the peaks of the spectrum.

95 Methods

96 All of the methods used are iterative schemes of the form $x^{k+1} = x^k + p^k$ where the step p^k
97 uniquely defines each algorithm:

- 98 ■ Newton's method: $p^k = (J_r^T J_r + H_r r) - 1 J_r^T r$ (Nocedal & Wright, 2006)
- 99 ■ Gauss-Newton method: $p^k = (J_r^T J_r)^{-1} J_r^T r$ (Nocedal & Wright, 2006)
- 100 ■ Lift and Projection Method: $p^k = B^{-1} J_r^T r$ (Bloor Riley et al., 2025a)

101 Where the matrix B is the Gram matrix formed from the frobenius inner products of the basis
102 matrices: $B_{ij} = \langle A_i, A_j \rangle_F$. The Lift and Projection method is a Riemannian Gradient descent
103 method (Bloor Riley et al., 2025a), inspired by the Lift and Projection method (Chu & Golub,
104 2005), specifically designed for solving IEPs. In (Bloor Riley et al., 2025a) it is proven that the
105 method is a strictly descending algorithm, that is it reduces the value of the objective function
106 every step. Both the deflated Gauss-Newton method and the Riemannian Gradient descent

107 Lift and Projection method are new methods designed for this package (Bloor Riley et al.,
108 2025a, 2025b).

109 Deflation

110 The number of eigenvalues that can be probed via INS experiments varies depending on the
111 equipment and sample in question, meaning that the fitting problem is often under or even
112 over determined. The IEP is also highly nonlinear and due to the experimental nature of the
113 data may be ill-posed. One consequence of this is that the solution space may be very 'bumpy',
114 that is there may exist many local minimisers to the problem. For example in Figure 1, there
115 are clearly 4 distinct solutions (for more details see Example 1 and the file Example1_Mn12.m
116 in the examples folder). We seek to solve the problem of multiple local minima by the use
117 of Deflation, a numerical technique used to find multiple solutions to systems of equations
118 (Farrell et al., 2015). Fortunately it is cheap to apply deflation for the above methods, it is
119 simply a change to the length of the step - notably this means that the direction of each step
120 does not change. It is proven in (Bloor Riley et al., 2025b) that the deflated methods will not
121 converge to deflated points. The usual requirements still apply to the convergence of the new
122 methods - that the initial guess is close enough to the new minimum, and that the Jacobian is
123 full rank in a neighbourhood around that minimum. The rate of convergence of the deflated
124 methods is also more complicated, although the number of iterations required to converge can
125 go up with the number of deflations this is not a strict correlation, as can be seen in Figure 2.




Figure 1: Contour plot of how F varies with the two parameters B_2^2 and B_4^4 for the molecule Mn_12 as described in Example 1. There are four locally minimising parameter pairs corresponding to the four blue regions




Figure 2: Comparison of the convergence behaviour for computing each solution in Example 1. The gradient of the lines as the method approaches the solution shows how the methods are quadratically convergent local to a minimum.

Examples

Example 1 - Mn12

The first example we will look at is Manganese-12-acetate. This is a well known example in the INS and magnetism community, as one of the first molecules that behaves like a nano-sized magnet with a molecular magnetic coercivity as well as its role in the research of quantum tunnelling of magnetisation (Friedman et al., 1996; Sessoli et al., 1993).

The Spin Hamiltonian of this system, using the giant spin approximation, can be represented as a 21×21 matrix modelled using 4 Stevens operators (Bircher et al., 2004):

$$H = B_2^0 O_2^0 + B_4^0 O_4^0 + B_2^2 O_2^2 + B_4^4 O_4^4 \in \mathbb{R}^{21 \times 21}$$

We utilise the same spin system syntax as easyspin, so to set up the problem we first set up the model, along with initial guesses for the parameters:

```
%One spin centre (because giant spin approximation)
Sys0.S=10;
%Four Stevens operators
Sys0.B2 = [-100,0,-1000,0,0];
Sys0.B4 = [-1,0,0,0,-1,0,0,0,0];
```

Then we input the experimental eigenvalues - these are typically shifted such that the smallest eigenvalue is zero - and define which parameters to fit. Note that all values given must be in Hertz, so it may be useful to use conversions.

```
rcm = 29979.2458; meV = rcm*8.065; %Conversions values
%Input calculated eigenvalues:
Exp.ev = [0,0,1.24,1.24,2.3,2.3,3.18,3.18,3.91,3.91,4.5,4.5,
4.97,4.97,5.32,5.32,5.54,5.59,5.69,5.75,5.78].*meV;
```

```
%Vary all non zero parameters (no Fixed parameters):
Vary = Sys0;
```

139 Then all that is required is to call INS_IIEP with these three inputs:

```
SysOut = INS_IIEP(Sys0,Vary,Exp);
```

140 If we wish to find all four solutions as shown in Figure 1 then we use the additional option:

```
Opt.NDeflations = 4;
SysOut = INS_IIEP(Sys0,Vary,Exp,Opt);
```

141 In this case SysOut will be an array of four spin structures each containing a distinct locally optimal solution. It is possible to access information about the convergence of each deflation by using SysOut.Output. For example by utilising the iterates recorded, stored in SysOut.Output.Iterates it is possible to plot a graph of convergence, as can be seen in Figure 2. The Output structure also contains the value of F at the final point, as well as the number of iterations it took to get there.

147 A full list of options is provided in the help of INS_IIEP.

148 Example 2 - Chromium(iii) Horseshoes

149 The second example concerns antiferro-magnetically coupled chain of six chromium(III) ions (M. L. Baker et al., 2011). Because there are multiple spin centres an electron-electron interaction term is required. The spin hamiltonian is a 4096×4096 matrix composed of two Stevens operators and one interaction term, since it is known a priori that each spin centre will have the same value parameters we pin the parameters here, by setting the initial guess as the same value:

```
Sys0.S = [1.5 1.5 1.5 1.5 1.5 1.5];
Sys0.B2 = [1 0 -1 0 0;
1 0 -1 0 0;
1 0 -1 0 0;
1 0 -1 0 0;
1 0 -1 0 0;
1 0 -1 0 0];
Sys0.J = [100,0,0,0,0,100,0,0,0,100,0,0,100,0,100];
Vary = Sys0;
Exp.ev = [0,0.355,0.457,0.497,1.576,1.577,1.592,1.629,1.632,
2.97,2.98,3.002,3.004,3.01,3.038,3.821,3.824,3.827,
3.837,3.856,3.879,3.888,3.895,3.903].*meV;
```

155 Note that only 24 eigenvalues were found experimentally, so this will form a partial LSIEP. To find the solution system is as simple as:

```
Opt.GradientTolerance = 1e-3; %Use additional stopping criterion.
SysOut= INS_IIEP(Sys1,Vary1,Exp,Opt);
```

157 It is possible to find multiple minimising systems even if they do not make any sense physically, however due to the scaling of the problem a change in the default deflation parameters is necessary:

```
Opt = struct('NDeflations',5,'Sigma',1e-7,'StepTolerance',1e-3 );
SysOut= INS_IIEP(Sys0,Vary,Exp,Opt);
```

160 The output contains five different spin systems that all have the same eigenvalues as input. Like in example one some of the minima found differ only by a sign change of the B_2^2 parameter, but all minima have the same exchange term.

163 Additional examples can be found in the Examples folder.

Note also that using mint (M. L. Baker, 2022), which is fully compatible with INS_IEP it is possible to simulate the INS spectrum of any calculated system - which can then be compared to the experimental spectrum. A description of how to do this can also be found in the examples folder.

Acknowledgements

ABR thanks the University of Manchester for a Dean's Doctoral Scholarship. MW thanks the Polish National Science Centre (SONATA-BIS-9), project no. 2019/34/E/ST1/00390, for the funding that supported some of this research.

References

- Baker, M. L. (2022). *Mint: Simulate inelastic neutron scattering induced magnetic excitations for single-ions, molecular-magnets and finite exchange coupled spin-systems*. <https://mlbakerlab.co.uk/mint/>
- Baker, M. L., Bianchi, A., Carretta, S., Collison, D., Docherty, R. J., McInnes, E. J. L., McRobbie, A., Muryn, C. A., Mutka, H., Piligkos, S., Rancan, M., Santini, P., Timco, G. A., Tregenna-Piggott, P. L. W., Tuna, F., Güdel, H. U., & Winpenny, R. E. P. (2011). Varying spin state composition by the choice of capping ligand in a family of molecular chains: Detailed analysis of magnetic properties of chromium(iii) horseshoes. *Dalton Transactions*, 40(12), 2725. <https://doi.org/10.1039/c0dt01243b>
- Baker, M., & Mutka, H. (2012). Neutron spectroscopy of molecular nanomagnets. *The European Physical Journal Special Topics*, 213, 53–68. <https://doi.org/10.1140/epjst/e2012-01663-6>
- Bircher, R., Chaboussant, G., Sieber, A., Güdel, H. U., & Mutka, H. (2004). Transverse magnetic anisotropy in Mn 12 acetate: Direct determination by inelastic neutron scattering. *Phys. Rev. B*, 70(21), 212413. <https://doi.org/10.1103/PhysRevB.70.212413>
- Bloor Riley, A., Webb, M., & Baker, M. L. (2025a). *A Riemannian Gradient Descent Method for the Least Squares Inverse Eigenvalue Problem*. arXiv. <https://doi.org/10.48550/arXiv.2504.07809>
- Bloor Riley, A., Webb, M., & Baker, M. L. (2025b). *Deflation Techniques for Finding Multiple Local Minima of a Nonlinear Least Squares Problem*. arXiv. <https://doi.org/10.48550/arXiv.2409.14438>
- Chu, M. T.-C., & Golub, G. H. (2005). *Inverse Eigenvalue problems: Theory, algorithms, and applications*. Oxford University Press. ISBN: 978-0-19-856664-9
- Farrell, P. E., Á. Birkisson, & Funke, S. W. (2015). Deflation Techniques for Finding Distinct Solutions of Nonlinear Partial Differential Equations. *SIAM J. Sci. Comp.*, 37(4), A2026–A2045. <https://doi.org/10.1137/140984798>
- Friedman, J. R., Sarachik, M. P., Tejada, J., & Ziolo, R. (1996). Macroscopic Measurement of Resonant Magnetization Tunneling in High-Spin Molecules. *Phys. Rev. Lett.*, 76(20), 3830–3833. <https://doi.org/10.1103/physrevlett.76.3830>
- Furrer, A., & Waldmann, O. (2013). Magnetic cluster excitations. *Reviews of Modern Physics*, 85, 367–420. <https://doi.org/10.1103/RevModPhys.85.367>
- Launay, J.-P., & Verdaguer, M. (2014). *Electrons in molecules: From basic principles to molecular electronics* (First edition). Oxford University Press. ISBN: 978-0-19-929778-8
- Nocedal, J., & Wright, S. J. (2006). *Numerical optimization* (2nd ed). Springer. ISBN: 978-0-387-30303-1

- 208 Rudowicz, C., & Chung, C. Y. (2004). The generalization of the extended Stevens operators
209 to higher ranks and spins, and a systematic review of the tables of the tensor operators
210 and their matrix elements. *Journal of Physics: Condensed Matter*, 16(32), 5825–5847.
211 <https://doi.org/10.1088/0953-8984/16/32/018>
- 212 Sessoli, R., Gatteschi, D., Caneschi, A., & Novak, M. A. (1993). Magnetic bistability in a
213 metal-ion cluster. *Nature*, 365(6442), 141–143. <https://doi.org/10.1038/365141a0>
- 214 Spectroscopy Methods for Molecular Nanomagnets. (2014). In *Structure and Bonding* (pp.
215 231–291). Springer Berlin Heidelberg. https://doi.org/10.1007/430_2014_155
- 216 Squires, G. L. (2012). *Introduction to the Theory of Thermal Neutron Scattering* (3rd ed.).
217 Cambridge University Press. <https://doi.org/10.1017/CBO9781139107808>
- 218 Stoll, S., & Schweiger, A. (2006). EasySpin, a comprehensive software package for spectral
219 simulation and analysis in EPR. *Journal of Magnetic Resonance*, 178, 42–55. <https://doi.org/10.1016/j.jmr.2005.08.013>
220

DRAFT