

pynucastro: an interface to nuclear reaction rates and code generator for reaction network equations

Donald E. Willcox¹ and Michael Zingale¹

1 Department of Physics and Astronomy, Stony Brook University

DOI: 10.21105/joss.00588

Software

■ Review 🗗

Repository 2

■ Archive ♂

Submitted: 15 February 2018 Published: 10 March 2018

Licence

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC-BY).

Summary

pynucastro addresses two needs in the field of nuclear astrophysics: visual exploration of nuclear reaction rates or networks and automated code generation for integrating reaction network ODEs. pynucastro accomplishes this by interfacing with nuclear reaction rate parameterizations published by the JINA Reaclib project (Cyburt et al. 2010).

Interactive exploration is enabled by a set of classes that provide methods to visualize the temperature dependency of a rate, evaluate it at a particular temperature, and find the exponent, n, for a simple Tⁿ parameterization. From a collection of rates, the flow between the nuclei can be visualized interactively using Jupyter widgets. These features help both with designing a network for a simulation as well as for teaching nuclear astrophysics in the classroom.

After selecting a set of rates for a given problem, pynucastro can construct a reaction network from those rates consisting of Python code to calculate the ODE right hand side. Generated Python right hand sides evolve species in the reaction network, and pynucastro includes a Python example integrating the CNO cycle for hydrogen burning.

pynucastro can also generate Fortran code implementing reaction networks, using SymPy (Meurer et al. 2017) to determine the system of ODEs comprising the network. From the symbolic expressions for the ODE right hand side, pynucastro also generates a routine to compute the analytic Jacobian matrix for implicit integration.

Fortran networks incorporate weak, intermediate, and strong reaction rate screening for the Reaclib rates (Graboske et al. 1973; Alastuey and Jancovici 1978; Itoh et al. 1979). These networks can also include selected weak reaction rate tabulations (Suzuki, Toki, and Nomoto 2016). To calculate energy generation in Fortran networks, pynucastro uses nuclear binding energies from the Atomic Mass Data Center (Huang et al. 2017; Wang et al. 2017) and the 2014 CODATA recommended values for the fundamental physical constants (Mohr, Newell, and Taylor 2016).

pynucastro is capable of generating two kinds of Fortran reaction networks. The first type is a standalone network with a driver program to integrate species and energy generation using the variable-order ODE integration package VODE (Brown, Byrne, and Hindmarsh 1989). This Fortran driver program is designed to be easy to use and can integrate reaction networks significantly faster than is possible for the generated Python networks.

Secondly, pynucastro can generate a Fortran network consisting of right hand side and Jacobian modules that evolve species, temperature, and energy generation for the StarKiller Microphysics code. Via StarKiller Microphysics, astrophysical simulation codes such as Castro (Almgren et al. 2010) and Maestro (Nonaka et al. 2010) can directly use pynucastro reaction networks. pynucastro includes a carbon burning network with tabulated



A = 23 Urca weak reactions currently used for studying white dwarf convection with Maestro (Zingale et al. 2017).

Future work will focus on implementing nuclear partition functions to compute reverse reaction rates in the Reaclib library (Rauscher and Thielemann 2000; Rauscher 2003). It is also in some cases necessary to compute reverse reaction rates using detailed balance with a consistent nuclear mass model instead of using the parameterized reverse reaction rates in Reaclib (Lippuner and Roberts 2017). Additionally, work is ongoing to port the networks generated for StarKiller Microphysics to CUDA Fortran to support parallel reaction network integration on GPU systems (Zingale et al. 2017). We intend to implement this port directly into the pynucastro-generated networks.

This work was supported by DOE/Office of Nuclear Physics grant DE-FG02-87ER40317.

References

Alastuey, A., and B. Jancovici. 1978. "Nuclear reaction rate enhancement in dense stellar matter." The Astrophysical Journal 226 (December):1034-40. https://doi.org/10.1086/ 156681.

Almgren, A. S., V. E. Beckner, J. B. Bell, M. S. Day, L. H. Howell, C. C. Joggerst, M. J. Lijewski, A. Nonaka, M. Singer, and M. Zingale. 2010. "CASTRO: A New Compressible Astrophysical Solver. I. Hydrodynamics and Self-gravity." The Astrophysical Journal 715 (June):1221-38. https://doi.org/10.1088/0004-637X/715/2/1221.

Brown, Peter N., George D. Byrne, and Alan C. Hindmarsh. 1989. "VODE: A Variable-Coefficient ODE Solver." SIAM Journal on Scientific and Statistical Computing 10 (5):1038-51. https://doi.org/10.1137/0910062.

Cyburt, Richard H., A. Matthew Amthor, Ryan Ferguson, Zach Meisel, Karl Smith, Scott Warren, Alexander Heger, et al. 2010. "The JINA REACLIB Database: Its Recent Updates and Impact on Type-I X-ray Bursts." The Astrophysical Journal Supplement Series 189 (1):240. http://stacks.iop.org/0067-0049/189/i=1/a=240.

Graboske, H. C., H. E. Dewitt, A. S. Grossman, and M. S. Cooper. 1973. "Screening Factors for Nuclear Reactions. II. Intermediate Screening and Astrophysical Applications." The Astrophysical Journal 181 (April):457–74. https://doi.org/10.1086/152062.

Huang, W.J., G. Audi, Meng Wang, F.G. Kondev, S. Naimi, and Xing Xu. 2017. "The AME2016 atomic mass evaluation (I). Evaluation of input data; and adjustment procedures." Chinese Physics C 41 (3):030002. http://stacks.iop.org/1674-1137/41/i=3/a= 030002.

Itoh, N., H. Totsuji, S. Ichimaru, and H. E. Dewitt. 1979. "Enhancement of thermonuclear reaction rate due to strong screening. II - Ionic mixtures." The Astrophysical Journal 234 (December):1079-84. https://doi.org/10.1086/157590.

Lippuner, J., and L. F. Roberts. 2017. "SkyNet: A Modular Nuclear Reaction Network Library." The Astrophysical Journal Supplement Series 233 (December):18. https://doi. org/10.3847/1538-4365/aa94cb.

Meurer, Aaron, Christopher P. Smith, Mateusz Paprocki, Ondřej Čertík, Sergey B. Kirpichev, Matthew Rocklin, Amit Kumar, et al. 2017. "SymPy: Symbolic Computing in Python." PeerJ Computer Science 3 (January):e103. https://doi.org/10.7717/peerj-cs. 103.

Mohr, Peter J., David B. Newell, and Barry N. Taylor. 2016. "CODATA recommended values of the fundamental physical constants: 2014." Rev. Mod. Phys. 88 (3). American Physical Society:035009. https://doi.org/10.1103/RevModPhys.88.035009.



Nonaka, A., A. S. Almgren, J. B. Bell, M. J. Lijewski, C. M. Malone, and M. Zingale. 2010. "MAESTRO: An Adaptive Low Mach Number Hydrodynamics Algorithm for Stellar Flows." The Astrophysical Journal Supplement Series 188:358–83.

Rauscher, T. 2003. "Nuclear Partition Functions at Temperatures Exceeding 10^{10} K." The Astrophysical Journal Supplement Series 147 (August):403-8. https://doi.org/10. 1086/375733.

Rauscher, T., and F.-K. Thielemann. 2000. "Astrophysical Reaction Rates from Statistical Model Calculations." Atomic Data and Nuclear Data Tables 75 (May):1-351. https://doi.org/10.1006/adnd.2000.0834.

Suzuki, Toshio, Hiroshi Toki, and Ken'ichi Nomoto. 2016. "Electron-capture and β -decay Rates for sd-Shell Nuclei in Stellar Environments Relevant to High-density O-Ne-Mg Cores." The Astrophysical Journal 817 (2):163. http://stacks.iop.org/0004-637X/817/i= 2/a = 163.

Wang, Meng, G. Audi, F.G. Kondev, W.J. Huang, S. Naimi, and Xing Xu. 2017. "The AME2016 atomic mass evaluation (II). Tables, graphs and references." Chinese Physics C 41 (3):030003. http://stacks.iop.org/1674-1137/41/i=3/a=030003.

Zingale, M., A. S. Almgren, M. G. Barrios Sazo, V. E. Beckner, J. B. Bell, B. Friesen, A. M. Jacobs, et al. 2017. "Meeting the Challenges of Modeling Astrophysical Thermonuclear Explosions: Castro, Maestro, and the AMReX Astrophysics Suite." ArXiv E-Prints, November. http://adsabs.harvard.edu/abs/2017arXiv171106203Z.