

# SpectralUnmixing: A general Julia package for unmixing spectroscopy data

Philip G. Brodrick<sup>1</sup>, Francisco Ochoa<sup>2</sup>, Gregory Okin<sup>2</sup>, Vatsal A. Jhalani<sup>1</sup>, Winston Olson-Duvall<sup>1</sup>, Sarah R. Lundeen<sup>1</sup>, David R. Thompson<sup>1</sup>, and Robert O. Green<sup>1</sup>

<sup>1</sup> Jet Propulsion Laboratory, California Institute of Technology, USA <sup>2</sup> University of California, Los Angeles, USA

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Rachel Kurchin](#)

## Reviewers:

- [@1oly](#)
- [@LaurenceDenneulin](#)

Submitted: 06 June 2025

Published: unpublished

## License

Authors of papers retain copyright and release the work under a

Creative Commons Attribution 4.0

International License ([CC BY 4.0](#))

## Summary


On the Earth's surface, mixtures are the norm rather than the exception. Taken to the limit, virtually every surface can be split into multiple constituents. Spectral unmixing is the remote sensing retrieval process that attempts to quantitatively determine the relative fractions of various components that make up a surface based on optical data. Imaging spectroscopy, in particular, has demonstrated the capacity for robust fractional retrievals across a wide range of domains, including mineralogical maps (Combe et al., 2008; Yan et al., 2010), urban land cover (Myint & Okin, 2009), and vegetation (Okin et al., 2001).

Spectral unmixing is typically performed under the assumption of linear mixtures. Some set of candidate 'endmembers' - constituents with known absolute (often pure) quantities and spectral signatures - are provided, and each pixel within an image is linearly unmixed with these endmembers to retrieve the relative contribution of each constituent. In essence, this reduces to a simple linear algebra inversion where the known reference library can be inverted and multiplied by the observed reflectances to produce mixture fractions. However, details arise regarding the nature of the selection of endmembers, with strategies ranging from dimensionality reduction of endmember 'classes' (Roberts et al., 1998), bootstrapping (Asner & Lobell, 2000), combinatorial selection (Franke et al., 2009; Roberts et al., 1998), and spectral brightness normalization (Asner & Lobell, 2000). The exact matrix inversion strategy to use is also an open and problem-specific decision, with candidates ranging from direct algebraic inversion to a constrained and regularized optimization (Hastie et al., 2015).

## Statement of need

SpectralUnmixing is a one-stop-shop Julia package for all types of spectral unmixing strategies, focused on imaging spectroscopy data. The code was designed for NASA's Earth Surface Mineral Dust Source Investigation (EMIT) (Green et al., 2020) mission when, during the algorithm design phase, it became evident that there did not yet exist an optimized codebase that provided parallelized and flexible spectral unmixing strategies. In particular, no central framework exists in which different unmixing strategies could be efficiently tested against one another. SpectralUnmixing addresses this issue by drawing on a vast amount of existing literature and bringing the various proposed strategies together into a single codebase. The framework also leverages a rich set of Julia packages for linear algebra, optimization, remote sensing data IO and more, resulting in a highly flexible and scalable unmixing package. SpectralUnmixing has already supported the development of new unmixing strategies by accelerating the process of combining different components of the overall unmixing problem in novel ways (Ochoa et al., n.d.) The package's scalability and breadth will allow it to continue

42 providing this kind of coupled flexibility and operational capacity into the future.



**Figure 1:** Example usage of SpectralUnmixing: fractional cover from an EMIT spectral image. A) Red-green-blue (RGB) image of sample EMIT reflectance data observed near Sacramento, CA, USA with zoom-ins around labeled points (left). B) Fractional cover output of SpectralUnmixing driver script `unmix.jl` on EMIT reflectance image with zoom-ins around labeled points (right). The RGB values in the fractional cover correspond to fractions of endmember library classes: non-photosynthetic vegetation (npv), photosynthetic vegetation (pv), and soil, respectively. C) EMIT reflectance spectra of sample points labeled in A) and B), chosen to each have high fractions of each of the 3 classes. Each spectra is colored by the RGB value corresponding to their class fraction. Here, `unmix.jl` was run using the included example endmember library and with the arguments `--normalization brightness --mode sma-best --n_mc 20 --num_endmembers 30`

43 While SpectralUnmixing was originally created to be used operationally for the EMIT mission,  
44 it was also designed to be quickly adaptable for different researchers' needs. The package  
45 currently supports the industry standard ENVI file format for unmixing raster maps, with  
46 support for more formats planned for the future. The code is fully documented, including  
47 an example notebook, and features a script front-end that allows for arguments to be easily  
48 passed in and different options and datasets to be coupled together for rapid testing. The  
49 package ultimately aims to benefit students, educators, professional researchers, and operational  
50 missions alike.

## 51 Acknowledgements

52 EMIT is supported by the National Aeronautics and Space Administration Earth Venture  
53 Instrument program, USA, under the Earth Science Division of the Science Mission Directorate.  
54 A portion of this research was carried out at the Jet Propulsion Laboratory, California Institute  
55 of Technology, under a contract with the National Aeronautics and Space Administration, USA  
56 (80NM0018D0004). We acknowledge the support and assistance of NASA's International  
57 Space Station Program, USA. Any use of trade, firm, or product names is for descriptive  
58 purposes only and does not imply endorsement by the U.S. Government. Copyright 2024

59 California Institute of Technology. All rights reserved. US Government Support Acknowledged.

## 60 References

- 61 Asner, G. P., & Lobell, D. B. (2000). A biogeophysical approach for automated SWIR  
62 unmixing of soils and vegetation. *Remote Sensing of Environment*, 74(1), 99–112. [https://doi.org/10.1016/S0034-4257\(00\)00126-7](https://doi.org/10.1016/S0034-4257(00)00126-7)  
63
- 64 Combe, J.-P., Le Mouélic, S., Sotin, C., Gendrin, A., Mustard, J., Le Deit, L., Launeau, P.,  
65 Bibring, J.-P., Gondet, B., Langevin, Y., & others. (2008). Analysis of OMEGA/mars  
66 express data hyperspectral data using a multiple-endmember linear spectral unmixing model  
67 (MELSUM): Methodology and first results. *Planetary and Space Science*, 56(7), 951–975.  
68 <https://doi.org/10.1016/j.pss.2007.12.007>
- 69 Franke, J., Roberts, D. A., Halligan, K., & Menz, G. (2009). Hierarchical multiple endmember  
70 spectral mixture analysis (MESMA) of hyperspectral imagery for urban environments.  
71 *Remote Sensing of Environment*, 113(8), 1712–1723. <https://doi.org/10.1016/j.rse.2009.03.018>  
72
- 73 Green, R. O., Mahowald, N., Ung, C., Thompson, D. R., Bator, L., Bennet, M., Bernas, M.,  
74 Blackway, N., Bradley, C., Cha, J., & others. (2020). The earth surface mineral dust  
75 source investigation: An earth science imaging spectroscopy mission. *2020 IEEE Aerospace*  
76 *Conference*, 1–15. <https://doi.org/10.1109/AERO47225.2020.9172731>
- 77 Hastie, T., Tibshirani, R., & Wainwright, M. (2015). Statistical learning with sparsity.  
78 *Monographs on Statistics and Applied Probability*, 143(143), 8. <https://doi.org/10.1201/b18401>  
79
- 80 Myint, S. W., & Okin, G. S. (2009). Modelling land-cover types using multiple endmember  
81 spectral mixture analysis in a desert city. *International Journal of Remote Sensing*, 30(9),  
82 2237–2257. <https://doi.org/10.1080/01431160802549328>
- 83 Ochoa, F., Brodrick, P. G., Okin, G. S., Ben-Dor, E., Meyer, T., Thompson, D. R., & Green,  
84 R. O. (n.d.). Optimal soil and vegetation cover estimation for global imaging spectroscopy  
85 using spectral mixture analysis. *In Review*. <https://doi.org/10.22541/essoar.173193387.79878484/v1>  
86
- 87 Okin, G. S., Roberts, D. A., Murray, B., & Okin, W. J. (2001). Practical limits on hyper-  
88 spectral vegetation discrimination in arid and semiarid environments. *Remote Sensing of*  
89 *Environment*, 77(2), 212–225. [https://doi.org/10.1016/S0034-4257\(01\)00207-3](https://doi.org/10.1016/S0034-4257(01)00207-3)
- 90 Roberts, D. A., Gardner, M., Church, R., Ustin, S., Scheer, G., & Green, R. O. (1998).  
91 Mapping chaparral in the santa monica mountains using multiple endmember spectral  
92 mixture models. *Remote Sensing of Environment*, 65(3), 267–279. [https://doi.org/10.1016/S0034-4257\(98\)00037-6](https://doi.org/10.1016/S0034-4257(98)00037-6)  
93
- 94 Yan, B., Wang, R., Gan, F., & Wang, Z. (2010). Minerals mapping of the lunar surface with  
95 clementine UVVIS/NIR data based on spectra unmixing method and hapke model. *Icarus*,  
96 208(1), 11–19. <https://doi.org/10.1016/j.icarus.2010.01.030>