

cppdlr: Imaginary time calculations using the discrete Lehmann representation

Jason Kaye^{1,2*}, Nils Wentzell^{1*}, and Hugo U. R. Strand^{3,4}

¹ Center for Computational Quantum Physics, Flatiron Institute, New York, NY, USA ² Center for Computational Mathematics, Flatiron Institute, New York, NY, USA ³ School of Science and Technology, Örebro University, Örebro, Sweden ⁴ Institute for Molecules and Materials, Radboud University, 6525 AJ Nijmegen, the Netherlands ¶ Corresponding author * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [✉](#)

Submitted: 22 September 2023

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/))

Summary

Imaginary time Green's functions encode the static and dynamical response of quantum systems at thermal equilibrium to external perturbations, such as applied electromagnetic fields. They therefore represent a direct point of connection between theoretical calculations and experimental measurements. As a consequence, they appear routinely in quantum many-body calculations at finite temperature, both for model systems like the Hubbard model (Hubbard, 1963), and in ab-initio electronic structure calculations beyond density functional theory, e.g., using Hedin's GW method (Golze et al., 2019; Hedin, 1965). Highly compact and accurate representations of imaginary time Green's functions and related imaginary time-dependent response functions are therefore an important ingredient in the development of robust and efficient codes for quantum many-body calculations. However, obtaining such representations has traditionally been challenging, particularly for low temperature calculations, in which the functions develop steep gradients.

In the past several years, significant progress has been achieved using low-rank approximations of the spectral Lehmann representation, which is given by

$$G(\tau) = - \int_{-\infty}^{\infty} d\omega \frac{e^{-\tau\omega}}{1 + e^{-\beta\omega}} \rho(\omega).$$

Here, $G(\tau)$ is a fermionic single-particle imaginary time Green's function, and $\rho(\omega)$ is its corresponding spectral function, which encodes information about the single-particle excitations of the underlying quantum many-body system. The spectral function always exists, but is typically not known. However, the existence of this integral representation constrains the space of possible imaginary time Green's functions to lie within the image of the integral operator, which is numerically low-rank, enabling the construction of highly compact basis representations. The intermediate representation (IR) was introduced first, and used the singular value decomposition to obtain an orthogonal but non-explicit basis of imaginary time Green's functions (Chikano et al., 2018; Shinaoka et al., 2017). The recently-introduced discrete Lehmann representation (DLR) uses the interpolative decomposition to obtain a non-orthogonal basis consisting of known exponential functions (Kaye, Chen, & Parcollet, 2022). The number of basis functions required in both representations is similar, and typically significantly less than the previous state-of-the-art methods based on orthogonal polynomials (Boehnke et al., 2011; Dong et al., 2020; Gull et al., 2018).

The DLR's use of an explicit basis of simple functions makes many common operations, including interpolation, integration, Fourier transform, and convolution, simple and highly efficient. This has led to a variety of recent algorithmic advances, including in reducing the size of the Matsubara frequency mesh in dynamical mean-field theory calculations (Sheng et

al., 2023), stabilizing the calculation of the single-particle self-energy via the Dyson equation (LaBollita et al., 2023), improving the efficiency of the imaginary time discretization in the mixing Green's function of the Keldysh formalism (Kaye & Strand, 2023), and accelerating the evaluation of imaginary time Feynman diagrams (Kaye, Strand, et al., 2023). It has also yielded immediate applications in computational physics, for example in low-temperature studies of superconductivity (Cai et al., 2022; Hou et al., 2023; Tanjaroon Ly et al., 2023). The DLR can be straightforwardly integrated into existing algorithms and codes, often yielding significant improvements in efficiency, accuracy, and algorithmic simplicity.

Statement of need

cppdlr is a C++ library which constructs the DLR and implements its standard operations. The flexible yet high-level interface of cppdlr makes it appealing for use both in small-scale applications and in existing large-scale software projects. The DLR has previously been implemented in other programming languages, specifically in Python via pydlr, in Fortran via libdlr, and in Julia via Lehmann.jl (Chen, 2021; Kaye, Chen, & Strand, 2022; Kaye & Strand, 2021; Strand & Kaye, 2021), as well as in the sparse-ir library implementing the IR (Wallerberger et al., 2023). cppdlr nevertheless provides a needed platform for future developments. First, cppdlr is written in C++, a common language used by many large projects in the quantum many-body physics community. Second, it offers a high-level user interface simpler than that of libdlr, enabled by the use of C++ templating and the nda library ("nda," n.d.) for array types and BLAS/LAPACK compatibility. These features have, for example, enabled the implementation of the DLR in the TRIQS library (Parcollet et al., 2015) for quantum many-body calculations.

cppdlr is distributed under the Apache License Version 2.0 through a public Git repository (Kaye, Wentzell, et al., 2023a). The project documentation (Kaye, Wentzell, et al., 2023b) is extensive, containing background on the DLR, a user guide describing example programs packaged with the library, and application interface (API) reference documentation for all classes and functions. We envision cppdlr as a platform for future algorithmic developments involving the DLR, and as a go-to tool for applications employing the DLR.

Acknowledgements

We are thankful for helpful discussions with Kun Chen, Olivier Parcollet, Malte Rösner, and Yann in 't Veld. H.U.R.S. acknowledges funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 854843-FASTCORR). The Flatiron Institute is a division of the Simons Foundation.

Boehnke, L., Hafermann, H., Ferrero, M., Lechermann, F., & Parcollet, O. (2011). Orthogonal polynomial representation of imaginary-time Green's functions. *Phys. Rev. B*, 84, 075145. <https://doi.org/10.1103/PhysRevB.84.075145>

Cai, X., Wang, T., Prokof'ev, N. V., Svistunov, B. V., & Chen, K. (2022). Superconductivity in the uniform electron gas: Irrelevance of the Kohn-Luttinger mechanism. *Phys. Rev. B*, 106, L220502. <https://doi.org/10.1103/PhysRevB.106.L220502>

Chen, K. (2021). Julia implementation of the discrete lehmann representation (DLR). In *GitHub repository*. GitHub. <https://github.com/numericleft/Lehmann.jl>

Chikano, N., Otsuki, J., & Shinaoka, H. (2018). Performance analysis of a physically constructed orthogonal representation of imaginary-time Green's function. *Phys. Rev. B*, 98, 035104. <https://doi.org/10.1103/PhysRevB.98.035104>

Dong, X., Zgid, D., Gull, E., & Strand, H. U. R. (2020). Legendre-spectral Dyson equation solver with super-exponential convergence. *J. Chem. Phys.*, 152(13), 134107. <https://doi.org/10.1063/1.5138881>

- 87 [//doi.org/10.1063/5.0003145](https://doi.org/10.1063/5.0003145)
- 88 Golze, D., Dvorak, M., & Rinke, P. (2019). The *GW* Compendium: A Practical Guide to
89 Theoretical Photoemission Spectroscopy. *Front. Chem.*, 7, 377. <https://doi.org/10.3389/fchem.2019.00377>
- 90
- 91 Gull, E., Iskakov, S., Krivenko, I., Rusakov, A. A., & Zgid, D. (2018). Chebyshev polynomial
92 representation of imaginary-time response functions. *Phys. Rev. B*, 98, 075127. <https://doi.org/10.1103/PhysRevB.98.075127>
- 93
- 94 Hedin, L. (1965). New Method for Calculating the One-Particle Green's Function with
95 Application to the Electron-Gas Problem. *Phys. Rev.*, 139, A796–A823. <https://doi.org/10.1103/PhysRev.139.A796>
- 96
- 97 Hou, P., Cai, X., Wang, T., Deng, Y., Prokof'ev, N. V., Svistunov, B. V., & Chen, K. (2023).
98 *Precursory Cooper flow in ultralow-temperature superconductors*. <https://arxiv.org/abs/2303.03624>
- 99
- 100 Hubbard, J. (1963). Electron correlations in narrow energy bands. *Proc. R. Soc. Lon. Ser.-A*,
101 276(1365), 238–257.
- 102 Kaye, J., Chen, K., & Parcollet, O. (2022). Discrete Lehmann representation of imaginary
103 time Green's functions. *Phys. Rev. B*, 105, 235115. <https://doi.org/10.1103/PhysRevB.105.235115>
- 104
- 105 Kaye, J., Chen, K., & Strand, H. U. R. (2022). libdlr: Efficient imaginary time calculations
106 using the discrete Lehmann representation. *Comput. Phys. Commun.*, 280, 108458.
107 <https://doi.org/https://doi.org/10.1016/j.cpc.2022.108458>
- 108 Kaye, J., & Strand, H. U. R. (2021). Libdlr: Imaginary time calculations using the discrete
109 lehmann representation (DLR). In *GitHub repository*. GitHub. <https://github.com/jasonkaye/libdlr>
- 110
- 111 Kaye, J., & Strand, H. U. R. (2023). A fast time domain solver for the equilibrium dyson
112 equation. *Adv. Comput. Math.*, 49(4). <https://doi.org/10.1007/s10444-023-10067-7>
- 113 Kaye, J., Strand, H. U. R., & Golež, D. (2023). *Decomposing imaginary time feynman diagrams*
114 *using separable basis functions: Anderson impurity model strong coupling expansion*.
115 <https://arxiv.org/abs/2307.08566>
- 116 Kaye, J., Wentzell, N., & Strand, H. U. R. (2023a). Cppdlr: Imaginary time calculations using
117 the discrete lehmann representation. In *GitHub repository*. GitHub. <https://github.com/flatironinstitute/cppdlr>
- 118
- 119 Kaye, J., Wentzell, N., & Strand, H. U. R. (2023b). Cppdlr: Imaginary time calculations
120 using the discrete lehmann representation. In *GitHub-hosted documentation*. <https://flatironinstitute.github.io/cppdlr/>
- 121
- 122 LaBollita, H., Kaye, J., & Strand, H. U. R. (2023). Stabilizing the calculation of the self-energy
123 in dynamical mean-field theory using constrained residual minimization. In *In preparation*.
- 124 nda: C++ library for multi-dimensional arrays. (n.d.). In *GitHub repository*. <https://github.com/TRIQS/nda>
- 125
- 126 Parcollet, O., Ferrero, M., Ayrat, T., Hafermann, H., Krivenko, I., Messio, L., & Seth, P.
127 (2015). TRIQS: A toolbox for research on interacting quantum systems. *Comput. Phys.*
128 *Commun.*, 196, 398–415. <https://doi.org/http://dx.doi.org/10.1016/j.cpc.2015.04.023>
- 129 Sheng, N., Hampel, A., Beck, S., Parcollet, O., Wentzell, N., Kaye, J., & Chen, K. (2023).
130 Low-rank Green's function representations applied to dynamical mean-field theory. *Phys.*
131 *Rev. B*, 107, 245123. <https://doi.org/10.1103/PhysRevB.107.245123>

- 132 Shinaoka, H., Otsuki, J., Ohzeki, M., & Yoshimi, K. (2017). Compressing Green's function
133 using intermediate representation between imaginary-time and real-frequency domains.
134 *Phys. Rev. B*, 96, 035147. <https://doi.org/10.1103/PhysRevB.96.035147>
- 135 Strand, H. U. R., & Kaye, J. (2021). Pydlr: Imaginary time calculations using the discrete
136 lehmann representation (DLR). In *Python Package Index (PyPI) project*. [https://pypi.org/
137 project/pydlr/](https://pypi.org/project/pydlr/)
- 138 Tanjaroon Ly, A., Cohen-Stead, B., Malkaruge Costa, S., & Johnston, S. (2023). *A comparative
139 study of the superconductivity in the Holstein and optical Su-Schrieffer-Heeger models*.
140 <https://arxiv.org/abs/2307.10809>
- 141 Wallerberger, M., Badr, S., Hoshino, S., Huber, S., Kakizawa, F., Koretsune, T., Nagai, Y.,
142 Nogaki, K., Nomoto, T., Mori, H., Otsuki, J., Ozaki, S., Plaikner, T., Sakurai, R., Vogel,
143 C., Witt, N., Yoshimi, K., & Shinaoka, H. (2023). Sparse-ir: Optimal compression and
144 sparse sampling of many-body propagators. *SoftwareX*, 21, 101266. [https://doi.org/https:
145 //doi.org/10.1016/j.softx.2022.101266](https://doi.org/https://doi.org/10.1016/j.softx.2022.101266)

DRAFT