

# <sup>1</sup> pySnowClim: fast process-based snow modeling in Python

<sup>3</sup> Aranildo R. Lima  <sup>1</sup>, Abby C. Lute  <sup>2</sup>, and Rajesh R. Shrestha  <sup>1</sup>

<sup>4</sup> 1 Climate Research Division, Environment and Climate Change Canada, Victoria, BC, Canada 2

<sup>5</sup> Woodwell Climate Research Center, Falmouth, MA 02540, USA  Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: Jack Atkinson  

Submitted: 08 October 2025

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## <sup>6</sup> Summary

<sup>7</sup> pySnowClim is an open-source Python implementation of the process-based SnowClim model for simulating snow dynamics including accumulation and melt processes. The model achieves a balance between representing important physical processes, such as energy balance calculations, snow density evolution, albedo dynamics, and phase change processes, while simplifying other components to maintain computationally efficiency, delivering consistent snowpack simulations across diverse climatic conditions. These characteristics make pySnowClim ideal for large-scale regional to continental studies, studies requiring < 1km spatial resolution such as those in complex terrain, and studies evaluating the impact of alternate climate scenarios on snowpack.

## <sup>10</sup> Statement of need

<sup>11</sup> Snow is a critical component of the global water cycle. The accurate simulation of snowpack dynamics is not only essential for water resource management, but also for flood forecasting, ecological studies, and climate change impact assessments ([Caretta et al., 2022](#)).

<sup>12</sup> The target audience of pySnowClim includes hydrologists, climatologists, ecologists, water resource managers, and students who need reliable snow modeling capabilities for applications such as:

- <sup>13</sup> **Research:** Detailed energy balance studies and process investigations
- <sup>14</sup> **Operations:** Water resource forecasting and management
- <sup>15</sup> **Education:** Teaching snow physics and energy balance concepts
- <sup>16</sup> **Climate Studies:** Long-term snow evolution under changing conditions
- <sup>17</sup> **Adaptation Planning:** Anticipating and planning adaptations to impacts of warming on snow-dependent species, ecosystems, and communities

<sup>18</sup> In addition, the model can be used on different:

- <sup>19</sup> **Spatial Scales:** Point locations to continental domains
- <sup>20</sup> **Temporal Scales:** Sub-daily to daily timesteps, multi-decadal simulations
- <sup>21</sup> **Environments:** Diverse snow climates from maritime to continental

## <sup>22</sup> State of the field

<sup>23</sup> Many current snow models are either computationally efficient but only represent physical processes to a very limited extent (e.g. temperature index models), or represent many important <sup>24</sup> physical processes but are too computational burdensome for large-scale high-resolution <sup>25</sup> applications (e.g. most process-based models). In addition, many are proprietary, not easily <sup>26</sup> accessible, or difficult to integrate with modern scientific workflows ([Garen & Marks, 2005](#);

<sup>38</sup> Ikeda et al., 2021; Liston & Elder, 2006; Walter et al., 2005; Wrzesien et al., 2018). pySnowClim  
<sup>39</sup> addresses some of these problems by offering a flexible, efficient, and open alternative with a  
<sup>40</sup> good balance between representing physical processes and usability.

## <sup>41</sup> Software design


<sup>42</sup> The model architecture of pySnowClim package follows object-oriented design principles  
<sup>43</sup> with clear separation between forcing data handling, parameter management, core physics  
<sup>44</sup> calculations, and output generation. pySnowClim leverages NumPy's vectorized operations for  
<sup>45</sup> processing multiple grid points simultaneously, enhancing computational efficiency. It also uses  
<sup>46</sup> xarray and pandas for more efficient data manipulation and NetCDF I/O operations. The  
<sup>47</sup> package includes a comprehensive testing framework, featuring unit tests for individual physics  
<sup>48</sup> functions and integration tests for complete workflows. pySnowClim also provides extensive  
<sup>49</sup> documentation (<https://abbylute.github.io/pySnowClim>), including API references, example  
<sup>50</sup> datasets, and validation against observations from a snow monitoring site. Model outputs  
<sup>51</sup> include standard snow variables (SWE, depth, density, melt, albedo, temperature) as well as  
<sup>52</sup> detailed energy budget components.

## <sup>53</sup> Research impact statement

<sup>54</sup> There are several key improvements of pySnowClim compared to the original MATLAB-based  
<sup>55</sup> SnowClim model (Lute et al., 2022). A Python package makes the model more accessible  
<sup>56</sup> to a wider scientific and practical audience. Python is widely adopted in earth science  
<sup>57</sup> and data science communities, and a well-supported package enables easier integration,  
<sup>58</sup> reproducibility, and further development. An example is the current implementation of  
<sup>59</sup> coupling between pySnowClim and the Community Water Model (CWatM) (Burek et al., 2020)  
<sup>60</sup> (<https://github.com/iiasa/CWatM>). In addition, to make the model more realistic, a new  
<sup>61</sup> functionality was added to reduce excessive snow accumulation (i.e. snow towers) using an  
<sup>62</sup> optional radiation enhancement.

## <sup>63</sup> The model

<sup>64</sup> pySnowClim employs the fundamental principles of mass and energy conservation as its core  
<sup>65</sup> framework. The model requires meteorological forcing data including temperature, precipitation,  
<sup>66</sup> shortwave radiation, longwave radiation, wind speed, humidity, and pressure to simulate critical  
<sup>67</sup> snow variables such as snow water equivalent (SWE), snow depth, snow density, snow melt,  
<sup>68</sup> snowpack liquid water content, albedo, and energy fluxes. [Figure 1](#) shows the snow model  
<sup>69</sup> conceptual diagram.



**Figure 1:** Snow model conceptual diagram. Solid black arrows indicate mass fluxes and dashed black arrows indicate energy fluxes.  $T_s$  is snow surface temperature and  $T_{pack}$  is the temperature of the snowpack.

pySnowClim calculates the net energy flux to the snow surface accounting for shortwave and longwave radiation, sensible and latent heat fluxes with stability corrections, ground heat flux, and precipitation heat flux [Equation 1](#). The model is built around the surface energy balance equation:

$$Q_{net} = SW_{down} - SW_{up} + LW_{down} - LW_{up} + H + E_i + E_w + G + P \quad (1)$$

Where  $SW$  denotes shortwave radiation fluxes,  $LW$  denotes longwave radiation fluxes,  $H$  is the sensible heat flux,  $E$  is the latent heat flux of ice ( $i$ ) and water ( $w$ ),  $G$  is the ground heat flux, and  $P$  is the advected heat flux from precipitation.

The model tracks snow accumulation, sublimation and evaporation processes, snowmelt generation, and liquid water movement through the snowpack with mass conservation [Equation 2](#). Mass balance of the solid ( $M_s$ ) and liquid ( $M_l$ ) portions of the snowpack are governed by:

$$M_s = M_{snow} + M_{ref} - M_{melt} + M_{dep} - M_{sub} \quad (2)$$

$$M_l = M_{rain} - M_{ref} + M_{melt} - M_{runoff} + M_{cond} - M_{evap} \quad (3)$$

Where  $M_{snow}$  is the mass of new snowfall,  $M_{ref}$  is the mass of the snowpack liquid water that has been refrozen,  $M_{melt}$  is the mass of snow that has melted,  $M_{dep}$  is the mass of deposition,  $M_{sub}$  is the mass of sublimation,  $M_{rain}$  is the mass of rain added to the snowpack,  $M_{runoff}$  is the mass of liquid water that has left the snowpack as runoff,  $M_{cond}$  is the mass of condensation, and  $M_{evap}$  is the mass of evaporation.

Other key model components include precipitation phase determination using bivariate logistic regression ([Jennings et al., 2018](#)), fresh snow density calculation based on temperature ([Anderson, 1976](#)), snow density evolution through compaction processes, liquid water retention and drainage. pySnowClim has three different albedo scheme implementations including aging

90 effects, melting conditions, and seasonal variations with options for different complexity levels  
91 ([Essery et al., 2013](#); [Hamman et al., 2018](#); [Liang et al., 1994](#); [Tarboton & Luce, 1996](#)).

## 92 AI usage disclosure

93 In preparing this work, the authors used ChatGPT (OpenAI) and Claude (Anthropic) to assist  
94 with the initial MATLAB-to-Python translation and generation of code documentation. The  
95 unit tests of the package were carried out by using Claude. After using these tools, the authors  
96 thoroughly reviewed, edited and refined the content, and take full responsibility for the content  
97 of this publication. No generative AI tools were used in the development of the writing of this  
98 manuscript.

## 99 Acknowledgements

100 We would like to thank Dr. Alex Cannon and Dr. Narayan Shrestha for their constructive  
101 comments and suggestions. In addition, we thank Lincoln Lute for providing the conceptual  
102 diagram of the snow model.

## 103 References

- 104 Anderson, E. A. (1976). *A point energy and mass balance model of a snow cover* (No. 19;  
105 NOAA Technical Report NWS). <https://repository.library.noaa.gov/view/noaa/6392>
- 106 Burek, P., Satoh, Y., Kahil, T., Tang, T., Greve, P., Smilovic, M., Guillaumot, L., Zhao,  
107 F., & Wada, Y. (2020). Development of the community water model (CWatM v1.04)  
108 – a high-resolution hydrological model for global and regional assessment of integrated  
109 water resources management. *Geoscientific Model Development*, 13(7), 3267–3298. <https://doi.org/10.5194/gmd-13-3267-2020>
- 110 Caretta, M. A., Mukherji, A., Arfanuzzaman, M., Betts, R. A., Gelfan, A., Hirabayashi, Y.,  
111 Lissner, T. K., Liu, J., Lopez Gunn, E., Morgan, R., Mwanga, S., & Supratid, S. (2022).  
112 Water. In H.-O. P"ortner, D. C. Roberts, M. Tignor, E. S. Poloczanska, K. Mintenbeck, A.  
113 Alegr'ia, M. Craig, S. Langsdorf, S. L"oschke, V. M"oller, A. Okem, & B. Rama (Eds.),  
114 *Climate change 2022: Impacts, adaptation and vulnerability. Contribution of working group  
115 II to the sixth assessment report of the intergovernmental panel on climate change* (pp.  
116 551–712). Cambridge University Press. <https://doi.org/10.1017/9781009325844.006>
- 117 Essery, R., Morin, S., Lejeune, Y., & B Ménard, C. (2013). A comparison of 1701 snow  
118 models using observations from an alpine site. *Advances in Water Resources*, 55, 131–148.  
119 <https://doi.org/10.1016/j.advwatres.2012.07.013>
- 120 Garen, D. C., & Marks, D. (2005). Spatially distributed energy balance snowmelt modelling in  
121 a mountainous river basin: Estimation of meteorological inputs and verification of model  
122 results. *Journal of Hydrology*, 315, 126–153. <https://doi.org/10.1016/j.jhydrol.2005.03.026>
- 123 Hamman, J. J., Nijssen, B., Bohn, T. J., Gergel, D. R., & Mao, Y. (2018). The variable  
124 infiltration capacity model version 5 (VIC-5): Infrastructure improvements for new  
125 applications and reproducibility. *Geoscientific Model Development*, 11(8), 3481–3496.  
126 <https://doi.org/10.5194/gmd-11-3481-2018>
- 127 Ikeda, K., Rasmussen, R., Liu, C., & al., et. (2021). Snowfall and snowpack in the western u.s.  
128 As captured by convection permitting climate simulations: Current climate and pseudo  
129 global warming future climate. *Climate Dynamics*, 57, 2191–2215. <https://doi.org/10.1007/s00382-021-05805-w>
- 130 Jennings, K. S., Winchell, T. S., & Livneh, B. (2018). Spatial variation of the rain–snow  
131

- 133 temperature threshold across the northern hemisphere. *Nature Communications*, 9, 1148.  
134 <https://doi.org/10.1038/s41467-018-03629-7>
- 135 Liang, X., Lettenmaier, D. P., Wood, E. F., & Burges, S. J. (1994). A simple hydrologically  
136 based model of land surface water and energy fluxes for general circulation models. *Journal*  
137 *of Geophysical Research: Atmospheres*, 99(D7), 14415–14428. <https://doi.org/10.1029/94JD00483>
- 138
- 139 Liston, G. E., & Elder, K. (2006). A distributed snow-evolution modeling system (SnowModel).  
140 *Journal of Hydrometeorology*, 7(6), 1259–1276. <https://doi.org/10.1175/JHM548.1>
- 141 Lute, A. C., Abatzoglou, J., & Link, T. (2022). SnowClim v1.0: High-resolution snow  
142 model and data for the western United States. *Geoscientific Model Development*, 15(13),  
143 5045–5071. <https://doi.org/10.5194/gmd-15-5045-2022>
- 144 Tarboton, D. G., & Luce, C. H. (1996). *Utah energy balance snow accumulation and*  
145 *melt model (UEB): Computer model technical description and user guide*. Utah Water  
146 Research Laboratory; USDA Forest Service Rocky Mountain Research Station. <https://hydrology.usu.edu/dtarb/snow/snowrep.pdf>
- 147
- 148 Walter, M. T., Brooks, E. S., McCool, D. K., King, L. G., Molnau, M., & Boll, J. (2005).  
149 Process-based snowmelt modeling: Does it require more input data than temperature-index  
150 modeling? *Journal of Hydrology*, 300(1-4), 65–75. <https://doi.org/10.1016/j.jhydrol.2004.05.002>
- 151
- 152 Wrzesien, M. L., Durand, M. T., Pavelsky, T. M., Kapnick, S. B., Zhang, Y., Guo, J., &  
153 Shum, C. K. (2018). A new estimate of north american mountain snow accumulation  
154 from regional climate model simulations. *Geophysical Research Letters*, 45, 1423–1432.  
155 <https://doi.org/10.1002/2017GL076664>