

Sunny.jl: A Julia Package for Spin Dynamics*

David Dahlbom^{1,2¶}, Hao Zhang³, Cole Miles⁴, Sam Quinn^{5,6}, Alin Niraula⁷, Bhushan Thipe⁷, Matthew Wilson⁸, Sakib Matin³, Het Mankad⁹, Steven Hahn⁹, Daniel Pajerowski¹, Steve Johnston^{2,10}, Zhentao Wang¹¹, Harry Lane^{12,13,14}, Ying Wai Li¹⁵, Xiaojian Bai⁷, Martin Mourigal⁵, Cristian D. Batista², and Kipton Barros^{3¶}

¹ Neutron Scattering Division, Oak Ridge National Laboratory ² Department of Physics and Astronomy, University of Tennessee ³ Theoretical Division and CNLS, Los Alamos National Laboratory ⁴ Kodiak Robotics ⁵ School of Physics, Georgia Institute of Technology ⁶ Department of Physics and Astronomy, University of California, Los Angeles ⁷ Department of Physics and Astronomy, Louisiana State University ⁸ X-Computational Physics Division, Los Alamos National Laboratory ⁹ Computer Science and Mathematics Division, Oak Ridge National Laboratory ¹⁰ Institute for Advanced Materials and Manufacturing, University of Tennessee ¹¹ Center for Correlated Matter and School of Physics, Zhejiang University ¹² Department of Physics and Astronomy, University of Manchester ¹³ The University of Manchester at Harwell, University of Manchester ¹⁴ School of Physics and Astronomy, University of St Andrews ¹⁵ Computer, Computational, and Statistical Sciences Division, Los Alamos National Laboratory ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: Sophie Beck ↗ 

Reviewers:

- [@jabirali](#)
- [@Chronum94](#)

Submitted: 23 December 2024

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Summary

Sunny is a Julia package designed to serve the needs of the quantum magnetism community. It supports the specification of a very broad class of spin models and a diverse suite of numerical solvers. These include powerful methods for simulating spin dynamics both in and out of equilibrium. Uniquely, it features a broad generalization of classical and semiclassical approaches to $SU(N)$ coherent states, which is useful for studying systems exhibiting strong spin-orbit coupling or local entanglement effects. Sunny also offers a well-developed framework for calculating the dynamical spin structure factor, enabling direct comparison with scattering experiments. Ease of use is a priority, with tools for symmetry-guided modeling and interactive visualization.

Statement of need

Progress in quantum magnetism depends on the development of accurate models of magnetic materials. Scattering techniques, such as inelastic neutron scattering (INS) and resonant inelastic X-ray scattering (RIXS), are among the most informative methods available for probing the dynamics of quantum magnets, yielding the dynamical spin structure factor $\mathcal{S}(\mathbf{q}, \omega)$ as experimental output. To evaluate the validity of a hypothetical model, it is necessary to calculate $\mathcal{S}(\mathbf{q}, \omega)$ theoretically. This is generally an intractable problem that must be treated numerically or with various approximation schemes. The difficulty of this step represents a bottleneck in the development of accurate models and impedes the advancement of our understanding of quantum materials.

*This manuscript has been authored by UT-Battelle, LLC, under contract DE-AC05-00OR22725 with the US Department of Energy (DOE). The US government retains and the publisher, by accepting the article for publication, acknowledges that the US government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for US government purposes. DOE will provide public access to these results of federally sponsored research in accordance with the DOE Public Access Plan (<https://www.energy.gov/doe-public-access-plan>).

The Sunny project is a collaborative effort among theorists, experimentalists and computational scientists aimed at developing theoretical and numerical methodologies for modeling realistic quantum magnets. The central product of this effort is the Sunny software package, which makes recent theoretical advances available in a form readily accessible to students and researchers. Distinguishing features of Sunny include:

- Symmetry analysis tools that facilitate model specification, visualization and data retrieval.
- A suite of optimizers, Monte Carlo samplers, and spin dynamics solvers that can all be applied to the same system specification.
- Implementation of the $SU(N)$ coherent state formalism for classical and semiclassical calculations.
- An interface tailored toward the needs of scattering scientists, with tools for integrating scattering intensities over regions of reciprocal space.
- Code written entirely in Julia, a language that can achieve speeds comparable to C++ or Fortran while offering an interactive workflow that will be familiar to users of Python and Matlab.
- A well documented codebase, an extensive collection of correctness tests, a website featuring many tutorials, and an active Slack channel where users can ask questions.

There are a number of existing codes that can calculate $\mathcal{S}(\mathbf{q}, \omega)$ using linear spin wave theory (LSWT), some of which have served as inspiration to the Sunny project [Rotter (2004); (2024a); Petit & Damay (2016); weber:2016; (2024b)]. The symmetry analysis tools of SpinW in particular have served as a model (Toth & Lake, 2015). There are also codes that perform classical spin simulations using Landau-Lifshitz (LL) dynamics (2024c; Evans et al., 2014; Müller et al., 2019). Sunny is unique in offering both approaches and generalizing them through a formalism based on $SU(N)$ coherent states (Muniz et al., 2014; H. Zhang & Batista, 2021). Sunny additionally permits completely general single-ion anisotropies and coupling of multipolar moments; provides an efficient implementation of long-range dipole-dipole interactions; automates the application of a number of quantum renormalizations (Dahlbom et al., 2023); and offers iterative solvers for efficient LSWT on large magnetic cells (Lane et al., 2024).

The value of collecting all these tools together in a modern, easy-to-use package is evidenced by the large number of publications that have already made use of Sunny, a partial list of which is maintained on the GitHub wiki (2024d). We note a number of experimental studies that have relied on Sunny for analysis (Bai et al., 2021, 2023; Do et al., 2023; Kim et al., 2023; Lee et al., 2023; Na et al., 2024; Nagl et al., 2024; Paddison et al., 2024; Park et al., 2023; Park, Sala, et al., 2024; Park, Ghioldi, et al., 2024; Park, Cho, et al., 2024; Scheie et al., 2023); as well as theoretical and methodological works (Dahlbom, Brooks, et al., 2024; Dahlbom, Thomas, et al., 2024; H. Zhang et al., 2023; Hao Zhang & Lin, 2024). Additional papers documenting the theoretical and algorithmic advances that have enabled the development of Sunny are discussed below.

Feature Overview

Symmetry analysis

By unifying and extending existing open source frameworks for the symmetry analysis of crystals – including Spglib (Togo et al., 2024), Brillouin.jl (2024e), and CrystalInfoFramework.jl (2024f) – Sunny facilitates the process of determining the complete set of interactions allowed by spacegroup symmetries. Similarly, any interaction specified on a site or bond will be automatically propagated to all symmetry-equivalent sites and bonds, as required by the spacegroup symmetries. Models may also be specified according to symmetry properties and subsequently made “inhomogenous,” allowing the arbitrary modification of pair interactions and site properties without regard to symmetry constraints. This greatly facilitates the modeling of

88 systems exhibiting chemical disorder. Finally, the symmetry information enables convenient
89 specification of paths and slices through reciprocal space, aiding visualization and comparison
90 to experimental data. All these tools can be applied just as easily to a user-specified crystal or
91 to a crystal loaded from an industry-standard CIF file (Hall et al., 1991).




Figure 1: a) Ground state of FeI₂, found using Sunny's minimize_energy! function and visualized with plot_spins. b) The crystal of FeI₂ visualized with the view_crystal function. Hovering the cursor over a bond reveals the exchange interaction, if already assigned, or a general expression for all symmetry-allowed interactions.

Visualization

Both the symmetry analysis and data retrieval features of Sunny include 3D visualization tools built on the Makie package (Danisch & Krumbiegel, 2021). These can be used to plot spin configurations, investigate the symmetries of a crystal Figure 1, generate animations of dynamic behavior, and plot the predicted results of scattering experiments Figure 2.

SU(N) Formalism and System Modes

Traditional classical and semiclassical approaches to spin dynamics are based on the assignment of a classical dipole to each lattice site. Recent theoretical work has generalized this picture, replacing dipoles with richer objects, namely SU(N) coherent states. Such states capture the full structure of an N-level quantum system. Setting $N = 2s + 1$ enables the faithful representation of a quantum spin-s and of the crystal field levels of a single-ion. The formalism can also be adapted to model local entanglement effects, where this entanglement may be between the spin and orbital degrees of freedom on a single site or within a cluster of spins on different sites.

The SU(N) formalism applies equally to LSWT calculations (Muniz et al., 2014) and classical spin dynamics (H. Zhang & Batista, 2021). Users can access this framework simply by setting the "mode" of a spin system to :SUN. Sunny also offers a :dipole mode, which is similar to the traditional classical approach but includes quantum renormalizations of biquadratic and single-ion anisotropy terms (Dahlbom et al., 2023). Finally, there is a mode that implements the traditional approach without any additional corrections, :dipole_uncorrected. Most Sunny features are supported in all modes.




Figure 2: Left: Scattering intensities of FeI_2 as measured on the SEQUOIA instrument at the Spallation Neutron Source, Oak Ridge National Laboratory (Bai et al., 2021). Right: Predicted scattering intensities calculated with Sunny's $\text{SU}(N)$ linear spin wave solver. The figure was generated with Sunny's data retrieval and plotting functions.

Optimization and Monte Carlo Tools

Identifying a classical ground state is often the first step when calculating the scattering response of a magnet. Sunny provides several tools for finding such states, including gradient optimizers built on the Optim.jl package (Mogensen & Riseth, 2018). These optimizers work on supercells as well as spiral orderings. Sophisticated Monte Carlo tools are also provided, which can be used both to anneal into ground states and to estimate finite temperature statistics. In particular, the classical dynamics can be run with Langevin coupling to a thermal bath (Dahlbom, Miles, et al., 2022), and samplers are provided that implement the Wang-Landau (Wang & Landau, 2001) and parallel tempering algorithms (Swendsen & Wang, 1986).

Linear Spin Wave Theory (LSWT)

Sunny has extensive support for LSWT calculations, including for systems with arbitrarily complex single-ion anisotropies, general bilinear and biquadratic interactions, and long-range dipole-dipole interactions. Like SpinW, Sunny provides efficient LSWT calculations for systems that exhibit incommensurate spiral orderings (Toth & Lake, 2015). Additionally, Sunny provides tools to efficiently calculate $\mathcal{S}(\mathbf{q}, \omega)$ on very large magnetic cells using iterative matrix-vector multiplications (Lane et al., 2024). The simulation of large supercells is essential to study systems with chemical disorder or complex magnetic orderings.

Classical Dynamics

The efficiency of LSWT calculations makes it the preferred tool when studying magnets near zero temperature. At elevated temperatures or in out-of-equilibrium conditions, however, classical dynamics becomes a valuable technique. Sunny supports both traditional Landau-Lifshitz dynamics and its generalization to $\text{SU}(N)$ coherent states (H. Zhang & Batista, 2021). Dissipationless trajectories are calculated using a symplectic integration scheme (Dahlbom, Zhang, et al., 2022), and a generalization of the stochastic Landau-Lifshitz-Gilbert equations to $\text{SU}(N)$ coherent states (Dahlbom, Miles, et al., 2022) enables the simulation of dynamics coupled to a thermal bath. This is particularly valuable for simulating, e.g., thermal transport, pump-probe experiments, and spin-glass relaxation.

Sunny as a Platform for Future Developments

To make these existing features more widely available, work at ORNL is underway to integrate Sunny into the Calvera platform for neutron data analysis (Watson et al., 2022). Sunny

143 itself can serve as a platform for new solvers and analysis techniques, building on its mature
 144 model specification and data retrieval features. Current efforts are directed at supporting: the
 145 self-consistent Gaussian approximation for diffuse scattering, enabling functionality inspired by
 146 (Paddison, 2023); the modeling of local entanglement effects generated by spin-orbit coupling
 147 or strongly coupled clusters of spins; non-perturbative corrections to LSWT for the modeling of
 148 continua and bound states, which can be probed in INS and terahertz spectroscopy experiments
 149 (Bai et al., 2023; Legros et al., 2021); and observables relevant to RIXS experiments.

150 Acknowledgements

151 We thank Mosé Giordano and Simon Danisch for valuable discussions. This work was supported
 152 by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under
 153 Award Numbers DE-SC0022311, DE-SC-0018660, and DE-SC0025426. The work was also
 154 sponsored by the Laboratory Directed Research and Development Programs (LDRD) at Los
 155 Alamos National Laboratory, managed by Triad National Security, LLC, and at Oak Ridge
 156 National Laboratory, managed by UT-Battelle, LLC, for the U. S. Department of Energy.
 157 Applications to inverse scattering were partially supported by the National Science Foundation
 158 Materials Research Science and Engineering Center program through the UT Knoxville Center for
 159 Advanced Materials and Manufacturing (DMR-2309083). Z.W. acknowledges support from the
 160 National Key Research and Development Program of China (Grant No. 2024YFA1408303) and
 161 the National Natural Science Foundation of China (Grant No. 12374124). H.L. acknowledges
 162 funding from the Royal Commission for the Exhibition of 1851. The data shown in Figure 2
 163 was collected at the the Spallation Neutron Source, a DOE Office of Science User Facility
 164 operated by the Oak Ridge National Laboratory. Beam time was allocated to the SEQUOIA
 165 instrument on proposal number IPTS-21166.

166 References

- 167 (2024d). In *GitHub repository*. GitHub. <https://github.com/SunnySuite/Sunny.jl>
- 168 (2024a). In *GitHub repository*. GitHub. <https://github.com/SpinWaveGenie/SpinWaveGenie>
- 169 (2024c). In *GitHub*. GitHub. <https://github.com/UppASD/UppASD>
- 170 (2024b). In *GitHub repository*. GitHub. <https://github.com/bingli621/pyLiSW>
- 171 (2024f). In *GitHub repository*. GitHub. <https://github.com/jamesrhester/CrystalInfoFramework.jl>
- 172
- 173 (2024e). In *GitHub repository*. GitHub. <https://github.com/thchr/Brillouin.jl>
- 174 Bai, X., Zhang, S.-S., Dun, Z., Zhang, H., Huang, Q., Zhou, H., Stone, M. B., Kolesnikov,
 175 A. I., Ye, F., Batista, C. D., & others. (2021). Hybridized quadrupolar excitations in
 176 the spin-anisotropic frustrated magnet FeI₂. *Nature Physics*, 17(4), 467–472. <https://doi.org/10.1038/s41567-020-01110-1>
- 177
- 178 Bai, X., Zhang, S.-S., Zhang, H., Dun, Z., Phelan, W. A., Garlea, V. O., Mourigal, M., &
 179 Batista, C. D. (2023). Instabilities of heavy magnons in an anisotropic magnet. *Nature*
 180 *Communications*, 14(1), 4199. <https://doi.org/10.1038/s41467-023-39940-1>
- 181 Dahlbom, D. A., Brooks, F. T., Wilson, M. S., Chi, S., Kolesnikov, A. I., Stone, M. B., Cao, H.,
 182 Li, Y. W., Barros, K., Mourigal, M., & others. (2024). Quantum-to-classical crossover in
 183 generalized spin systems: Temperature-dependent spin dynamics of FeI₂. *Physical Review*
 184 *B*, 109(1), 014427. <https://doi.org/10.1103/PhysRevB.109.014427>
- 185 Dahlbom, D. A., Miles, C., Zhang, H., Batista, C. D., & Barros, K. (2022). Langevin dynamics
 186 of generalized spins as SU(N) coherent states. *Physical Review B*, 106(23), 235154.
 187 <https://doi.org/10.1103/PhysRevB.106.235154>

- 188 Dahlbom, D. A., Thomas, J., Johnston, S., Barros, K., & Batista, C. D. (2024). Classical
189 dynamics of the antiferromagnetic Heisenberg $S=1/2$ spin ladder. *Physical Review B*,
190 110(10), 104403. <https://doi.org/10.1103/PhysRevB.110.104403>
- 191 Dahlbom, D. A., Zhang, H., Laraib, Z., Pajeroski, D. M., Barros, K., & Batista, C. D. (2023).
192 Renormalized classical theory of quantum magnets. *arXiv Preprint arXiv:2304.03874*.
193 <https://doi.org/10.48550/arXiv.2304.03874>
- 194 Dahlbom, D. A., Zhang, H., Miles, C., Bai, X., Batista, C. D., & Barros, K. (2022). Geometric
195 integration of classical spin dynamics via a mean-field Schrödinger equation. *Physical*
196 *Review B*, 106(5), 054423. <https://doi.org/10.1103/PhysRevB.106.054423>
- 197 Danisch, S., & Krumbiegel, J. (2021). Makie.jl: Flexible high-performance data visualization for
198 Julia. *Journal of Open Source Software*, 6(65), 3349. <https://doi.org/10.21105/joss.03349>
- 199 Do, S.-H., Zhang, H., Dahlbom, D. A., Williams, T. J., Garlea, V. O., Hong, T., Jang, T.-H.,
200 Cheong, S.-W., Park, J.-H., Barros, K., & others. (2023). Understanding temperature-
201 dependent $SU(3)$ spin dynamics in the $S=1$ antiferromagnet $Ba_2FeSi_2O_7$. *Npj Quantum*
202 *Materials*, 8(1), 5. <https://doi.org/10.1038/s41535-022-00526-7>
- 203 Evans, R. F. L., Fan, W. J., Churemart, P., Ostler, T. A., Ellis, M. O. A., & Chantrell, R. W.
204 (2014). Atomistic spin model simulations of magnetic nanomaterials. *Journal of Physics:*
205 *Condensed Matter*, 26(10), 103202. <https://doi.org/10.1088/0953-8984/26/10/103202>
- 206 Hall, S. R., Allen, F. H., & Brown, I. D. (1991). The crystallographic information file (CIF):
207 A new standard archive file for crystallography. *Foundations of Crystallography*, 47(6),
208 655–685. <https://doi.org/10.1107/S010876739101067X>
- 209 Kim, C., Kim, S., Park, P., Kim, T., Jeong, J., Ohira-Kawamura, S., Murai, N., Nakajima,
210 K., Chernyshev, A. L., Mourigal, M., & others. (2023). Bond-dependent anisotropy and
211 magnon decay in cobalt-based Kitaev triangular antiferromagnet. *Nature Physics*, 19(11),
212 1624–1629. <https://doi.org/10.1038/s41567-023-02180-7>
- 213 Lane, H., Zhang, H., Dahlbom, D., Quinn, S., Somma, R., Mourigal, M., Batista, C. D., &
214 Barros, K. (2024). Kernel polynomial method for linear spin wave theory. *SciPost Physics*,
215 17(5), 145. <https://doi.org/10.21468/SciPostPhys.17.5.145>
- 216 Lee, M., Schönemann, R., Zhang, H., Dahlbom, D., Jang, T.-H., Do, S.-H., Christianson, A.
217 D., Cheong, S.-W., Park, J.-H., Brosha, E., & others. (2023). Field-induced spin level
218 crossings within a quasi-XY antiferromagnetic state in $Ba_2FeSi_2O_7$. *Physical Review B*,
219 107(14), 144427. <https://doi.org/10.1103/PhysRevB.107.144427>
- 220 Legros, A., Zhang, S.-S., Bai, X., Zhang, H., Dun, Z., Phelan, W. A., Batista, C. D., Mourigal,
221 M., & Armitage, N. (2021). Observation of 4-and 6-magnon bound states in the spin-
222 anisotropic frustrated antiferromagnet FeI_2 . *Physical Review Letters*, 127(26), 267201.
223 <https://doi.org/10.1103/PhysRevLett.127.267201>
- 224 Mogensen, P., & Riseth, A. (2018). Optim: A mathematical optimization package for julia.
225 *Journal of Open Source Software*, 3(24). <https://doi.org/10.21105/joss.00615>
- 226 Müller, G. P., Hoffmann, M., DiBelknap, C., Schürhoff, D., Mavros, S., Sallermann, M.,
227 Kiselev, N. S., Jónsson, H., & Blügel, S. (2019). Spirit: Multifunctional framework for
228 atomistic spin simulations. *Physical Review B*, 99(22), 224414. <https://doi.org/10.1103/PhysRevB.99.224414>
- 229
- 230 Muniz, R. A., Kato, Y., & Batista, C. D. (2014). Generalized spin-wave theory: Application to
231 the bilinear–biquadratic model. *Progress of Theoretical and Experimental Physics*, 2014(8),
232 083I01. <https://doi.org/10.1093/ptep/ptu109>
- 233 Na, W., Park, P., Oh, S., Kim, J., Scheie, A., Tennant, D. A., Lee, H. C., Park, J.-G.,
234 & Cheong, H. (2024). Direct observation and analysis of low-energy magnons with
235 Raman spectroscopy in atomically thin $NiPS_3$. *ACS Nano*, 18(31), 20482–20492. <https://doi.org/10.1021/acsnano.3c05000>

- 236 [//doi.org/10.1021/acsnano.4c04824](https://doi.org/10.1021/acsnano.4c04824)
- 237 Nagl, J., Flavián, D., Hayashida, S., Povarov, K. Y., Yan, M., Murai, N., Ohira-Kawamura, S.,
238 Simutis, G., Hicken, T. J., Luetkens, H., & others. (2024). Excitation spectrum and spin
239 Hamiltonian of the frustrated quantum Ising magnet Pr_3BWO_9 . *Physical Review Research*,
240 6(2), 023267. <https://doi.org/10.1103/PhysRevResearch.6.023267>
- 241 Paddison, J. A. M. (2023). Spinteract: A program to refine magnetic interactions to diffuse
242 scattering data. *Journal of Physics: Condensed Matter*, 35(49), 495802. <https://doi.org/10.1088/1361-648X/acf261>
- 243
- 244 Paddison, J. A. M., Zhang, H., Yan, J., Cliffe, M. J., McGuire, M. A., Do, S.-H., Gao, S.,
245 Stone, M. B., Dahlbom, D. A., Barros, K., & others. (2024). Cubic double perovskites
246 host noncoplanar spin textures. *Npj Quantum Materials*, 9(1), 48. <https://doi.org/10.1038/s41535-024-00650-6>
- 247
- 248 Park, P., Cho, W., Kim, C., An, Y., Iida, K., Kajimoto, R., Matin, S., Zhang, S.-S., Batista,
249 C. D., & Park, J.-G. (2024). Contrasting dynamical properties of single-Q and triple-Q
250 magnetic orderings in a triangular lattice antiferromagnet. *arXiv Preprint arXiv:2410.02180*.
251 <https://doi.org/10.48550/arXiv.2410.02180>
- 252 Park, P., Cho, W., Kim, C., An, Y., Kang, Y.-G., Avdeev, M., Sibille, R., Iida, K., Kajimoto,
253 R., Lee, K. H., & others. (2023). Tetrahedral triple-Q magnetic ordering and large
254 spontaneous Hall conductivity in the metallic triangular antiferromagnet $\text{Co}_{1/3}\text{TaS}_2$. *Nature*
255 *Communications*, 14(1), 8346. <https://doi.org/10.1038/s41467-023-43853-4>
- 256 Park, P., Ghioldi, E. A., May, A. F., Kolopus, J. A., Podlesnyak, A. A., Calder, S., Paddison,
257 J. A. M., Trumper, A. E., Manuel, L. O., Batista, C. D., & others. (2024). Anomalous
258 continuum scattering and higher-order van Hove singularity in the strongly anisotropic
259 $S=1/2$ triangular lattice antiferromagnet. *Nature Communications*, 15(1), 7264. <https://doi.org/10.1103/PhysRevResearch.6.033184>
- 260
- 261 Park, P., Sala, G., Pajerowski, D. M., May, A. F., Kolopus, J. A., Dahlbom, D. A., Stone, M.
262 B., Halász, G. B., & Christianson, A. D. (2024). Quantum and classical spin dynamics
263 across temperature scales in the $S=1/2$ Heisenberg antiferromagnet. *Physical Review*
264 *Research*, 6(3), 033184. <https://doi.org/10.1103/PhysRevResearch.6.033184>
- 265 Petit, S., & Damay, F. (2016). SpinWave, a software dedicated to spin wave simulations.
266 *Neutron News*, 27(4), 27–28. <https://doi.org/10.1080/10448632.2016.1233020>
- 267 Rotter, M. (2004). Using McPhase to calculate magnetic phase diagrams of rare earth
268 compounds. *Journal of Magnetism and Magnetic Materials*, 272, E481–E482. <https://doi.org/10.1016/j.jmmm.2003.12.1394>
- 269
- 270 Scheie, A., Park, P., Villanova, J. W., Granroth, G. E., Sarkis, C. L., Zhang, H., Stone, M.
271 B., Park, J.-G., Okamoto, S., Berlijn, T., & others. (2023). Spin wave Hamiltonian and
272 anomalous scattering in NiPS_3 . *Physical Review B*, 108(10), 104402. <https://doi.org/10.1103/PhysRevB.108.104402>
- 273
- 274 Swendsen, R. H., & Wang, J.-S. (1986). Replica Monte Carlo simulation of spin-glasses.
275 *Physical Review Letters*, 57(21), 2607. <https://doi.org/10.1103/PhysRevLett.57.2607>
- 276 Togo, A., Shinohara, K., & Tanaka, I. (2024). Spglib: A software library for crystal symmetry
277 search. *Science and Technology of Advanced Materials: Methods*, 4(1), 2384822. <https://doi.org/10.1080/27660400.2024.2384822>
- 278
- 279 Toth, S., & Lake, B. (2015). Linear spin wave theory for single-Q incommensurate magnetic
280 structures. *Journal of Physics: Condensed Matter*, 27(16), 166002. <https://doi.org/10.1088/0953-8984/27/16/166002>
- 281
- 282 Wang, F., & Landau, D. P. (2001). Efficient, multiple-range random walk algorithm to
283 calculate the density of states. *Physical Review Letters*, 86(10), 2050. <https://doi.org/10.1103/PhysRevLett.86.2050>

- 284 [1103/PhysRevLett.86.2050](#)
- 285 Watson, G. R., Cage, G., Fortney, J., Granroth, G. E., Hughes, H., Maier, T., McDonnell,
286 M., Ramirez-Cuesta, A., Smith, R., Yakubov, S., & others. (2022). Calvera: A plat-
287 form for the interpretation and analysis of neutron scattering data. *Smoky Mountains*
288 *Computational Sciences and Engineering Conference*, 137–154. [https://doi.org/10.1007/](https://doi.org/10.1007/978-3-031-23606-8_9)
289 [978-3-031-23606-8_9](https://doi.org/10.1007/978-3-031-23606-8_9)
- 290 Zhang, H., & Batista, C. D. (2021). Classical spin dynamics based on SU(N) coherent states.
291 *Physical Review B*, 104(10), 104409. <https://doi.org/10.1103/PhysRevB.104.104409>
- 292 Zhang, Hao, & Lin, S.-Z. (2024). Multipolar skyrmion crystals in non-Kramers doublet systems.
293 *Physical Review Letters*, 133(19), 196702. <https://doi.org/PhysRevLett.133.196702>
- 294 Zhang, H., Wang, Z., Dahlbom, D. A., Barros, K., & Batista, C. D. (2023). CP² skyrmions
295 and skyrmion crystals in realistic quantum magnets. *Nature Communications*, 14(1), 3626.
296 <https://doi.org/10.1038/s41467-023-39232-8>

DRAFT