

PolSAR tools: A QGIS plugin for generating SAR descriptors

Narayanarao Bhogapurapu¹, Subhadip Dey¹, Dipankar Mandal¹, Avik Bhattacharya¹, and Y. S. Rao¹

¹ Microwave Remote Sensing Lab, Centre of Studies in Resources Engineering, Indian Institute of Technology Bombay, Mumbai-400076, India

DOI: [10.21105/joss.02970](https://doi.org/10.21105/joss.02970)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Hugo Ledoux](#) ↗

Reviewers:

- [@liberostelios](#)
- [@HenrikJanPersson](#)

Submitted: 28 December 2020

Published: 08 April 2021

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Statement of need

The demand for processing tools increases with the increasing number of **Synthetic Aperture Radar (SAR)** satellite missions and datasets. However, to process SAR data, a minimal number of free tools are available ([PolSARpro](#), [SNAP](#)), which consolidates all necessary pre-processing steps. Bearing this in mind, there is a need to develop specific tools for the remote sensing user community to derive polarimetric descriptors like the vegetation indices and decomposition parameters. Besides, to the best of our knowledge, there are no such free tools available on a GIS platform, which is often necessary for the processing of SAR remote sensing datasets.

Hence we have developed a plugin for QGIS that supports data for the three available polarimetric modes (i.e., full-, compact, and dual). The PolSAR tools plugin generates polarimetric descriptors (viz., vegetation indices, polarimetric decomposition parameters) from the 3x3 (C3/T3) or the 2x2 (C2/T2) covariance (coherency) matrices obtained from the ESA's [PolSARpro](#) software. The input data needs to be in PolSARpro format (*.bin and *.hdr). The plugin is coded in Python and is dependant on the QGIS framework. It uses the following Python libraries (bundled with QGIS): [numpy](#), and [gdal](#).

Background

Conventional model-based decomposition methods utilize diverse scattering models and typical hierarchical rule to enumerate power components leading to numerous limitations. The **polarimetric decomposition techniques** incorporated in this QGIS plugin are model-free, i.e., no prior scattering models are assumed to compute the powers. The proposed decomposition techniques utilize certain novel roll-invariant target characterization parameters to decompose the total power into even bounce, odd bounce, and diffused power components. It is guaranteed that the proposed technique's powers are non-negative, which is seldom true with the existing methodologies.

In terms of target descriptors, we often use **vegetation indices** as plant growth proxies. While appreciating the potential of vegetation indices derived from optical remote sensing sensors, regional to global products have been supported for operational uses. These days, the Earth Observation (EO) community relies on SAR imaging technology due to its all-weather imaging capability among its numerous advantages. The SAR images are presently processed by several downstream users and are more frequently interpreted by non-radar specialists. This paradigm shift allows the utility of radar-derived vegetation indices towards a quintessential goal of Analysis Ready Data (ARD) products.

Recently, we proposed three vegetation indices: GRVI (Generalized Radar Vegetation Index) ([Ratha et al., 2019](#)), CpRVI (Compact-pol Radar Vegetation Index) ([Mandal, Ratha, et al., 2020](#)), and Dual-pol Radar Vegetation Index (DpRVI) ([Mandal, Kumar, et al., 2020](#)) for distinct acquisition modes. These vegetation indices have provided a better opportunity to estimate biophysical parameters with fitted models directly. The retrieval of biophysical parameters from SAR observations is of vital importance for in-season monitoring of crop growth.

PoISAR tools Audience

PoISAR tools are intended for students, researchers, and polarimetry experts to derive different SAR descriptors, utilizing the QGIS and Python ecosystem of diverse tools. Especially for non-domain and application users, the plugin interface provides an easy way to process the pre-processed *polarimetric SAR data*.

PoISAR tools Functionality

The functionalities of the PoISAR tools are organized into three modules to handle the data from three different SAR polarization modes. The following is the list of the available functions in the PoISAR tools:

- Full-pol :
 - Radar Vegetation Index (RVI) ([Kim & Zyl, 2009](#))
 - Generalized volume Radar Vegetation Index (GRVI) ([Ratha et al., 2019](#))
 - Polarimetric Radar Vegetation Index (PRVI) ([Chang et al., 2018](#))
 - Model free 3-Component decomposition for full-pol data (MF3CF) ([Dey et al., 2020](#))
 - Model free 4-Component decomposition for full-pol data (MF4CF) ([Dey et al., 2021](#))
 - Degree of Polarization (DOP) ([Barakat, 1977](#))
- Compact-pol :
 - Model free 3-Component decomposition for compact-pol data (MF3CC) ([Dey et al., 2020](#))
 - Improved S-Omega decomposition for compact-pol data (iS-Omega) ([Kumar et al., 2020](#))
 - Compact-pol Radar Vegetation Index (CpRVI) ([Mandal, Ratha, et al., 2020](#))
 - Degree of Polarization (DOP)
- Dual-pol :
 - Radar Vegetation Index (RVI)
 - Dual-pol Radar Vegetation Index (DpRVI) ([Mandal, Kumar, et al., 2020](#)),
 - Polarimetric Radar Vegetation Index (PRVI)
 - Degree of Polarization (DOP) ([Barakat, 1977](#))

Acknowledgements

The authors would like to thank the developers of [QGIS Plugin Builder](#). Authors acknowledge the [GEO-AWS Earth Observation Cloud Credits Program](#), which supported the computation, development, and testing of PoISAR tools on AWS cloud platform through the project: *AWS4AgriSAR-Crop inventory mapping from SAR data on cloud computing platform*.

References

- Barakat, R. (1977). Degree of polarization and the principal idempotents of the coherency matrix. *Optics Communications*, 23(2), 147–150. [https://doi.org/10.1016/0030-4018\(77\)90292-9](https://doi.org/10.1016/0030-4018(77)90292-9)
- Chang, J. G., Shoshany, M., & Oh, Y. (2018). Polarimetric radar vegetation index for biomass estimation in desert fringe ecosystems. *IEEE Transactions on Geoscience and Remote Sensing*, 56(12), 7102–7108. <https://doi.org/10.1109/tgrs.2018.2848285>
- Dey, S., Bhattacharya, A., Frery, A. C., Lopez-Martinez, C., & Rao, Y. S. (2021). A Model-free Four Component Scattering Power Decomposition for Polarimetric SAR Data. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 1–1. <https://doi.org/10.1109/JSTARS.2021.3069299>
- Dey, S., Bhattacharya, A., Ratha, D., Mandal, D., & Frery, A. C. (2020). Target characterization and scattering power decomposition for full and compact polarimetric SAR data. *IEEE Transactions on Geoscience and Remote Sensing*. <https://doi.org/10.1016/j.isprsjprs.2020.09.010>
- Kim, Y., & Zyl, J. J. van. (2009). A time-series approach to estimate soil moisture using polarimetric radar data. *IEEE Trans. Geosci. Remote Sens.*, 47(8), 2519–2527. <https://doi.org/10.1109/TGRS.2009.2014944>
- Kumar, V., Mandal, D., Bhattacharya, A., & Rao, Y. (2020). Crop characterization using an improved scattering power decomposition technique for compact polarimetric SAR data. *International Journal of Applied Earth Observation and Geoinformation*, 88, 102052. <https://doi.org/10.1016/j.jag.2020.102052>
- Mandal, D., Kumar, V., Ratha, D., Dey, S., Bhattacharya, A., Lopez-Sanchez, J. M., McNairn, H., & Rao, Y. S. (2020). Dual polarimetric radar vegetation index for crop growth monitoring using sentinel-1 SAR data. *Remote Sensing of Environment*, 247, 111954. <https://doi.org/10.1016/j.rse.2020.111954>
- Mandal, D., Ratha, D., Bhattacharya, A., Kumar, V., McNairn, H., Rao, Y. S., & Frery, A. C. (2020). A radar vegetation index for crop monitoring using compact polarimetric SAR data. *IEEE Transactions on Geoscience and Remote Sensing*. <https://doi.org/10.1109/tgrs.2020.2976661>
- Ratha, D., Mandal, D., Kumar, V., McNairn, H., Bhattacharya, A., & Frery, A. C. (2019). A generalized volume scattering model-based vegetation index from polarimetric SAR data. *IEEE Geoscience and Remote Sensing Letters*, 16(11), 1791–1795. <https://doi.org/10.1109/lgrs.2019.2907703>