




# PinCFlow.jl: An idealized-atmospheric-flow solver coupled to the 3D transient gravity-wave model MS-GWaM

Felix Jochum<sup>1</sup>, Irmgard Knop<sup>1</sup>, Sandra Klewinghaus<sup>1</sup>, Jonas Rothermel<sup>1</sup>, Marco Artiano<sup>2</sup>, Arpit Babbar<sup>2</sup>, Hendrik Ranocha<sup>2</sup>, Stamen Dolaptchiev<sup>1</sup>, and Ulrich Achatz<sup>1</sup>

<sup>1</sup> Institute for Atmospheric and Environmental Sciences, Goethe University Frankfurt, Germany <sup>2</sup> Institute of Mathematics, Johannes Gutenberg University Mainz, Germany ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: 

Submitted: 17 November 2025

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Summary

PinCFlow.jl (*Pseudo-inCompressible Flow* solver) is an atmospheric-flow solver that was designed for conducting idealized simulations. It integrates the Boussinesq, pseudo-incompressible, and compressible equations in a conservative flux form (Klein, 2009; Rieper et al., 2013), using a semi-implicit method that combines explicit and implicit time-stepping schemes (Benacchio & Klein, 2019; Chew et al., 2022; Schmid et al., 2021). Spatially, the equations are discretized with a finite-volume method, such that all quantities are represented by spatial averages over grid cells and fluxes are computed on the respective cell interfaces. The grid is staggered (Arakawa & Lamb, 1977) so that the velocity components are defined at the same points as the corresponding fluxes of scalar quantities. PinCFlow.jl operates in a vertically stretched terrain-following coordinate system based on Gal-Chen & Somerville (1975a), Gal-Chen & Somerville (1975b), and Clark (1977).

The Lagrangian gravity-wave parameterization MS-GWaM (*Multi-Scale Gravity-Wave Model*) is interactively coupled to the dynamical core of PinCFlow.jl, so that unresolved gravity waves may be parameterized in a manner that accounts for transient wave-mean-flow interaction and horizontal wave propagation. The resolved fields are updated with tendencies computed by MS-GWaM at the beginning of every time step. A description of the theory behind MS-GWaM can be found in Achatz et al. (2017) and Achatz et al. (2023). For a numerical perspective and more information on the development, see Muraschko et al. (2015), Bölöni et al. (2016), Wilhelm et al. (2018), Wei et al. (2019), and Jochum et al. (2025).

## Statement of need

Atmospheric gravity waves are generated by various sources, including convection, flow over mountains, and jet-frontal imbalances. As they propagate upward, they transport and deposit momentum and energy, thus impacting the large-scale flow. A significant part of the gravity-wave spectrum cannot be resolved by weather and climate models at their respective standard resolutions and must therefore be parameterized (Achatz et al., 2024; Alexander et al., 2010; Fritts & Alexander, 2003).

Most gravity-wave parameterizations rely on single-column and steady-state approximations, since these yield a comparatively simple and efficient method of calculating the mean-flow impact. However, it has become increasingly apparent that transient wave-mean-flow interaction and horizontal gravity-wave propagation need to be accounted for in an accurate description of gravity-wave-mean-flow interaction (Bölöni et al., 2016; Ehard et al., 2017; Muraschko et al.,

2015; Sato et al., 2009; Senf & Achatz, 2011; Wei et al., 2019). The Multi-Scale Gravity-Wave Model (MS-GWaM) has been created for this reason. Essential to its development, as to the improvement of the fundamental understanding of atmospheric dynamics, is the study of gravity waves in idealized simulations. PinCFlow.jl provides a modeling infrastructure tailored to this exact need. Moreover, it is interactively coupled to MS-GWaM, thus allowing for direct comparison between high-resolution wave-resolving simulations and coarse-resolution wave-parameterizing simulations.

## Features

Figure 1 shows an overview of PinCFlow.jl's features. As has been mentioned above, the model can be used to integrate the Boussinesq, pseudo-incompressible or compressible equations. For each of these, it provides a number of background atmospheres defined by model parameters. The initial deviations of the prognostic variables from the background state are set with user-defined functions. In addition to the momentum, continuity, and potential-temperature equations, the flow solver integrates the prognostic equation for an arbitrary passive tracer with direct gravity-wave impact parameterized using MS-GWaM according to Knop et al. (2026).




Figure 1: Overview of PinCFlow.jl's features.

PinCFlow.jl's grid is determined from a user-defined surface topography, using a height-based terrain-following coordinate system. Fluxes are computed with a finite-volume (MUSCL) scheme (e.g., Leer, 2003), using the monotized-centered variant limiter (e.g., Kemm, 2011). In addition to the advection terms, the model includes molecular friction, heat conduction, and a turbulent viscosity.

MS-GWaM can be run in a 1D steady-state mode (which represents classic gravity-wave

parameterizations), a 1D transient mode (so that the impact of transient wave-mean-flow interaction may be investigated separately, i.e., without horizontal propagation), and a 3D transient mode (the full implementation of the theory). It is interactively coupled to the resolved dynamics, so that the evolution of the gravity-wave field depends on the current state of the large-scale flow and in turn changes it via tendencies that are added to the prognostic equations. This coupling includes the impact of gravity-wave fluxes on the large-scale tracer. The parameterization has an inbuilt orographic source which launches mountain waves generated by the resolved flow passing over unresolved terrain. The latter is represented by an arbitrary spectrum computed by a user-defined function. Moreover, MS-GWaM may be initialized with any wave field, also via a user-defined function.

By default, the flow solver uses an adaptive time step determined from several stability criteria, including CFL conditions with respect to the resolved flow and the maximum group velocities of the unresolved gravity-wave field computed by MS-GWaM. The CFL numbers are configurable, however, one may also run the model with a fixed time step.

MPI parallelization is supported in all grid dimensions. The number of MPI tasks is set per dimension, using corresponding model parameters. Furthermore, one may pass a base communicator to the model, facilitating the setup of ensemble simulations in a single Julia script.

## Software design

PinCFlow.jl's design philosophy aims to balance a user-friendly and minimal Public API with a well-structured and easily extendible model. The code is organized in nested modules that group related functions and types. Most functions take an instance of PinCFlow.jl's type `State` as input, which contains all information on the model's configuration and current simulation state, organized in nested composite types. Julia's multiple dispatch is heavily used to implement algorithms that are specialized for the configuration provided by the user. This facilitates the continued development of the model, since adding a new feature can be as simple as defining a new type for an existing model parameter and dispatching to an appropriate method where the feature comes into play. All of PinCFlow.jl's functions have docstrings with detailed descriptions of each of their methods, which can be found in the reference section of the documentation.

The Public API consists of all objects that are exported by PinCFlow.jl's main module, including the `State` type, a composite type called `Namelists` and the types of its fields (which are used to create a `State` instance in a particular configuration), model parameter types, and the `integrate` function (which also takes an instance of `Namelists` as input). An extension for the visualization package Makie.jl (Danisch & Krumbiegel, 2021) is provided, which exports tools for plotting the model output. A user guide and a number of examples that illustrate the usage of PinCFlow.jl are given in the documentation.

## Research impact statement

PinCFlow.jl originated from a pseudo-incompressible flow solver with implicit turbulence parameterization (PincFloIt), written in Fortran by Rieper et al. (2013). PincFloIt was used by Bölöni et al. (2016) to conduct wave-resolving simulations of gravity-wave packets, which provided reference data for the validation of an early version of MS-GWaM. An important result of this study was that, due to its accounting for transience, MS-GWaM yielded a more accurate forcing of the mean flow than the classic steady-state approach. Wilhelm et al. (2018) modified PincFloIt to also integrate the Boussinesq equations, coupled MS-GWaM to it, and validated the latter against submesoscale-resolving simulations of mesoscale inertia gravity waves radiated by a submesoscale gravity-wave packet. Wei et al. (2019) used this new framework to show that the common approach of using the gravity-wave pseudomomentum-flux

convergence to force the mean flow and thus neglect elastic and thermal effects is less reliable than using a direct approach which includes these contributions. Due to the implicit turbulence parameterization no longer being used, the model was renamed PincFlow when Schmid et al. (2021) implemented a semi-implicit time scheme (based on Benacchio & Klein, 2019) and verified its accuracy and efficiency in baroclinic life-cycle simulations. Using this scheme in conjunction with a newly added terrain-following coordinate system, Jochum et al. (2025) extended PincFlow's implementation of MS-GWaM with an orographic source and showed that the parameterization is able to capture highly transient interaction between mountain waves and mean flow in an idealized setting, with wave-resolving simulations providing a reference. Finally, Knop et al. (2026) showed that MS-GWaM can improve the representation of gravity-wave-induced tracer transport through the inclusion of a tracer Stokes drift due to unresolved gravity waves.

PinCFlow.jl is based on a translation of PincFlow from Fortran to Julia. Apart from time-dependent orography, heating above flat terrain in the pseudo-incompressible regime, alternative turbulence schemes, and a fully explicit time scheme, it offers all the features that were essential to the studies listed above. In addition, it introduces several new features, most notably access to the compressible regime (allowing for heating above arbitrary terrain), vertical grid stretching, and arbitrary initialization of the deviations from the atmospheric background. The implementation of the compressible mode necessitated slight modifications to the time scheme, based on the temporal discretization in the models introduced by Benacchio & Klein (2019) and Chew et al. (2022). Note that in contrast to these models, PinCFlow.jl does not provide seamless access to soundproof and hydrostatic dynamics (instead, the prognostic equations are set in the initialization and not changed during the simulation) or any data assimilation capabilities.

## State of the field

What distinguishes PinCFlow.jl from other software is its implementation of MS-GWaM. The latter was first developed as an isolated code (Böloni et al., 2016; Muraschko et al., 2015) and later as a part of the predecessor of PinCFlow.jl (by Jochum et al., 2025; Knop et al., 2026; Wei et al., 2019; Wilhelm et al., 2018) and within the NWP-and-climate model ICON (by Böloni et al., 2021; Kim et al., 2021, 2024; Voelker et al., 2024). PinCFlow.jl's version of MS-GWaM differs from the ICON version in several components, most notably its coordinate system, discretization, and sources, however, the underlying theory (Achatz et al., 2017, 2023) is the same. In contrast to other gravity-wave parameterizations, MS-GWaM relaxes the single-column and steady-state approximations in the description of the gravity-wave field and uses a direct approach (as opposed to the above-mentioned pseudomomentum approach) to calculate the mean-flow impact, resulting in additional force terms in the horizontal-momentum and potential-temperature equations (e.g., Wei et al., 2019). The conservation of gravity-wave energy is encoded in a prognostic equation for a wave-action density in the topological product of physical space and spectral space. This equation is integrated in a Lagrangian manner, so that wave action is essentially traced along rays in a six-dimensional phase space. Via this method, MS-GWaM avoids so-called caustics, i.e., ray intersections leading to an ill-defined wave-action-density equation, in a unique and physically sound manner (e.g., Muraschko et al., 2015). The numerical implementation involves the definition of rectangular ray volumes, which may overlap, move, stretch, split, and merge (e.g., Jochum et al., 2025).

The code presented here provides the first open access to a full implementation of MS-GWaM coupled to a flow solver within an idealized and highly configurable framework.

## AI usage disclosure

No generative AI tools were used in the software creation or writing of the paper. Parts of the reference section of the documentation were initially generated with AI but manually rewritten (in their entirety) later on.

## Acknowledgements

F. J. and U. A. are grateful for the support of Eric and Wendy Schmidt through the Schmidt Sciences VESRI “DataWave” project. S. D., I. K., and U. A. thank the German Research Foundation (DFG) for partial support through CRC 301 “TPChange” (Project No. 428312742 and Projects B06 “Impact of small-scale dynamics on UTLS transport and mixing”, B07 “Impact of cirrus clouds on tropopause structure”, and Z03 “Joint model development and modelling synthesis”), and U. A. thanks DFG for support through the CRC 181 “Energy transfers in Atmosphere and Ocean” (Project No. 274762653 and Projects W01 “Gravity-wave parameterization for the atmosphere” and S02 “Improved Parameterizations and Numerics in Climate Models”). M. A., A. B., and H. R. were supported by the German Research Foundation (Deutsche Forschungsgemeinschaft DFG, project numbers 528753982 and 526031774 within the SPP 2410 “Hyperbolic Balance Laws in Fluid Mechanics: Complexity, Scales, Randomness” (CoScaRa)). S. K. thanks Ray Chew for his help with the implementation of the compressible mode.

## References

- Achatz, U., Alexander, M. J., Becker, E., Chun, H.-Y., Dörnbrack, A., Holt, L., Plougonven, R., Polichtchouk, I., Sato, K., Sheshadri, A., Stephan, C. C., Niekerk, A. van, & Wright, C. J. (2024). Atmospheric gravity waves: Processes and parameterization. *Journal of the Atmospheric Sciences*, 81(2), 237–262. <https://doi.org/10.1175/jas-d-23-0210.1>
- Achatz, U., Kim, Y.-H., & Voelker, G. S. (2023). Multi-scale dynamics of the interaction between waves and mean flows: From nonlinear WKB theory to gravity-wave parameterizations in weather and climate models. *Journal of Mathematical Physics*, 64(11). <https://doi.org/10.1063/5.0165180>
- Achatz, U., Ribstein, B., Senf, F., & Klein, R. (2017). The interaction between synoptic-scale balanced flow and a finite-amplitude mesoscale wave field throughout all atmospheric layers: Weak and moderately strong stratification. *Quarterly Journal of the Royal Meteorological Society*, 143(702), 342–361. <https://doi.org/10.1002/qj.2926>
- Alexander, M. J., Geller, M., McLandress, C., Polavarapu, S., Preusse, P., Sassi, F., Sato, K., Eckermann, S., Ern, M., Hertzog, A., Kawatani, Y., Pulido, M., Shaw, T. A., Sigmond, M., Vincent, R., & Watanabe, S. (2010). Recent developments in gravity-wave effects in climate models and the global distribution of gravity-wave momentum flux from observations and models. *Quarterly Journal of the Royal Meteorological Society*, 136(650), 1103–1124. <https://doi.org/10.1002/qj.637>
- Arakawa, A., & Lamb, V. R. (1977). Computational design of the basic dynamical processes of the UCLA general circulation model. In *General circulation models of the atmosphere* (pp. 173–265). Elsevier. <https://doi.org/10.1016/b978-0-12-460817-7.50009-4>
- Benacchio, T., & Klein, R. (2019). A semi-implicit compressible model for atmospheric flows with seamless access to soundproof and hydrostatic dynamics. *Monthly Weather Review*, 147(11), 4221–4240. <https://doi.org/10.1175/mwr-d-19-0073.1>
- Bölöni, G., Kim, Y.-H., Borchert, S., & Achatz, U. (2021). Toward transient subgrid-scale gravity wave representation in atmospheric models. Part I: Propagation model including



- 201 nondissipative wave–mean-flow interactions. *Journal of the Atmospheric Sciences*, 78(4),  
202 1317–1338. <https://doi.org/10.1175/jas-d-20-0065.1>
- 203 Bölöni, G., Ribstein, B., Muraschko, J., Sgoff, C., Wei, J., & Achatz, U. (2016). The interaction  
204 between atmospheric gravity waves and large-scale flows: An efficient description beyond  
205 the nonacceleration paradigm. *Journal of the Atmospheric Sciences*, 73(12), 4833–4852.  
206 <https://doi.org/10.1175/jas-d-16-0069.1>
- 207 Chew, R., Benacchio, T., Hastermann, G., & Klein, R. (2022). A one-step blended soundproof-  
208 compressible model with balanced data assimilation: Theory and idealized tests. *Monthly*  
209 *Weather Review*, 150(9), 2231–2254. <https://doi.org/10.1175/mwr-d-21-0175.1>
- 210 Clark, T. L. (1977). A small-scale dynamic model using a terrain-following coordinate  
211 transformation. *Journal of Computational Physics*, 24(2), 186–215. [https://doi.org/10.1016/0021-9991\(77\)90057-2](https://doi.org/10.1016/0021-9991(77)90057-2)
- 212
- 213 Danisch, S., & Krumbiegel, J. (2021). Makie.jl: Flexible high-performance data visualization for  
214 Julia. *Journal of Open Source Software*, 6(65), 3349. <https://doi.org/10.21105/joss.03349>
- 215 Ehard, B., Kaifler, B., Dörnbrack, A., Preusse, P., Eckermann, S. D., Bramberger, M., Gisinger,  
216 S., Kaifler, N., Liley, B., Wagner, J., & Rapp, M. (2017). Horizontal propagation of  
217 large-amplitude mountain waves into the polar night jet. *Journal of Geophysical Research:*  
218 *Atmospheres*, 122(3), 1423–1436. <https://doi.org/10.1002/2016jd025621>
- 219 Fritts, D. C., & Alexander, M. J. (2003). Gravity wave dynamics and effects in the middle  
220 atmosphere. *Reviews of Geophysics*, 41(1). <https://doi.org/10.1029/2001rg000106>
- 221 Gal-Chen, T., & Somerville, R. C. J. (1975a). On the use of a coordinate transformation for  
222 the solution of the Navier-Stokes equations. *Journal of Computational Physics*, 17(2),  
223 209–228. [https://doi.org/10.1016/0021-9991\(75\)90037-6](https://doi.org/10.1016/0021-9991(75)90037-6)
- 224 Gal-Chen, T., & Somerville, R. C. J. (1975b). Numerical solution of the Navier-Stokes  
225 equations with topography. *Journal of Computational Physics*, 17(3), 276–310. [https://doi.org/10.1016/0021-9991\(75\)90054-6](https://doi.org/10.1016/0021-9991(75)90054-6)
- 226
- 227 Jochum, F., Chew, R., Lott, F., Voelker, G. S., Weinkaemmerer, J., & Achatz, U. (2025). The  
228 impact of transience in the interaction between orographic gravity waves and mean flow.  
229 *Journal of the Atmospheric Sciences*, 82(2), 425–442. <https://doi.org/10.1175/jas-d-24-0158.1>
- 230
- 231 Kemm, F. (2011). A comparative study of TVD-limiters—well-known limiters and an  
232 introduction of new ones. *International Journal for Numerical Methods in Fluids*, 67(4),  
233 404–440. <https://doi.org/10.1002/flid.2357>
- 234 Kim, Y.-H., Bölöni, G., Borchert, S., Chun, H.-Y., & Achatz, U. (2021). Toward transient  
235 subgrid-scale gravity wave representation in atmospheric models. Part II: Wave intermittency  
236 simulated with convective sources. *Journal of the Atmospheric Sciences*, 78(4), 1339–1357.  
237 <https://doi.org/10.1175/jas-d-20-0066.1>
- 238 Kim, Y.-H., Voelker, G. S., Bölöni, G., Zängl, G., & Achatz, U. (2024). Crucial role of  
239 obliquely propagating gravity waves in the quasi-biennial oscillation dynamics. *Atmospheric*  
240 *Chemistry and Physics*, 24(5), 3297–3308. <https://doi.org/10.5194/acp-24-3297-2024>
- 241 Klein, R. (2009). Asymptotics, structure, and integration of sound-proof atmospheric flow  
242 equations. *Theoretical and Computational Fluid Dynamics*, 23(3), 161–195. <https://doi.org/10.1007/s00162-009-0104-y>
- 243
- 244 Knop, I., Dolaptchiev, S., & Achatz, U. (2026). Impact of small-scale gravity waves on tracer  
245 transport. *Quarterly Journal of the Royal Meteorological Society*. <https://doi.org/10.1002/qj.70091>
- 246
- 247 Leer, B. van. (2003, June). Upwind and high-resolution methods for compressible flow: From

- 248 donor cell to residual-distribution schemes. *16<sup>th</sup> AIAA Computational Fluid Dynamics*  
 249 *Conference*. <https://doi.org/10.2514/6.2003-3559>
- 250 Muraschko, J., Fruman, M. D., Achatz, U., Hickel, S., & Toledo, Y. (2015). On the application  
 251 of Wentzel–Kramer–Brillouin theory for the simulation of the weakly nonlinear dynamics of  
 252 gravity waves. *Quarterly Journal of the Royal Meteorological Society*, 141(688), 676–697.  
 253 <https://doi.org/10.1002/qj.2381>
- 254 Rieper, F., Hickel, S., & Achatz, U. (2013). A conservative integration of the pseudo-  
 255 incompressible equations with implicit turbulence parameterization. *Monthly Weather*  
 256 *Review*, 141(3), 861–886. <https://doi.org/10.1175/mwr-d-12-00026.1>
- 257 Sato, K., Watanabe, S., Kawatani, Y., Tomikawa, Y., Miyazaki, K., & Takahashi, M. (2009).  
 258 On the origins of mesospheric gravity waves. *Geophysical Research Letters*, 36(19).  
 259 <https://doi.org/10.1029/2009gl039908>
- 260 Schmid, F., Gagarina, E., Klein, R., & Achatz, U. (2021). Toward a numerical laboratory for  
 261 investigations of gravity wave–mean flow interactions in the atmosphere. *Monthly Weather*  
 262 *Review*, 149(12), 4005–4026. <https://doi.org/10.1175/mwr-d-21-0126.1>
- 263 Senf, F., & Achatz, U. (2011). On the impact of middle-atmosphere thermal tides on the  
 264 propagation and dissipation of gravity waves. *Journal of Geophysical Research: Atmospheres*,  
 265 116(D24). <https://doi.org/10.1029/2011jd015794>
- 266 Voelker, G. S., Bölöni, G., Kim, Y.-H., Zängl, G., & Achatz, U. (2024). MS-GWaM: A  
 267 three-dimensional transient gravity wave parametrization for atmospheric models. *Journal*  
 268 *of the Atmospheric Sciences*, 81(7), 1181–1200. <https://doi.org/10.1175/jas-d-23-0153.1>
- 269 Wei, J., Bölöni, G., & Achatz, U. (2019). Efficient modeling of the interaction of mesoscale  
 270 gravity waves with unbalanced large-scale flows: Pseudomomentum-flux convergence  
 271 versus direct approach. *Journal of the Atmospheric Sciences*, 76(9), 2715–2738. <https://doi.org/10.1175/jas-d-18-0337.1>
- 272
- 273 Wilhelm, J., Akylas, T. R., Bölöni, G., Wei, J., Ribstein, B., Klein, R., & Achatz, U.  
 274 (2018). Interactions between mesoscale and submesoscale gravity waves and their efficient  
 275 representation in mesoscale-resolving models. *Journal of the Atmospheric Sciences*, 75(7),  
 276 2257–2280. <https://doi.org/10.1175/jas-d-17-0289.1>