

# EmissV: a R package to create vehicular and other emissions for air quality models

Daniel Schuch<sup>1</sup>, Sergio Ibarra-Espinosa<sup>1</sup>, and Edmilson Dias de Freitas<sup>1</sup>

<sup>1</sup> Departamento de Ciências Atmosféricas, Universidade de São Paulo, Brasil

DOI: [10.21105/joss.00662](https://doi.org/10.21105/joss.00662)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Submitted: 14 March 2018

Published: 07 August 2018

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

## Summary

Air quality models need input data containing information about atmosphere (such as temperature, wind, humidity), terrestrial data (such as terrain, land use, soil types) and emissions. Therefore, the emission inventories are easily seen as the scapegoat if a mismatch is found between modelled and observed concentrations of air pollutants (Pulles and Heslinga 2010). The anthropogenic emissions, especially vehicular emissions, are highly dependent on human activity and constantly changing due to various factors ranging from economic (such as the state of conservation of the fleet, renewal of the fleet and the price of fuel) to legal aspects (such as the vehicle routing).

The EmissV is a R package that estimates vehicular emissions by a top-down approach, the emissions are calculated using the statistical description of the fleet at available level (National, State, City, etc). The following steps show an example of the workflow for calculating vehicular emissions, this emissions are initially temporally and spatially disaggregated, and then distributed spatially and temporally to be used as input in numeric air quality models such WRF-Chem (Grell et al. 2005).

**I.** Total: emission of pollutants is estimated from the fleet (number, type and year of vehicles), vehicular activity (km/day) and emission factors (g/km) by pollutant for each interest area (cities, states, countries, etc) or alternatively the totals of some inventory can be used.

**II.** Spatial distribution: the package has functions to read information from tables, georeferenced images (tiff), shapefiles (sh), openstreetmap data (osm), global inventories in NetCDF format (nc) to calculate point, line and area sources.

**III.** Emission calculation: calculates the final emission from all different sources and converts to model units and resolution.

**IV.** Temporal distribution: a set of hourly profiles that represents the mean activity (by hour and day of the week) calculated from traffic counts of toll stations located at São Paulo city available for apply in the emissions.

The package has additional functions for creating emissions from individual sources (including plume rise parameterizations) and to estimate the vehicular emissions of volatile organic compounds from exhaust (through the exhaust pipe), liquid (carter and evaporative) and vapor (fuel transfer operations).

## Functions and data

EmissV counts with the following functions:

Function	Description
areaSource	Distribution of emissions by area
emission	Emissions in the format for atmospheric models
emissionFactors	Tool to set-up vehicle emission factors
gridInfo	Read grid information from a NetCDF file
lineSource	Distribution of emissions by streets
perfil	Dataset with temporal profile for vehicular emissions
plumeRise	Calculate plume rise height
pointSource	Emissions from point sources
rasterSource	Distribution of emissions by a georeferenced image
read	Read NetCDF data from global inventories
streedDist	Distribution by OpenStreetMap street
totalEmission	Calculate total emissions
totalVOC	Calculate total VOCs emissions
vehicles	Tool to set-up vehicle data frame

## Examples

The following example creates an area source for São Paulo State (Brasil). The **vehicles** function creates a **data.frame** with information about the São Paulo Fleet using data from (DETRAN 2015), the **emissionFactors** creates a **data.frame** with emission factors for CO and PM (CETESB 2015). The **totalEmission** calculates the total emissions of CO for these vehicles and this emission factors. The next 3 lines opens different data: wrf file, a raster and the area shapefiles. These data are the input for **areaSouce** that creates an area source based on an image of persistent lights of the Defense Meteorological Satellite Program (DMSP) for São Paulo and Minas Gerais (Brasil) states and finally the function **emission** calculates the CO emissions.

```
library(EmissV)
```

```
fleet <- vehicles(example = T)
```

```
# using a example of vehicles (DETRAN 2016 data and SP vehicle distribution):
#
# Category      Type Fuel      Use      SP ...
# Light Duty Vehicles Gasohol LDV_E25 LDV E25 41 km/d 11624342 ...
# Light Duty Vehicles Ethanol LDV_E100 LDV E100 41 km/d 874627 ...
# Light Duty Vehicles Flex LDV_F LDV FLEX 41 km/d 9845022 ...
# Diesel Trucks TRUCKS_B5 TRUCKS B5 110 km/d 710634 ...
# Diesel Urban Busses CBUS_B5 BUS B5 165 km/d 792630 ...
# Diesel Intercity Busses MBUS_B5 BUS B5 165 km/d 21865 ...
# Gasohol Motorcycles MOTO_E25 MOTO E25 140 km/d 3227921 ...
# Flex Motorcycles MOTO_F MOTO FLEX 140 km/d 235056 ...
```

```
# dropping the fleet from Rio de Janeiro (RJ), Parana (PR) and Santa Catarina (SC)
fleet <- fleet[,c(-6,-8,-9)]
```

```
EF <- emissionFactor(example = T)
```

```
# using a example emission factor (values calculated from CETESB 2015):
#
# CO      PM
# Light Duty Vehicles Gasohol 1.75 g/km 0.0013 g/km
# Light Duty Vehicles Ethanol 10.04 g/km 0.0000 g/km
# Light Duty Vehicles Flex 0.39 g/km 0.0010 g/km
# Diesel Trucks 0.45 g/km 0.0612 g/km
# Diesel Urban Busses 0.77 g/km 0.1052 g/km
```

```
# Diesel Intercity Busses      1.48 g/km 0.1693 g/km
# Gasohol Motorcycles         1.61 g/km 0.0000 g/km
# Flex Motorcycles            0.75 g/km 0.0000 g/km

TOTAL <- totalEmission(fleet,EF,pol = c("CO"),verbose = T)
# [1] "Total of CO : 1127548.9048935 t year-1"

grid <- gridInfo(paste(system.file("extdata", package = "EmissV"),
                               "/wrfinput_d02",sep=""))
# [1] "Grid information from: ../EmissV/extdata/wrfinput_d02"

raster <- raster::raster(paste(system.file("extdata", package = "EmissV"),
                               "/dmsp_hi-res.tiff",sep=""))

shape <- raster::shapefile(paste(system.file("extdata", package = "EmissV"),
                               "/BR.shp",sep=""),verbose = F)[12,1]

Minas_Gerais <- areaSource(shape,raster,grid,name = "Minas Gerais")
# [1] "processing Minas Gerais area ... "
# [1] "fraction of Minas Gerais area inside the domain = 0.0147607845622591"

shape <- raster::shapefile(paste(system.file("extdata", package = "EmissV"),
                               "/BR.shp",sep=""),verbose = F)[22,1]


Sao_Paulo <- areaSource(shape,raster,grid,name = "Sao Paulo")
# [1] "processing Sao Paulo area ... "
# [1] "fraction of Sao Paulo area inside the domain = 0.473260323300595"

sp::spplot(raster::merge(TOTAL[[1]][[1]] * Sao_Paulo, TOTAL[[1]][[2]] * Minas_Gerais),
            scales = list(draw=TRUE),ylab="Lat",xlab="Lon",
            main=list(label="Emissions of CO [g/d]"),
            col.regions = c("#031638", "#001E48", "#002756", "#003062",
                           "#003A6E", "#004579", "#005084", "#005C8E",
                           "#006897", "#0074A1", "#0081AA", "#008FB3",
                           "#009EBD", "#00AFC8", "#00C2D6", "#00E3F0"))


CO_emissions <- emission(TOTAL,"CO",list(SP = Sao_Paulo, MG = Minas_Gerais),
                          grid,mm=28, plot = T)
# [1] "calculating emissions for CO using molar mass = 28 ..."
```

The emissions of CO calculated in this example can be seen in Figure 1 in g/d (by pixel) and the final emissions on Figure 2 in MOL h<sup>-1</sup> km<sup>-1</sup> (by model grid cell). This emissions can be written to WRF-Chem emission files using some package that makes the interface with NetCDF format such as **ncdf4** (Pierce 2017), **RNetCDF** (Michna and Milton Woods 2017), **ncdf.tools** (Buttlar 2015) or with the **eixport** (Ibarra-Espinosa and Schuch 2018).

The R package EmissV is available at the repository <https://github.com/atmoschem/EmissV>. And this installation is tested automatically on Linux via [TravisCI](#) and Windows via [Appveyor](#) continuous integration systems. Also, EmissV is already on CRAN <https://CRAN.R-project.org/package=EmissV>.



**Figure 1:** Emissions of CO using nocturnal lights.



**Figure 2:** CO emissions ready for use in air quality model.

## Acknowledgements

The development of EmissV was supported by postdoc grants from the Fundação da Universidade de São Paulo and Fundação Coordenação de Aperfeiçoamento de Pessoal de Nível Superior.

## References

- Buttler, Jannis v. 2015. *Ncdf.tools: Easier 'Netcdf' File Handling*. <https://CRAN.R-project.org/package=ncdf.tools>.
- CETESB. 2015. *Emissões Veiculares No Estado de São Paulo*. Série Relatórios. <http://cetesb.sp.gov.br/veicular/relatorios-e-publicacoes/>.
- DETRAN. 2015. *Relatórios Estatísticos: Frota de Veículos de 2015*. <https://www.denatran.gov.br/estatistica/257-frota-2015>.
- Grell, Georg A, Steven E Peckham, Rainer Schmitz, Stuart A McKeen, Gregory Frost, William C Skamarock, and Brian Eder. 2005. “Fully Coupled ‘Online’ Chemistry Within the Wrf Model.” *Atmospheric Environment* 39 (37). Elsevier:6957–75.
- Ibarra-Espinosa, Sergio, and Daniel Schuch. 2018. *Eixport: Export Emissions to Atmospheric Models*. <https://doi.org/10.21105/joss.00607>.
- Michna, Pavel, and with contributions from Milton Woods. 2017. *RNetCDF: Interface to Netcdf Datasets*. <https://CRAN.R-project.org/package=RNetCDF>.
- Pierce, David. 2017. *Ncdf4: Interface to Unidata netCDF (Version 4 or Earlier) Format Data Files*. <https://CRAN.R-project.org/package=ncdf4>.
- Pulles, Tim, and Dick Heslinga. 2010. “The Art of Emission Inventorying.” *TNO, Utrecht*. <https://doi.org/doi.org/10.13140/RG.2.1.2082.8007>.