

GPUE: Graphics Processing Unit Gross–Pitaevskii Equation solver

James R. Schloss¹ and Lee J. O’Riordan¹

¹ Okinawa Institute of Science and Technology Graduate University, Onna-son, Okinawa 904-0495, Japan.

DOI: [10.21105/joss.01037](https://doi.org/10.21105/joss.01037)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Submitted: 15 October 2018

Published: 11 December 2018

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

Summary

Bose–Einstein Condensates (BECs) are superfluid systems consisting of bosonic atoms that have been cooled and condensed into a single, macroscopic ground state (Fetter, 2009; Pethick & Smith, 2008). These systems can be created in an experimental laboratory and allow for the the exploration of physical phenomenon such as superfluid turbulence (Navon, Gaunt, Smith, & Hadzibabic, 2016; Roche & Barenghi, 2008; White, Anderson, & Bagnato, 2014), chaotic dynamics (Gardiner, 2002; Kyriakopoulos, Koukoulouyannis, Skokos, & Kevrekidis, 2014; Zhang, 2017), and analogues of other quantum systems (Dalibard, Gerbier, Juzeliūnas, & Öhberg, 2011). Numerical simulations of BECs that directly mimic experiments are valuable to fundamental research in these areas and allow for theoretical advances before experimental validation. The dynamics of BEC systems can be found by solving the non-linear Schrödinger equation known as the Gross–Pitaevskii Equation (GPE),

$$i\hbar \frac{\partial \Psi(\mathbf{r}, t)}{\partial t} = \left(-\frac{\hbar^2}{2m} \nabla^2 + V(\mathbf{r}) + g|\Psi(\mathbf{r}, t)|^2 \right) \Psi(\mathbf{r}, t),$$

where $\Psi(\mathbf{r}, t)$ is the three-dimensional many-body wavefunction of the quantum system, $\mathbf{r} = (x, y, z)$, m is the atomic mass, $V(\mathbf{r})$ is an external potential, $g = \frac{4\pi\hbar^2 a_s}{m}$ is a coupling factor, and a_s is the scattering length of the atomic species. Here, the GPE is shown in three dimensions, but it can easily be modified to one or two dimensions (Pethick & Smith, 2008). One of the most straightforward methods for solving the GPE is the split-operator method, which has previously been accelerated with GPU devices (Bauke & Keitel, 2011; Ruf, Bauke, & Keitel, 2009). No software packages are available using this method on GPU devices that allow for user-configurable simulations and a variety of different system types; however, several software packages exist to simulate BECs with other methods and on different architectures, including GPULab (Antoine & Duboscq, 2014) the Massively Parallel Trotter-Suzuki Solver (Witteck & Cucchietti, 2013), and XMDS (Dennis, Hope, & Johnsson, 2013).

GPUE is a GPU-based Gross–Pitaevskii Equation solver via the split-operator method for superfluid simulations of both linear and non-linear Schrödinger equations, emphasizing superfluid vortex dynamics in two and three dimensions. GPUE is a fast, robust, and accessible software suite to simulate physics for fundamental research in the area of quantum systems and has been used to manipulate large vortex lattices in two dimensions (O’Riordan & Busch, 2016; O’Riordan, White, & Busch, 2016) along with ongoing studies of vortex dynamics.

For these purposes, GPUE provides a number of unique features: 1. Dynamic field generation for trapping potentials and other variables on the GPU device. 2. Vortex tracking in 2D and vortex highlighting in 3D. 3. Configurable gauge fields for the generation of artificial magnetic fields and corresponding vortex distributions (Dalibard et al., 2011; Ghosh & Sachdeva, 2014). 4. Vortex manipulation via direct control of the wavefunction phase (Dobrek et al., 1999).

All of these features enable GPUE to simulate a wide variety of linear and non-linear dynamics of quantum systems. GPUE additionally features a numerical solver that improves on other available suites (O’Riordan, 2017; Wittek, 2016). All of GPUE’s features and functionality have been described in further detail in the documentation (Schloss & O’Riordan, 2018).

Acknowledgements

This work has been supported by the Okinawa Institute of Science and Technology Graduate University and by JSPS KAKENHI Grant Number JP17J01488. We would also like to thank Thomas Busch, Rashi Sachdeva, Tiantian Zhang, Albert Benseney, and Angela White for discussions on useful physical systems to simulate with the GPUE codebase, along with Peter Wittek and Tadhg Morgan for contributions to the code, itself. These acknowledgements can be found in `GPUE/acknowledgements.md`.

References

- Antoine, X., & Duboscq, R. (2014). GPELab, a Matlab toolbox to solve Gross–Pitaevskii equations I: Computation of stationary solutions. *Computer Physics Communications*, 185(11), 2969–2991. doi:[10.1016/j.cpc.2014.06.026](https://doi.org/10.1016/j.cpc.2014.06.026)
- Bauke, H., & Keitel, C. H. (2011). Accelerating the Fourier split operator method via graphics processing units. *Computer Physics Communications*, 182(12), 2454–2463. doi:[10.1016/j.cpc.2011.07.003](https://doi.org/10.1016/j.cpc.2011.07.003)
- Dalibard, J., Gerbier, F., Juzeliūnas, G., & Öhberg, P. (2011). Colloquium: Artificial gauge potentials for neutral atoms. *Rev. Mod. Phys.*, 83(4), 1523–1543. doi:[10.1103/RevModPhys.83.1523](https://doi.org/10.1103/RevModPhys.83.1523)
- Dennis, G., Hope, J., & Johnsson, M. (2013). XMDS2: Fast, scalable simulation of coupled stochastic partial differential equations. *Computer Physics Communications*, 184(1), 201–208. doi:[10.1016/j.cpc.2012.08.016](https://doi.org/10.1016/j.cpc.2012.08.016)
- Dobrek, L., Gajda, M., Lewenstein, M., Sengstock, K., Birkel, G., & Ertmer, W. (1999). Optical generation of vortices in trapped Bose-Einstein condensates. *Phys. Rev. A*, 60(5), R3381–R3384. doi:[10.1103/PhysRevA.60.R3381](https://doi.org/10.1103/PhysRevA.60.R3381)
- Fetter, A. L. (2009). Rotating trapped Bose-Einstein condensates. *Rev. Mod. Phys.*, 81(2), 647–691. doi:[10.1103/RevModPhys.81.647](https://doi.org/10.1103/RevModPhys.81.647)
- Gardiner, S. A. (2002). (Quantum) chaos in Bose-Einstein condensates. *Journal of Modern Optics*, 49(12), 1971–1977. doi:[10.1080/09500340210140777](https://doi.org/10.1080/09500340210140777)
- Ghosh, S., & Sachdeva, R. (2014). Synthetic Gauge Fields for Ultra Cold Atoms: A Primer. *Journal of the Indian Institute of Science*, 94(2), 217–232. Retrieved from <http://journal.iisc.ernet.in/index.php/iisc/article/view/4502>
- Kyriakopoulos, N., Koukouloyannis, V., Skokos, C., & Kevrekidis, P. G. (2014). Chaotic behavior of three interacting vortices in a confined Bose-Einstein condensate. *Chaos*, 24(10), 104410. doi:[10.1063/1.4882169](https://doi.org/10.1063/1.4882169)

- Navon, N., Gaunt, A. L., Smith, R. P., & Hadzibabic, Z. (2016). Emergence of a turbulent cascade in a quantum gas. *Nature*, 539, 72. doi:[10.1038/nature20114](https://doi.org/10.1038/nature20114)
- O’Riordan, L. J. (2017). *Non-equilibrium vortex dynamics in rapidly rotating Bose–Einstein condensates* (PhD thesis). Okinawa Institute of Science; Technology Graduate University. Retrieved from <http://ci.nii.ac.jp/naid/500001054902>
- O’Riordan, L. J., & Busch, T. (2016). Topological defect dynamics of vortex lattices in Bose–Einstein condensates. *Phys. Rev. A*, 94(5), 053603. doi:[10.1103/PhysRevA.94.053603](https://doi.org/10.1103/PhysRevA.94.053603)
- O’Riordan, L. J., White, A. C., & Busch, T. (2016). Moiré superlattice structures in kicked Bose–Einstein condensates. *Phys. Rev. A*, 93(2), 023609. doi:[10.1103/PhysRevA.93.023609](https://doi.org/10.1103/PhysRevA.93.023609)
- Pethick, C. J., & Smith, H. (2008). *Bose–Einstein Condensation in Dilute Gases*. Cambridge University Press. doi:[10.1017/CBO9780511802850](https://doi.org/10.1017/CBO9780511802850)
- Roche, P.-E., & Barenghi, C. F. (2008). Vortex spectrum in superfluid turbulence: Interpretation of a recent experiment. *EPL (Europhysics Letters)*, 81(3), 36002. Retrieved from <http://stacks.iop.org/0295-5075/81/i=3/a=36002>
- Ruf, M., Bauke, H., & Keitel, C. H. (2009). A real space split operator method for the Klein–Gordon equation. *Journal of Computational Physics*, 228(24), 9092–9106. doi:[10.1016/j.jcp.2009.09.012](https://doi.org/10.1016/j.jcp.2009.09.012)
- Schloss, J., & O’Riordan, L. J. (2018). GPUE documentation website. <https://gpue-group.github.io/>.
- White, A. C., Anderson, B. P., & Bagnato, V. S. (2014). Vortices and turbulence in trapped atomic condensates. *Proceedings of the National Academy of Sciences*. doi:[10.1073/pnas.1312737110](https://doi.org/10.1073/pnas.1312737110)
- Wittek, P. (2016). Comparing three numerical solvers of the Gross-Pitaevskii equation. <https://web.archive.org/web/20171120181431/https://peterwittek.com/gpe-comparison.html>.
- Wittek, P., & Cucchietti, F. M. (2013). A second-order distributed Trotter–Suzuki solver with a hybrid CPU–GPU kernel. *Computer Physics Communications*, 184(4), 1165–1171. doi:[10.1016/j.cpc.2012.12.008](https://doi.org/10.1016/j.cpc.2012.12.008)
- Zhang, T. (2017). *Chaotic Vortex Dynamics in Bose-Einstein Condensates* (Master’s thesis). Norwegian University of Science; Technology, Norway. Retrieved from <http://hdl.handle.net/11250/2467021>