








# ThermoCodegen: a python/C++ package for the generation of custom thermodynamic models

Marc Spiegelman <sup>1,2¶</sup>, Owen Evans<sup>\*1,3</sup>, Mark Ghiorso <sup>4</sup>, Lucy Tweed <sup>2</sup>, and Cian Wilson <sup>5</sup>

<sup>1</sup> Dept. of Applied Physics and Applied Math, Columbia University <sup>2</sup> Dept. of Earth and Environmental Science, Columbia University <sup>3</sup> Now at Google <sup>4</sup> OFM Research <sup>5</sup> Earth and Planets Laboratory, Carnegie Institution for Science ¶ Corresponding author

DOI: [10.21105/joss.04874](https://doi.org/10.21105/joss.04874)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Kyle Niemeyer](#) 

## Reviewers:

- [@bocklund](#)
- [@rdguha1995](#)

Submitted: 31 July 2022

Published: 30 June 2023

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

Many important problems in solid-Earth science require a consistent coupling between thermodynamics and geodynamics. Examples include global mantle convection with thermodynamically consistent phase changes, the generation and flow of magma in the mantle, reactive fluid flow and geological carbon sequestration in the crust and even the coupled flow of water and ice in glaciers and ice sheets. Given the complexity of these systems, it is important that the individual user/scientist have considerable control over the formulation and computational choices involved for both the thermodynamic and geodynamic components of models while maintaining transparency and reproducibility. Access to flexible open-source software for the creation and exploration of thermodynamic models and their incorporation into large scale reactive flow simulations is essential for these endeavors.

## Statement of need

ThermoCodegen is a software package for the reproducible generation of custom thermodynamic and kinetic reaction models for minerals, melts and some aqueous fluids commonly encountered in the solid-Earth sciences. It is part of the larger [ENKI](#) project for thermodynamic modeling whose flagship code [ThermoEngine](#) ([Ghiorso & Wolf, 2019](#); [Wolf & Ghiorso, 2019](#)) provides a [JupyterHub](#), cloud-based environment for the interactive generation and exploration of thermodynamic models in Earth sciences. In particular, ThermoEngine provides a critical code-generation module, `coder.py` for generating high-performance C code from [SymPy](#) expressions for the free energies of thermodynamic endmembers and phases.

ThermoCodegen shares the same underlying code-generation infrastructure as ThermoEngine however its design is driven more by the need for ready incorporation of thermodynamics into large scale geodynamics simulation codes for both equilibrium and disequilibrium transport. Key additional features of ThermoCodegen include:

- *Portable model storage*: Thermodynamic model generation for endmembers and phases are described in detailed [Jupyter](#) notebooks using `coder.py` and [SymPy](#), however, the final models are stored in structured xml files that contain all the information required to describe and auto-generate code for thermodynamic databases of endmembers and phases.
- *Kinetic modeling*: ThermoCodegen also provides code generation and storage of custom kinetic reaction models built on the phase databases.

<sup>\*</sup>co-first author

- *C++ support*: The code also generates C++ libraries for inclusion in other modeling systems such as [TerraFERMA](#) (Wilson et al., 2017; Wilson & Spiegelman, 2015) or [ASPECT](#) (Bangerth et al., 2021; Heister et al., 2017; Kronbichler et al., 2012).
- *Python bindings*: Python bindings of the C++ classes are generated using [pybind11](#) (Wenkel et al., 2017) for direct import into python applications.
- *ThermoEngine support*: Python bindings of ThermoCodegen objects can also be imported back into ThermoEngine as a custom database for use in equilibration or calibration calculations (requires installing the [tcg\\_model\\_integration](#) branch of [ThermoEngine](#)).

### Relationship to other Software packages for Solid-Earth Thermodynamic modeling

Computational thermodynamic software has been a central tool for solid-Earth science for over 40 years and has focused primarily on calculating phase equilibrium for several distinct self-consistent “Thermodynamic Datasets” such as those for solid mineral phases (Berman, 1988; Holland & Powell, 1998, 2011; Stixrude & Lithgow-Bertelloni, 2005, 2011), silicate liquids (Ghiorso et al., 2002; Ghiorso & Sack, 1995; Holland et al., 2018), and ionic species in aqueous solutions (Shock et al., 1992; Tanger & Helgeson, 1988).

Many of these datasets are associated with specific software for calculating equilibrium assemblages using a range of Gibbs free energy minimization algorithms. Some of the most commonly used software includes the MELTs family of models (Asimow & Ghiorso, 1998; Ghiorso et al., 2002; Ghiorso & Gualda, 2015; Ghiorso & Sack, 1995; Gualda et al., 2012) for equilibration of silicate minerals and melts, which has also been repackaged as alphaMELTS (Antoshechkina & Ghiorso, 2018; Smith & Asimow, 2005) as well as in its current incarnation within the ENKI project. Primarily used for metamorphic petrology, [THERMOCALC](#)<sup>1</sup> (Powell et al., 1998) is closely associated with the thermodynamic models of Holland and Powell (Holland & Powell, 1998, 2011), while [HeFESTo](#) is used for the calculation of phase equilibria of the high pressure/temperature solid-state models of Stixrude and Lithgow-Bertelloni (Stixrude & Lithgow-Bertelloni, 2005, 2011). Note that not all of these packages have open-source licenses. In addition, there are several more general Gibbs free energy minimization packages that can be used with multiple data sets. [Purple\\_X](#) (Connolly, 1990, 2005, 2009) is one of the most widely used. More recent additions include [THERIAK-DOMINO](#) (Capitani & Petrakakis, 2010) which also supports the models of Holland and Powell, [Thermolab](#) (Vrijmoed & Podladchikov, 2022) a MATLAB based toolkit, and [MAGEMin](#) (Riel et al., 2022) an open-source Gibbs energy minimization package written in C (with [julia](#) and [MATLAB](#) bindings) for adaptive phase-diagram generation with a range of modern optimization techniques. In particular, [MAGEMin](#) includes the igneous thermodynamic database of Holland et al. (2018).

These packages provide critical thermodynamic modeling infrastructure for solid-Earth science but are complementary and distinct in design and applicability to ThermoCodegen. In particular, the primary features added by ThermoCodegen are:


- the ability to more easily create and explore custom, consistent models of all thermodynamic parameters through the use of symbolic differentiation and automatic code generation
- the ability to generate code for disequilibrium reaction kinetics
- the ability to create custom C++ and python libraries for direct inclusion in geodynamics models.

Given the complexity of these high-dimensional open-system problems, having effective open-source software for reproducible exploration of both equilibrium phase diagrams and disequilibrium transport remains a central challenge and goal in geodynamics and ThermoCodegen is designed to address several of these critical challenges.

<sup>1</sup>Not to be confused with the commercial package [Thermo-Calc](#)

## Software design/overview

The design of ThermoCodegen follows the hierarchical structure of thermodynamic modeling commonly used in Earth Sciences and illustrated in Figure 1.



**Figure 1:** The hierarchical structure of thermodynamic modeling.  $\mu_i^k$  is an endmember's chemical potential model and  $M_i^k$  its mass.  $G_i$  is a phase's Gibbs free energy model and  $\rho_i, C_{Pi}, s_i$  and  $\alpha_i$  are (for example) its density, heat capacity, specific entropy, and thermal expansivity, derived consistently by symbolic differentiation of  $G_i$ .  $\nu_j$  describes the stoichiometry of reactions between the endmembers of the phases,  $A_j$  the resulting reaction affinities,  $R_j$  their rates, and  $\Gamma_i^k$  and  $\Gamma_i$  the resulting endmember and phase source terms. The Thermodynamic Database and Reactive System are functions of the temperature,  $T$ , pressure  $P$ , the number of moles,  $n$ , or mass fractions,  $C$ , of the endmembers, and presence (e.g. volume fraction) of the phases,  $\phi$ . These can be supplied by coupling to a dynamic model that may additionally calculate phase velocities, e.g.,  $v_s$  and  $v_f$ . Image adapted from Tweed (2021).

At the lowest level are thermodynamic **Endmembers** which provide chemical potentials,  $\mu_i^k(T, P)$ , for stoichiometric chemical components. The actual components can be arbitrary (e.g., oxides, minerals, ionic species) but have fixed chemical composition and thermodynamic properties that only depend on temperature and pressure (or temperature and volume, for models of the Helmholtz free energy).

Given a space of thermodynamic endmembers, we can construct homogeneous **Phases** as combinations of endmembers and develop thermodynamic models for their Gibbs free energy,  $G_i(T, P, n_i)$ , that depend on temperature, pressure and the number of moles of each endmember  $n_i = n_i^k$ .

Given  $G_i(T, P, n_i)$  for each phase, all of the other thermodynamic properties (and their derivatives) can be derived by partial differentiation of  $G_i$ . For example, entropy  $S_i$ , volume  $V_i$ , chemical potentials  $\bar{\mu}_i^k$ , heat capacity  $C_{Pi}$  and thermal expansivity  $\alpha_i$  are all related to  $G_i$  by:

$$\begin{aligned} S_i &= -\frac{\partial G_i}{\partial T} & V_i &= \frac{\partial G_i}{\partial P} & \bar{\mu}_i^k &= \frac{\partial G_i}{\partial n_i^k} \\ C_{Pi} &= T \frac{\partial S_i}{\partial T} = -T \frac{\partial^2 G_i}{\partial T^2} & \alpha_i &= \frac{1}{V_i} \frac{\partial V_i}{\partial T} = \frac{1}{V_i} \frac{\partial^2 G_i}{\partial P^2} \end{aligned}$$

Thus if we can define an analytic model for  $G_i(T, P, n_i)$  for any phase, we can use symbolic differentiation and code-generation to provide custom routines to generate all thermodynamic properties consistently.

We define a set of phases, together with their constituent thermodynamic endmembers as a **Thermodynamic Database** corresponding to the blue box in Figure 1. Standard databases available in the literature include the mineral database of Berman (1988) and its extension to silicate liquids through the MELTs family of models (Ghiorso et al., 2002; Ghiorso & Sack, 1995), as well as the high-pressure mineral phases of Stixrude and Lithgow-Bertelloni (Stixrude & Lithgow-Bertelloni, 2005, 2011) or metamorphic phases of, for example, Holland & Powell (2011). All of these are currently available through python bindings to ObjectiveC code in

ThermoEngine. However, the ObjectiveC objects are not readily incorporated within other codes.

ThermoEngine also provides support for generating custom thermodynamic models of endmembers and phases through the `thermoengine.coder` module written by Mark Ghiorso. `coder` uses [SymPy](#) to describe the free energies of endmembers and phases and then uses automatic differentiation and code-generation to write efficient C source code (and wrap it in [Cython](#)) for all thermodynamic properties of phases. Our extensions in `thermocodegen.coder` generate the same underlying C code and interfaces as `thermoengine.coder` but extend its functionality in several critical ways for better integration with large-scale geodynamic modeling.

## ThermoCodegen

ThermoCodegen is designed to produce code for just the colored boxes in [Figure 1](#).

### Custom Thermodynamic Databases

For **Thermodynamic Databases** of endmembers and phases, ThermoCodegen extends `thermoengine.coder` to record the internal state of each thermodynamic model as a python dictionary in order to store and generate code from xml files that completely describe all variables, parameters and SymPy models for endmembers and phases. These xml files are parsed using the science neutral options package [Spud](#) ([Ham et al., 2009](#)). Endmembers are stored with the extension `.emml` and phases are stored as `.phml` files, which can be viewed with the Spud GUI `diamond`. Given a set of consistent endmember and phase description files, the ThermoCodegen script `tcg_buildddb` can be used to inspect, validate and autogenerate code for a custom Thermodynamic Database as well as provide C++ wrappers and python bindings using [pybind11](#).

At this point, all source code, together with `.emml` and `.phml` files describing the full thermodynamic models, and optionally compiled C++ and python libraries are made available as a compressed tar-ball. This object is considered a Thermodynamic Database and can be used locally or made available as a web-accessible URL, for example with a [zenodo](#) DOI.

Given a custom thermodynamic database, these can then be used in optimization codes such as `thermoengine.equilibrate` to find equilibrium assemblages that minimize the global Gibbs free energy.

### Reaction Kinetic Objects

For better integration with open-system geodynamics codes that do not assume equilibrium, ThermoCodegen also provides code for generating C++ libraries and python bindings for custom kinetics models that describe a set of reactions between endmembers and phases within a given thermodynamic database. The software provides interfaces for calculating affinities, reaction rates, and thermodynamic properties of phases as well as their exact derivatives for use in non-linear solvers that take Jacobians. These reaction kinetic models are stored as xml files with the extension `.rxml` from which custom C++ libraries and python bindings are generated using the build script `tcg_buildrx`.

These reaction libraries, along with the underlying database objects can then be incorporated in external modeling software such as ThermoEngine, TerraFERMA, ASPECT, Jupyter notebooks and general python or C++ codes. Inclusion in dynamic models is considered external to ThermoCodegen, however, the ThermoCodegen distribution also provides containers that include a complete installation of TerraFERMA along with some example test cases.

Detailed [documentation](#) for ThermoCodegen is available that includes a more detailed description of the software design as well as instructions for using the package through the provided docker and singularity containers (local installation guidelines are also available). Most importantly, the documentation also includes fully worked out examples illustrating the

work flow for using ThermoCodegen including a set of Jupyter notebooks for constructing and exploring a range of thermodynamic models.

## Acknowledgements

This work was supported by NSF SI<sup>2</sup> award NSF-ACI1550337, NSF-EAR1520732, NSF-EAR1049905 and the Sloan Foundation project on Net-Zero and Negative Emissions Technologies.

## References

- Antoshechkina, P. M., & Ghiorso, M. S. (2018). *MELTS for MATLAB: A new Educational and Research Tool for Computational Thermodynamics*. 2018, ED44B–23. <https://ui.adsabs.harvard.edu/abs/2018AGUFMED44B..23A>
- Asimow, P. D., & Ghiorso, M. S. (1998). Algorithmic modifications extending MELTS to calculate subsolidus phase relations. *American Mineralogist*, 83(9–10), 1127–1132. <https://doi.org/10.2138/am-1998-9-1022>
- Bangerth, W., Dannberg, J., Fraters, M., Gassmoeller, R., Glerum, A., Heister, T., & Naliboff, J. (2021). *ASPECT v2.3.0*. Zenodo. <https://doi.org/10.5281/ZENODO.5131909>
- Berman, R. G. (1988). Internally-Consistent Thermodynamic Data for Minerals in the System Na<sub>2</sub>O-K<sub>2</sub>O-CaO-MgO-FeO-Fe<sub>2</sub>O<sub>3</sub>-Al<sub>2</sub>O<sub>3</sub>-SiO<sub>2</sub>-TiO<sub>2</sub>-H<sub>2</sub>O-CO<sub>2</sub>. *Journal of Petrology*, 29(2), 445–522. <https://doi.org/10.1093/petrology/29.2.445>
- Capitani, C. de, & Petrakakis, K. (2010). The computation of equilibrium assemblage diagrams with Theriak/Domino software. *American Mineralogist*, 95(7), 1006–1016. <https://doi.org/10.2138/am.2010.3354>
- Connolly, J. A. D. (1990). Multivariable phase diagrams; an algorithm based on generalized thermodynamics. *American Journal of Science*, 290(6), 666–718. <https://doi.org/10.2475/ajs.290.6.666>
- Connolly, J. A. D. (2005). Computation of phase equilibria by linear programming: A tool for geodynamic modeling and its application to subduction zone decarbonation. *Earth and Planetary Science Letters*, 236(1–2), 524–541. <https://doi.org/10.1016/j.epsl.2005.04.033>
- Connolly, J. A. D. (2009). The geodynamic equation of state: What and how. *Geochemistry, Geophysics, Geosystems*, 10(10), Q10014. <https://doi.org/10.1029/2009GC002540>
- Ghiorso, M. S., & Gualda, G. A. R. (2015). An H<sub>2</sub>O–CO<sub>2</sub> mixed fluid saturation model compatible with rhyolite-MELTS. *Contributions to Mineralogy and Petrology*, 169(6), 53. <https://doi.org/10.1007/s00410-015-1141-8>
- Ghiorso, M. S., Hirschmann, M. M., Reiners, P. W., & Kress, V. C. (2002). The pMELTS: A revision of MELTS for improved calculation of phase relations and major element partitioning related to partial melting of the mantle to 3 GPa. *Geochemistry, Geophysics, Geosystems*, 3(5), 1–35. <https://doi.org/10.1029/2001GC000217>
- Ghiorso, M. S., & Sack, R. O. (1995). Chemical mass transfer in magmatic processes IV. A revised and internally consistent thermodynamic model for the interpolation and extrapolation of liquid-solid equilibria in magmatic systems at elevated temperatures and pressures. *Contributions to Mineralogy and Petrology*, 119(2–3), 197–212. <https://doi.org/10.1007/BF00307281>
- Ghiorso, M. S., & Wolf, A. S. (2019). *Thermodynamic Modeling Using ENKI: 1. Overview and Phase Equilibrium Applications*. 2019, ED53F–0897. <https://ui.adsabs.harvard.edu/abs/2019AGUFMED53F0897G>

- Gualda, G. A. R., Ghiorso, M. S., Lemons, R. V., & Carley, T. L. (2012). Rhyolite-MELTS: A Modified Calibration of MELTS Optimized for Silica-rich, Fluid-bearing Magmatic Systems. *Journal of Petrology*, 53(5), 875–890. <https://doi.org/10.1093/petrology/egr080>
- Ham, D. A., Farrell, P. E., Gorman, G. J., Maddison, J. R., Wilson, C. R., Kramer, S. C., Shipton, J., Collins, G. S., Cotter, C. J., & Piggott, M. D. (2009). Spud 1.0: Generalising and automating the user interfaces of scientific computer models. *Geoscientific Model Development*, 2(1), 33–42. <https://doi.org/10.5194/gmd-2-33-2009>
- Heister, T., Dannberg, J., Gassmüller, R., & Bangerth, W. (2017). High accuracy mantle convection simulation through modern numerical methods – II: Realistic models and problems. *Geophysical Journal International*, 210(2), 833–851. <https://doi.org/10.1093/gji/ggx195>
- Holland, T. J. B., Green, E. C. R., & Powell, R. (2018). Melting of Peridotites through to Granites: A Simple Thermodynamic Model in the System KNCFMASHTOCr. *Journal of Petrology*, 59(5), 881–900. <https://doi.org/10.1093/petrology/egy048>
- Holland, T. J. B., & Powell, R. (1998). An internally consistent thermodynamic data set for phases of petrological interest. *Journal of Metamorphic Geology*, 16(3), 309–343. <https://doi.org/10.1111/j.1525-1314.1998.00140.x>
- Holland, T. J. B., & Powell, R. (2011). An improved and extended internally consistent thermodynamic dataset for phases of petrological interest, involving a new equation of state for solids. *Journal of Metamorphic Geology*, 29(3), 333–383. <https://doi.org/10.1111/j.1525-1314.2010.00923.x>
- Kronbichler, M., Heister, T., & Bangerth, W. (2012). High accuracy mantle convection simulation through modern numerical methods: High accuracy mantle convection simulation. *Geophysical Journal International*, 191(1), 12–29. <https://doi.org/10.1111/j.1365-246X.2012.05609.x>
- Powell, R., Holland, T., & Worley, B. (1998). Calculating phase diagrams involving solid solutions via non-linear equations, with examples using THERMOCALC. *Journal of Metamorphic Geology*, 16(4), 577–588. <https://doi.org/10.1111/j.1525-1314.1998.00157.x>
- Riel, N., Kaus, B. J. P., Green, E. C. R., & Berlie, N. (2022). MAGEMin, an Efficient Gibbs Energy Minimizer: Application to Igneous Systems. *Geochemistry, Geophysics, Geosystems*, 23(7). <https://doi.org/10.1029/2022GC010427>
- Shock, E. L., Oelkers, E. H., Johnson, J. W., Sverjensky, D. A., & Helgeson, H. C. (1992). Calculation of the thermodynamic properties of aqueous species at high pressures and temperatures. Effective electrostatic radii, dissociation constants and standard partial molal properties to 1000°C and 5 kbar. *Journal of the Chemical Society, Faraday Transactions*, 88(6), 803–826. <https://doi.org/10.1039/FT9928800803>
- Smith, P. M., & Asimow, P. D. (2005). Adibat\_1ph: A new public front-end to the MELTS, pMELTS, and pHMELTS models. *Geochemistry, Geophysics, Geosystems*, 6(2). <https://doi.org/10.1029/2004GC000816>
- Stixrude, L., & Lithgow-Bertelloni, C. (2005). Thermodynamics of mantle minerals - I. Physical properties. *Geophysical Journal International*, 162(2), 610–632. <https://doi.org/10.1111/j.1365-246X.2005.02642.x>
- Stixrude, L., & Lithgow-Bertelloni, C. (2011). Thermodynamics of mantle minerals - II. Phase equilibria. *Geophysical Journal International*, 184(3), 1180–1213. <https://doi.org/10.1111/j.1365-246X.2010.04890.x>
- Tanger, J. C., & Helgeson, H. C. (1988). Calculation of the thermodynamic and transport properties of aqueous species at high pressures and temperatures; revised equations of



- state for the standard partial molal properties of ions and electrolytes. *American Journal of Science*, 288(1), 19–98. <https://doi.org/10.2475/ajs.288.1.19>
- Tweed, L. E. L. (2021). *Coupling the Thermodynamics, Kinetics and Geodynamics of Multiphase Reactive Transport in Earth's Interior* [PhD thesis, Columbia University]. <https://doi.org/10.7916/d8-wh72-vb08>
- Vrijmoed, J. C., & Podladchikov, Y. Y. (2022). Thermolab: A Thermodynamics Laboratory for Nonlinear Transport Processes in Open Systems. *Geochemistry, Geophysics, Geosystems*, 23(4). <https://doi.org/10.1029/2021GC010303>
- Wenkel, J., Rhineland, J., & Moldovan, D. (2017). *pybind11 – Seamless operability between C++11 and Python*. <https://github.com/pybind/pybind11>
- Wilson, C. R., & Spiegelman, M. (2015). *Software: TerraFERMA v1.0 — The Transparent Finite Element Rapid Model Assembler, open-source repository: terraferma.github.io*. [terraferma.github.io](https://github.com/terraferma/terraferma)
- Wilson, C. R., Spiegelman, M., & Keken, P. E. van. (2017). TerraFERMA: The Transparent Finite Element Rapid Model Assembler for multiphysics problems in Earth sciences. *Geochemistry, Geophysics, Geosystems*, 18(2), 769–810. <https://doi.org/10.1002/2016GC006702>
- Wolf, A. S., & Ghiorso, M. S. (2019). *Thermodynamic Modeling Using ENKI: 2. Extending Existing Models*. 2019, ED52A–05. <https://ui.adsabs.harvard.edu/abs/2019AGUFMED52A.05W>