

# Tools21cm: A python package to analyse the large-scale 21 cm signal from the Epoch of Reionization and Cosmic Dawn

Sambit K. Giri<sup>1, 2</sup>, Garrelt Mellema<sup>1</sup>, and Hannes Jensen<sup>3</sup>

<sup>1</sup> Department of Astronomy and Oskar Klein Centre, Stockholm University, AlbaNova, SE-106 91 Stockholm, Sweden <sup>2</sup> Institute for Computational Science, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland <sup>3</sup> Self

DOI: [10.21105/joss.02363](https://doi.org/10.21105/joss.02363)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Amy Roberts](#) ↗

## Reviewers:

- [@nicholebarry](#)
- [@aureliocarnero](#)
- [@ziotom78](#)

Submitted: 08 June 2020

Published: 07 August 2020

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Summary

The Cosmic Dawn (CD) and Epoch of Reionization (EoR) are among the least understood epochs from the history of our Universe. Large-scale cosmological simulations are useful to understand this era and help prepare for future observations. `Tools21cm` is a data manipulation and analysis package for studying such cosmological simulations. It is written in the python programming language and designed to be very user-friendly.

The 21 cm signal, produced by the spin-flip transition of neutral hydrogen, is a unique tracer of matter in large-scale structures present during the EoR and CD (e.g. Furlanetto, Oh, & Briggs, 2006; Pritchard & Loeb, 2012). This signal will be cosmologically redshifted and thus found at low radio frequencies. Much of the functionality of `Tools21cm` is focused on the construction and analysis of mock 21 cm observations in the context of current and upcoming radio telescopes, such as the Low Frequency Array (LOFAR; Haarlem et al., 2013), the Murchison Widefield Array (MWA; Tingay et al., 2013; Wayth et al., 2018), Hydrogen Epoch of Reionization Array (HERA; DeBoer et al., 2017) and the Square Kilometre Array (SKA; Mellema et al., 2013). `Tools21cm` post-processes cosmological simulation data to create mock 21 cm observations.

Radio telescopes typically observe the redshifted 21 cm signal over a range of frequencies and therefore produce 21 cm images at different redshifts. A sequence of such images from different redshifts is known as a tomographic data set and is three dimensional. `Tools21cm` can construct such tomographic data sets from simulation snapshots (Datta et al., 2012; Giri et al., 2018a). When constructing these data sets, it can also add the impact of peculiar velocities, leading to an effect known as redshift space distortions (e.g. Jensen et al., 2013, 2016; Giri et al., 2018a). See Giri (2019) for a detailed description of tomographic 21 cm data sets.

`Tools21cm` also includes tools to calculate a wide range of statistical quantities from simulated 21 cm data sets. These include one-point statistics, such as the global or sky-averaged signal as a function of frequency, as well the variance, skewness and kurtosis (e.g. Ross, Dixon, Iliev, & Mellema, 2017; Ross, Dixon, Ghara, Iliev, & Mellema, 2019). It can also characterise the spatial fluctuations in the signal through spherically and cylindrically averaged power spectra (Giri et al., 2019; Jensen et al., 2013; Ross et al., 2017) and position dependent power spectra (Giri, D'Aloisio, et al., 2019). It also has the capability to find interesting features, such as ionized regions, in (tomographic) image data (Giri et al., 2019, 2018b) and from these to derive statistical quantities, such as size distributions (Giri et al., 2018a) and topological quantities such as the Euler characteristic (Giri et al., 2019). Such statistical characterisations of the data are required when comparing observations with simulations using a Bayesian inference framework in order to derive constraints on model parameters (e.g. Greig & Mesinger, 2015).

## Acknowledgements

We acknowledge the contributions and feedback from Keri Dixon, Hannah Ross, Raghunath Ghara, Ilian Iliev, Catherine Watkinson and Michele Bianco. We are also thankful for support by Swedish Research Council grant 2016-03581.

## References

- Datta, K. K., Mellema, G., Mao, Y., Iliev, I. T., Shapiro, P. R., & Ahn, K. (2012). Light-cone effect on the reionization 21-cm power spectrum. *Monthly Notices of the Royal Astronomical Society*, 424(3), 1877–1891. doi:[10.1111/j.1365-2966.2012.21293.x](https://doi.org/10.1111/j.1365-2966.2012.21293.x)
- DeBoer, D. R., Parsons, A. R., Aguirre, J. E., Alexander, P., Ali, Z. S., Beardsley, A. P., Bernardi, G., et al. (2017). Hydrogen epoch of reionization array (HERA). *Publications of the Astronomical Society of the Pacific*, 129(974), 45001.
- Furlanetto, S. R., Oh, S. P., & Briggs, F. H. (2006). Cosmology at low frequencies: The 21 cm transition and the high-redshift Universe, 433, 181–301. doi:[10.1016/j.physrep.2006.08.002](https://doi.org/10.1016/j.physrep.2006.08.002)
- Giri, S. K. (2019). *Tomographic studies of the 21-cm signal during reionization: Going beyond the power spectrum* (PhD thesis). Department of Astronomy, Stockholm University.
- Giri, S. K., D'Aloisio, A., Mellema, G., Komatsu, E., Ghara, R., & Majumdar, S. (2019). Position-dependent power spectra of the 21-cm signal from the epoch of reionization. *Journal of Cosmology and Astroparticle Physics*, 2019(02), 058–058. doi:[10.1088/1475-7516/2019/02/058](https://doi.org/10.1088/1475-7516/2019/02/058)
- Giri, S. K., Mellema, G., Aldheimer, T., Dixon, K. L., & Iliev, I. T. (2019). Neutral island statistics during reionization from 21-cm tomography. *Monthly Notices of the Royal Astronomical Society*, 489(2), 1590–1605. doi:[10.1093/mnras/stz2224](https://doi.org/10.1093/mnras/stz2224)
- Giri, S. K., Mellema, G., Dixon, K. L., & Iliev, I. T. (2018a). Bubble size statistics during reionization from 21-cm tomography. *MNRAS*, 473, 2949–2964. doi:[10.1093/mnras/stx2539](https://doi.org/10.1093/mnras/stx2539)
- Giri, S. K., Mellema, G., & Ghara, R. (2018b). Optimal identification of H II regions during reionization in 21-cm observations. *Monthly Notices of the Royal Astronomical Society*, 479(4), 5596–5611. doi:[10.1093/mnras/sty1786](https://doi.org/10.1093/mnras/sty1786)
- Greig, B., & Mesinger, A. (2015). 21CMMC: An MCMC analysis tool enabling astrophysical parameter studies of the cosmic 21 cm signal. *Monthly Notices of the Royal Astronomical Society*, 449(4), 4246–4263. doi:[10.1093/mnras/stv571](https://doi.org/10.1093/mnras/stv571)
- Haarlem, M. P. van, Wise, M. W., Gunst, A. W., Heald, G., McKean, J. P., Hessels, J. W. T., Bruyn, A. G. de, et al. (2013). LOFAR: The LOw-Frequency ARray. *Astronomy & Astrophysics*, 556, A2. doi:[10.1051/0004-6361/201220873](https://doi.org/10.1051/0004-6361/201220873)
- Jensen, H., Datta, K. K., Mellema, G., Chapman, E., Abdalla, F. B., Iliev, I. T., Mao, Y., et al. (2013). Probing reionization with LOFAR using 21-cm redshift space distortions. *Monthly Notices of the Royal Astronomical Society*, 435(1), 460–474. doi:[10.1093/mnras/stt1341](https://doi.org/10.1093/mnras/stt1341)
- Jensen, H., Majumdar, S., Mellema, G., Lidz, A., Iliev, I. T., & Dixon, K. L. (2016). The wedge bias in reionization 21-cm power spectrum measurements, 456, 66–70. doi:[10.1093/mnras/stv2679](https://doi.org/10.1093/mnras/stv2679)
- Mellema, G., Koopmans, L. V. E., Abdalla, F. A., Bernardi, G., Ciardi, B., Daiboo, S., Bruyn, A. G. de, et al. (2013). Reionization and the Cosmic Dawn with the Square Kilometre Array. *Experimental Astronomy*, 36, 235–318. doi:[10.1007/s10686-013-9334-5](https://doi.org/10.1007/s10686-013-9334-5)

- Pritchard, J. R., & Loeb, A. (2012, August). 21cm cosmology in the 21st century. doi:[10.1088/0034-4885/75/8/086901](https://doi.org/10.1088/0034-4885/75/8/086901)
- Ross, H. E., Dixon, K. L., Ghara, R., Iliev, I. T., & Mellema, G. (2019). Evaluating the QSO contribution to the 21-cm signal from the Cosmic Dawn. *MNRAS*, 487, 1101–1119. doi:[10.1093/mnras/stz1220](https://doi.org/10.1093/mnras/stz1220)
- Ross, H. E., Dixon, K. L., Iliev, I. T., & Mellema, G. (2017). Simulating the impact of X-ray heating during the cosmic dawn. *Monthly Notices of the Royal Astronomical Society*, 468(4), 3785–3797. doi:[10.1093/mnras/stx649](https://doi.org/10.1093/mnras/stx649)
- Tingay, S. J., Goeke, R., Bowman, J. D., Emrich, D., Ord, S. M., Mitchell, D. A., Morales, M. F., et al. (2013). The Murchison Widefield Array: The Square Kilometre Array Precursor at Low Radio Frequencies. *Publications of the Astronomical Society of Australia*, 30, e007. doi:[10.1017/pasa.2012.007](https://doi.org/10.1017/pasa.2012.007)
- Wayth, R. B., Tingay, S. J., Trott, C. M., Emrich, D., Johnston-Hollitt, M., McKinley, B., Gaensler, B. M., et al. (2018). The Phase II Murchison Widefield Array: Design overview. *Publications of the Astronomical Society of Australia*, 35, 33. doi:[10.1017/pasa.2018.37](https://doi.org/10.1017/pasa.2018.37)