

- BatchPlanet: Batch access and processing of
- ² PlanetScope imagery for spatiotemporal analysis in R
- yiluan Song ^{1,2}, Alexandre R. Bevington ^{3,4}, Ken Reid ¹, Yi Liu ², Jiali
- ⁴ Zhu ¹ ², and Kai Zhu ¹ ²
- 1 Michigan Institute for Data and AI in Society, University of Michigan, Ann Arbor, MI, USA 2 School
- for Environment and Sustainability and Institute for Global Change Biology, University of Michigan, Ann
- ⁷ Arbor, MI, USA 3 Department of Geography, Earth and Environmental Sciences, University of Northern
- British Columbia, Prince George, BC, Canada 4 Ministry of Forests, Province of British Columbia, Prince
- 9 George, BC, Canada

DOI: 10.xxxxx/draft

Software

- Review 🗗
- Repository 🗗
- Archive ♂

Editor: William Gearty ♂ ⑩

Reviewers:

- @ajpelu
- @QuinnAsena

Submitted: 25 July 2025 **Published:** unpublished

License

Authors of papers retain copyright and release the work under a 21 Creative Commons Attribution 4.0 International License (CC BY 4.0)

Summary

The BatchPlanet R package offers a reproducible and scalable workflow for accessing and processing PlanetScope satellite imagery. It is designed for environmental researchers to efficiently perform spatiotemporal analysis on high-resolution remote sensing data. The package streamlines imagery ordering and downloading with the PlanetScope API, retrieval and cleaning of pixel-level time series, and computation of derived metrics like the Enhanced Vegetation Index (EVI) and phenological metrics (e.g., green-up/down time). This tool is particularly valuable for research involving large volumes of imagery across multiple sites and extended time periods.

Statement of Need

PlanetScope imagery provides global, high-resolution (~3-meter), near-daily data, making it valuable for scientific research and monitoring of phenology (Moon et al., 2021), land use change, disaster impacts, and more. While the Planet API facilitates access (Team, 2017), using this data remains challenging due to complex API interactions, limits on large-volume data downloads, and non-trivial processing workflows, which can hinder reproducibility.

Existing options include the official Planet Python SDK, cloud-based platforms like Sentinel Hub and Google Earth Engine (GEE), and the R package planetR. However, the Python and JavaScript used by the first three platforms may be less familiar to R users. Cloud platforms also restrict user control over processing environments and local data storage. Furthermore, existing tools typically require users to write extensive custom scripts for batch downloading and processing across multiple sites, a process often limited by computing and storage resources. BatchPlanet addresses these gaps by providing an R-native tool for batch access and processing of PlanetScope imagery (Table 1). Its streamlined, parallelized functions are designed for scalability, transparency, and reproducibility in scientific data pipelines. This package has supported peer-reviewed research in predicting reproductive phenology in wind-pollinated trees (Song et al., 2025).



Table 1: Table 1. Comparison of BatchPlanet with existing tools for PlanetScope data access and processing.


Feature /	Planet Python	Sentinel	Google Earth Engine	planetR	
Tool	SDK	Hub	(GEE)	(Bevington)	BatchPlanet
Primary Language	Python	Python	JavaScript / Python	R	R
Processing Environment	Local/Cloud	Cloud	Cloud	Local	Local
Data Control & Repro- ducibility	High	Moderate	Low	High	High
Batch Processing	Via script- ing/CLI	Supported for enterprise users	Via scripting	Via scripting	Streamlined
Time Series Analysis Tools	Not supported	Limited	Supported	Not supported	Supported
Interactive Visualization	Not supported	Limited	Supported	Not supported	Supported

Key Features

- The package is tailored for researchers and practitioners who: Conduct time series analyses across spatially dispersed monitoring sites. Work primarily in R and seek alternatives to Python-based tools. Prioritize reproducibility in remote sensing workflows. Use high-performance computing (HPC) infrastructure. Need interactive visualization of PlanetScope imagery and processed data products.
- A major hurdle in using PlanetScope data is the volume of data and the risk of hitting Planet
 API rate limits during mass downloads. This could happen, for example, when a phenological
 study requires years of time series over multiple locations. BatchPlanet provides solutions for
 streamlined batch ordering and downloading. It searches for images by month and automatically
 splits large requests into smaller orders, ensuring complete data retrieval without hitting API
 rate limits. It enables users to specify multiple, spatially distant sites, allowing for efficient,
 parallelized downloading of only the relevant imagery, minimizing unnecessary data volume.
- In addition to ordering and downloading, BatchPlanet facilitates the entire R-native workflow for PlanetScope imagery processing, with a focus on temporal analysis. These include functions to retrieve pixel-level time series data, clean reflectance time series, calculate the Enhanced Vegetation Index (EVI), and compute phenological metrics such as green-up and green-down dates. Apart from the streamlined batch processing functions, BatchPlanet provides individual functions for key steps of the workflow, allowing users to customize their data processing pipelines. BatchPlanet also enables interactive visualization of true color images and EVI time series (Fig. 1, 2).
- These batch functions significantly accelerate the process. For example, images for an approximately 9 km² area over one month were downloaded in 6.7 seconds. The total downloading time for a year's worth of images is comparable due to parallelization across months. Time series retrieval (reflectances, quality masks, and metadata) for 100 coordinates from one month of downloaded images took only 20.9 seconds, which can be parallelized across sites and coordinate groups.



Select Site:






Figure 1: Figure 1. A screenshot of the interactive PlanetScope imagery viewer in the BatchPlanet Song et al. (2025). BatchPlanet: Bakek bases having existing explanations in the BatchPlanet and Software, i VOL?(i ISSUE?), 9268. https://doi.org/10.xxxxx/draft.






Figure 2: Figure 2. Enhanced Vegetation Index (EVI) for three trees in San Joaquin Experimental range (SJER) NEON site with phenological metrics annotated, with indices and metrics calculated and visualized using the BatchPlanet package.

Example Usage

```
64 Install package in R and load package.
```

Read example coordinates.

66 Set download parameters and data directory.

```
setting <- set_planetscope_parameters(
    api_key = set_api_key(),
    item_name = "PSScene",
    asset = "ortho_analytic_4b_sr",
    product_bundle = "analytic_sr_udm2",
    cloud_lim = 0.3,
    harmonized = TRUE
)

download_sample_data()
dir_data <- "sample-data"</pre>
```



```
dir_data_NEON <- file.path(dir_data, "NEON")</pre>
67 Order and download imagery.
   order_planetscope_imagery_batch(
     dir = dir_data_NEON,
     df_coordinates = df_coordinates,
     v_site = c("HARV", "SJER"),
     v_{year} = 2024,
     setting = setting
   download_planetscope_imagery_batch(
     dir = dir data NEON,
     setting = setting,
     num\_cores = 12
   visualize_true_color_imagery_batch(
     dir = dir_data_NEON,
     df_coordinates = df_coordinates
  Retrieve time series at coordinates of interest.
   retrieve_planetscope_time_series_batch(
     dir = dir_data_NEON,
     df_coordinates = df_coordinates,
     num_cores = 12
   )
   df_ts <- read_data_product(</pre>
     dir = dir_data_NEON,
     product_type = "ts"
   visualize_time_series(
     df_ts = df_ts,
     var = "green",
     ylab = "Green reflectance",
     facet_var = "site",
     smooth = F)
69 Clean time series and calculate EVI.
   clean_planetscope_time_series_batch(
     dir = dir data NEON,
     num\_cores = 3,
     calculate_evi = T
   df_clean <- read_data_product(</pre>
     dir = dir_data_NEON,
     product_type = "clean"
   visualize_time_series(
     df ts = df clean,
```



```
var = "evi",
   ylab = "EVI",
   facet_var = "site",
   smooth = T
Calculate phenological metrics.
 df_thres <- set_thresholds(</pre>
   thres_up = c(0.3, 0.4, 0.5),
   thres_down = NULL
 )
calculate phenological metrics batch(
   dir = dir_data_NEON,
   v_site = "SJER",
   v_group = "Quercus",
  df_thres = df_thres,
  var_index = "evi",
   num\_cores = 12
v_id <- c("NEON.PLA.D17.SJER.06001",</pre>
           "NEON.PLA.D17.SJER.06337"
           "NEON.PLA.D17.SJER.06310")
df_doy <- read_data_product(</pre>
   dir = dir_data_NEON,
   product_type = "doy"
df_doy_sample <- df_doy[df_doy$id %in% v_id, ]</pre>
 df_evi <- read_data_product(</pre>
   dir = dir_data_NEON,
   product type = "clean"
df_evi_sample <- df_evi[df_evi$id %in% v_id, ]</pre>
 visualize_time_series(
   df_ts = df_evi_sample,
   df_doy = df_doy_sample,
   var = "evi",
   ylab = "EVI",
   facet_var = "id",
   smooth = T
```

Acknowledgements

- $_{72}$ Yiluan Song was supported by the Eric and Wendy Schmidt AI in Science Postdoctoral
- 73 Fellowship, a Schmidt Sciences program. Kai Zhu and Yiluan Song were supported by the
- National Science Foundation [grant numbers 2306198 (CAREER)]. We thank the Planet team
- 75 for providing access to their API.



References

- Moon, M., Richardson, A. D., & Friedl, M. A. (2021). Multiscale assessment of land surface phenology from harmonized landsat 8 and sentinel-2, PlanetScope, and PhenoCam imagery.

 Remote Sensing of Environment, 266, 112716. https://doi.org/10.1016/j.rse.2021.112716
- Song, Y., Katz, D. S., Zhu, Z., Beaulieu, C., & Zhu, K. (2025). Predicting reproductive phenology of wind-pollinated trees via PlanetScope time series. *Science of Remote Sensing*, 100205. https://doi.org/10.1016/j.srs.2025.100205
- Team, P. (2017). Planet application program interface: In space for life on earth. https://api.planet.com.

