

c14bazAAR: An R package for downloading and preparing C14 dates from different source databases

Clemens Schmid¹, Dirk Seidensticker², and Martin Hinz³

¹ Max Planck Institute for the Science of Human History ² Faculty of Arts and Philosophy, Department of Languages and Cultures, Ghent University ³ Institute of Archaeological Sciences, University of Bern

DOI: [10.21105/joss.01914](https://doi.org/10.21105/joss.01914)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Arfon Smith](#) ↗

Reviewers:

- [@arfon](#)

Submitted: 17 November 2019

Published: 25 November 2019

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

Background

Radiocarbon dating is one of the most important methods for absolute and relative chronological reconstruction of cultural development in prehistoric archaeology (Taylor & Bar-Yosef, 2016). This is true for the intrasite level, for regional comparisons and also in cases, where processes that have a large spatial and temporal reach are to be investigated. Prominent examples for the latter include the ‘neolithization’ of Europe or the ‘Bantu expansion’ in sub-Saharan Africa. They have been extensively analysed with radiocarbon data (e.g. Ammerman & Cavalli-Sforza (1971), Garcin et al. (2018), Jerardino, Fort, Isern, & Rondelli (2014), Lemmen, Gronenborn, & Wirtz (2011), Oslisly et al. (2013), Pinhasi, Fort, & Ammerman (2005), Russell, Silva, & Steele (2014), Weninger et al. (2009)).

Data selection for such models is complex and requires a thorough understanding of the archaeological questions. Most of the time it is not sufficient to include every date that vaguely fits into the context. Some dates have to be deliberately omitted. In order to make this important process of data filtering as transparent and reproducible as possible, the criteria for selection and especially the original data set must be generally accessible and well contextualized. Otherwise, peers can not evaluate the results in a meaningful way.

Fortunately, many archaeological institutions and individual authors are sharing their radiocarbon collections online (e.g. Hinz et al. (2012), Kneisel, Hinz, & Rinne (2013), Courtney Mustaphi (2016), Seidensticker (2016)) – some of them are archives with a long tradition of quality control and maintenance. Also, the boom of the Open Data movement in recent years has led to an increase of publications with raw data in archaeology (e.g. Palmisano, Bevan, & Shennan (2017)). These collections are an important archive for future research questions.

However, the entire data basis is currently highly decentralized and lacks basic standardisation. That results in an effective loss of the possible added value that could result from the intersection of data sets in terms of searchability, error checking and further analysis. The creation of a world-wide and centralised database of radiocarbon dates which could solve these issues is not to be expected for the near future.

Code Summary

c14bazAAR is an R package that attempts to tackle the problem at hand by providing an independent interface to access radiocarbon data and make it available for a reproducible research process: from modelling to publication to scientific discourse. It queries openly available ¹⁴C data archives, but not those behind pay- or login-walls.

The package includes download functions (accessible with the main interface `c14bazAAR::get_c14data()`) that – first of all – acquire databases from different sources online. They then reduce the tables to a set of common variables and store them in a dedicated R S3 class: `c14_date_list`. The `c14_date_list` is based on `tibble::tibble` to integrate well into the R [tidyverse](#) ecosystem. It also establishes standardised data types for the most important variables usually defined to describe radiocarbon data.

Beyond the download functions, `c14bazAAR` contains a multitude of useful helpers that can be applied to objects of class `c14_date_list`. These include methods for bulk calibration of radiocarbon dates with the `Bchron` R package (Haslett & Parnell, 2008), removal of duplicates, estimation of coordinate precision, or conversion to other useful R data types (e.g. `sf::sf` (Pebesma, 2018)). For the classification of sample material `c14bazAAR` provides a manually curated reference list that maps the inconsistent attributions in the source databases to a standardized set of material classes. Such a reference list exists as well to fix the country attribution value of dates – which is especially important in case of missing coordinate information. Methods to determine the source country based on coordinates fail on such dates.

`c14bazAAR` was already used for data acquisition and preparation in at least one research paper: Schmid (2019).

Acknowledgements

The package got valuable code input from several members of the [ISAAKiel group](#) (Initiative for Statistical Analysis in Archaeology Kiel), most notably: Daniel Knitter, David Matzig, Wolfgang Hamer, Kay Schmütz and Nils Müller-Scheeßel.

References

- Ammerman, A. J., & Cavalli-Sforza, L. L. (1971). Measuring the rate of spread of early farming in Europe. *Man*, 6(4), 674–688. doi:[10.2307/2799190](#)
- Courtney Mustaphi, C. (2016). Radiocarbon dates from Eastern Africa in the CARD2.0 format. Harvard Dataverse. doi:[10.7910/DVN/NJLNRJ](#)
- Garcin, Y., Deschamps, P., Ménot, G., de Saulieu, G., Schefuß, E., Sebag, D., Dupont, L. M., et al. (2018). Early anthropogenic impact on Western Central African rainforests 2,600 y ago. *Proceedings of the National Academy of Sciences of the United States of America*, 30, 201715336. doi:[10.1073/pnas.1715336115](#)
- Haslett, J., & Parnell, A. C. (2008). A simple monotone process with application to radiocarbon-dated depth chronologies. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 57(4), 399–418. doi:[10.1111/j.1467-9876.2008.00623.x](#)
- Hinz, M., Furholt, M., Müller, J., Rinne, C., Raetzl-Fabian, D., Sjögren, K.-G., & Wotzka, H.-P. (2012). RADON - Radiocarbon dates online 2012. Central European database of 14C dates for the Neolithic and the Early Bronze Age. *Journal of Neolithic Archaeology*, 14. doi:[10.12766/jna.2012.65](#)
- Jerardino, A., Fort, J., Isern, N., & Rondelli, B. (2014). Cultural diffusion was the main driving mechanism of the neolithic transition in Southern Africa. *PLOS ONE*, 9(12), 1–19. doi:[10.1371/journal.pone.0113672](#)
- Kneisel, J., Hinz, M., & Rinne, C. (2013). Radon-B. <http://radon-b.ufg.uni-kiel.de>.
- Lemmen, C., Gronenborn, D., & Wirtz, K. W. (2011). A simulation of the neolithic transition in Western Eurasia. *Journal of Archaeological Science*, 38(12), 3459–3470. doi:[10.1016/j.jas.2011.08.008](#)

- Oslisly, R., Bentaleb, I., Favier, C., Fontugne, M., Gillet, J. F., & Morin-Rivat, J. (2013). West Central African peoples: Survey of radiocarbon dates over the past 5000 years. *Radiocarbon*, 55(2–3). doi:[10.2458/azu_js_rc.55.16385](https://doi.org/10.2458/azu_js_rc.55.16385)
- Palmisano, A., Bevan, A., & Shennan, S. (2017). Regional demographic trends and settlement patterns in Central Italy: Archaeological sites and radiocarbon dates. *UCL Institute of Archaeology: London, UK*. (2017). doi:[10.14324/000.ds.1575442](https://doi.org/10.14324/000.ds.1575442)
- Pebesma, E. (2018). Simple features for R: Standardized support for spatial vector data. *The R Journal*, 10(1), 439–446. doi:[10.32614/RJ-2018-009](https://doi.org/10.32614/RJ-2018-009)
- Pinhasi, R., Fort, J., & Ammerman, A. J. (2005). Tracing the origin and spread of agriculture in Europe. *PLOS Biology*, 3(12). doi:[10.1371/journal.pbio.0030410](https://doi.org/10.1371/journal.pbio.0030410)
- Russell, T., Silva, F., & Steele, J. (2014). Modelling the spread of farming in the bantu-speaking regions of Africa: An archaeology-based phylogeography. *PLOS ONE*, 9(1), 1–9. doi:[10.1371/journal.pone.0087854](https://doi.org/10.1371/journal.pone.0087854)
- Schmid, C. (2019). Evaluating cultural transmission in Bronze Age burial rites of Central, Northern and Northwestern Europe using radiocarbon data. *Adaptive Behavior*. doi:[10.1177/1059712319860842](https://doi.org/10.1177/1059712319860842)
- Seidensticker, D. (2016). ADRAC. Zenodo. doi:[10.5281/zenodo.61113](https://doi.org/10.5281/zenodo.61113)
- Taylor, R. E., & Bar-Yosef, O. (2016). *Radiocarbon dating: An archaeological perspective*. Routledge. doi:[10.4324/9781315421216](https://doi.org/10.4324/9781315421216)
- Weninger, B., Clare, L., Rohling, E., Bar-Yosef, O., Böhner, U., Budja, M., Bundschuh, M., et al. (2009). The impact of rapid climate change on prehistoric societies during the Holocene in the Eastern Mediterranean. *Documenta Praehistorica*, 36, 7–59. doi:[10.4312/dp.36.2](https://doi.org/10.4312/dp.36.2)