

AQEval: R code for the analysis of discrete change in Air Quality time-series

Karl Ropkins¹, Anthony Walker², Ian Philips¹, Christopher E. Rushton¹, Tony Clark², and James E. Tate¹

¹ Institute for Transport Studies, University of Leeds, Leeds, LS2 9JT, UK ² Joint Air Quality Unit, Department for Transport & Department for Environment, Food and Rural Affairs, Marsham Street London, SW1P 4DF, UK

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Gabriele Bozzola](#)

Reviewers:

- [@eldemet](#)
- [@meenakshi-kushwaha](#)

Submitted: 02 June 2025

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

AQEval (Air Quality Evaluation) is an R package for the routine investigation of discrete changes in air quality time-series. The main functions use break-point/segmentation (BP/S) methods to detect, characterise and quantify change, while other functions build on these to provide a workflow to measure smaller changes and/or changes in more complex environments.

Statement of Need

Authorities responsible for air quality management are typically required to implement and evaluate air quality interventions they adopt (Bradley et al., 2019). These interventions are often costly, disruptive and unpopular (Glazener & Khreis, 2019), and the inherent variability monitoring data hinders impact assessments (Grange & Carslaw, 2019; Jones et al., 2012; Kelly et al., 2011; Pearce et al., 2011). Various methods have been developed to investigate discrete changes in a wide range of time-series (see e.g. Reeves et al., 2007; Truong et al., 2020) and several R (R Core Team & others, 2025) packages have been developed for their use, e.g. bcp (Erdman & Emerson, 2008), changepoint (Killick et al., 2016), segmented (Muggeo & others, 2008), and strucchange (Zeileis et al., 2002). Some have even been applied to air quality time-series (see e.g. Carslaw et al. (2006), Carslaw & Carslaw (2007)). However, many of those tasked with air quality policy assessment, although highly skilled in a wide range of monitoring activities, are unlikely to be able to dedicate sufficient time and resources to the development of in-house expertise in such specialist analyses. AQEval was developed to address this skill gap. It aligns the inputs and outputs of a number of statistical methods to provide a one-package option for anyone interested in using R to routinely investigate change in air quality data. As many air quality professionals already use the R package openair (Carslaw & Ropkins, 2012; Ropkins & Carslaw, 2012) for more conventional analysis and data visualisation, AQEval has also been written using openair coding conventions to reduce the learning-curve typically associated with learning new software.

Sources

AQEval is freely available under General Public License (GPL):

- The latest (stable) release version of AQEval is on the Comprehensive R Archive Network (CRAN) <https://CRAN.R-project.org/package=AQEval>;
- The developers' version and code are publicly on GitHub <https://github.com/karlropkins/AQEval>, where issues or change requests can also be posted; and
- The project website is at <https://karlropkins.github.io/AQEval/>.

40 Analytical Rationale

41 The AQEval Break-Point/Segment (BP/S) methods involve three steps: finding possible
42 'points-of-change', testing these and quantifying 'regions-of-change' about the most likely:

- 43 1. Breaks-points are determined using the strucchange methods of Zeileis and colleagues
44 (Zeileis et al., 2002, 2003). Here, a rolling-window approach is applied: a first subset
45 (time-series window TW_0 in Figure 1a) is selected and linear regression modelled; the
46 window advanced (TW_1 in Figure 1a) and a second model built, and so on through the
47 time-series; then, likely points-of-change assigned by comparing the F-Stat scores of
48 sequential models.
- 49 2. In addition to the standard Bayesian Information Criterion (BIC) testing used by
50 strucchange, AQEval also checks all individual break-points are statistically valid
51 ($p < 0.05$), and down-scores less likely combinations.
- 52 3. Finally, the segmented methods of Muggeo and colleagues (Muggeo, 2003, 2017; Muggeo
53 & others, 2008) are used to determine regions-of-change about break-points. Here, the
54 confidence intervals for the selected break-points are used as start points, and segments
55 assigned by random walk testing about these points as shown in Figure 1b.




Figure 1: Break-point/segment scheme: (a) break-point, and (b) segment modelling about break-points.

56 Figure 2 shows the break-point/segment analysis of an NO_2 time-series from a roadside site
57 where a change event ($\text{ca. } 25 \mu\text{g.m}^{-3}$; 31%) is detected between 2003-01-11 and 2003-02-19.



Figure 2: Standard AQEval break-point/segment analysis (graphical output and report) of an NO_2 1998-2005 time-series from a heavily trafficked roadside in the UK.

58 In some cases changes are small or local air inputs are complex, and time-series may require
59 additional pre-processing to successfully isolate obscured change-events. For these, AQEval uses
60 Generalized Additive Models (GAMs) (using mgcv methods, Wood, 2017, 2025) to subtract
61 associated variance, by default:

$$[pollutant] = s_1(day\ of\ year) + s_2(hour\ of\ day) + te_1(wind\ speed, wind\ direction)$$

$$[pollutant]_{isolated} = ([pollutant] - [pollutant]_{predicted}) + mean(pollutant)$$

63 Where the investigated pollutant concentration, $[pollutant]$, is modelled as a function of *day of year*, *hour of day* and *wind speed* and *direction* using a combination of spline (s_1 and s_2)
64 and tensor (te_1) fit-terms, and the unmodelled component, $[pollutant]_{isolated}$, is estimated as
65 the mean-centred residual of this model.
66

67 Figure 3a shows the break-point analysis of NO₂ from a nearby but less heavily trafficked site
68 where seasonality dominates the time-series, and Figure 3b shows the smaller (ca. 6.6 µg.m⁻³;
69 13%) underlying change-event observed at a similar time to the large change observed at the
70 more heavily trafficked site in Figure 2 using signal isolation and then break-point/segment
71 analysis (2002-09-09 to 2002-12-21 compared with 2003-01-11 and 2003-02-19).




Figure 3: AQEval analysis of NO₂ 1998-2005 time-series from a roadside site where: (a) standard (ambient air) break-point analysis exhibits a near-regular distribution of breaks typical of a site dominated by seasonal factors; and, (b) an underlying change-event is revealed using signal isolation and then break-point/segment analysis.

72 This default correction can also be modified to include other potential confounders, e.g. other
73 frequency terms (e.g. day of week and/or week of year), background contributions (as local

74 variance associated with trends at near-by site not affected by the investigated change), or
 75 proxies for other local contributors (e.g. other meteorological parameters like air temperature,
 76 markers for other sources, etc).

77 Related Outputs

78 The AQEval functions are described, along with worked examples of the code used to generate
 79 Figures 2 and 3, in the [extended package introduction](#). Other work using AQEval include:

- 80 ■ [Ropkins & Tate \(2021\)](#), a peer-reviewed article on the multi-species AQEval analysis of
 81 air quality during the UK COVID-19 lockdown.
- 82 ■ [Ropkins et al \(2022\)](#), a peer-reviewed article on the use of AQEval to measure the NO₂
 83 impact of bus fleet interventions.
- 84 ■ Also Clear Air Zone (CAZ) impact assessment reports include analyses using AQEval,
 85 see e.g.:
 86 — [CAZ Baseline Study](#)
 87 — [First Year Report](#)
 88 — (...)
 89 — [Report archive](#)

90 Acknowledgements

91 Initial AQEval development was funded by the UK Department for Environment, Food and
 92 Rural Affairs (Defra). The authors gratefully acknowledge contributions from colleagues at
 93 University of Leeds, Defra and IPSOS Mori, and internal review by the Defra/Department
 94 for Transport Joint Air Quality Unit (JAQU) Technical Independent Review Panel (T-IRP).
 95 The authors also gratefully acknowledge the work of the R core team and collaborators in
 96 developing and maintaining the open-source statistical language R and associated packages
 97 <http://www.r-project.org/>.

98 The views and opinions expressed herein by the authors are their own and do not necessarily
 99 reflect those of UK Government or any agency thereof.

100 References

- 101 Bradley, N., Dobney, A., Exley, K., Aldridge, J., Craswell, A., Dimitroulopoulou, S., Hodgson,
 102 G., Izon-Cooper, L., Mitchem, L., Mitsakou, C., & others. (2019). *Review of interventions*
 103 *to improve outdoor air quality and public health*. [https://assets.publishing.service.
 104 gov.uk/government/uploads/system/uploads/attachment_data/file/937341/Principal_
 105 interventions_for_local_authorities-air_quality_public_health.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/937341/Principal_interventions_for_local_authorities-air_quality_public_health.pdf)
- 106 Carslaw, D. C., & Carslaw, N. (2007). Detecting and characterising small changes in urban
 107 nitrogen dioxide concentrations. *Atmospheric Environment*, 41(22), 4723–4733. [https:
 108 //doi.org/10.1016/j.atmosenv.2007.03.034](https://doi.org/10.1016/j.atmosenv.2007.03.034)
- 109 Carslaw, D. C., & Ropkins, K. (2012). Openair—an r package for air quality data analysis.
 110 *Environmental Modelling & Software*, 27, 52–61. [https://doi.org/10.1016/j.envsoft.2011.
 111 09.008](https://doi.org/10.1016/j.envsoft.2011.09.008)
- 112 Carslaw, D. C., Ropkins, K., & Bell, M. C. (2006). Change-point detection of gaseous and
 113 particulate traffic-related pollutants at a roadside location. *Environmental Science &
 114 Technology*, 40(22), 6912–6918. <https://doi.org/10.1016/j.scitotenv.2018.10.344>
- 115 Erdman, C., & Emerson, J. W. (2008). Bcp: An r package for performing a bayesian
 116 analysis of change point problems. *Journal of Statistical Software*, 23, 1–13. [http:
 117 //www.jstatsoft.org/v23/i03/](http://www.jstatsoft.org/v23/i03/)

- 118 Glazener, A., & Khreis, H. (2019). Transforming our cities: Best practices towards clean
119 air and active transportation. *Current Environmental Health Reports*, 6, 22–37. <https://doi.org/10.1007/s40572-019-0228-1>
120
- 121 Grange, S. K., & Carslaw, D. C. (2019). Using meteorological normalisation to detect
122 interventions in air quality time series. *Science of the Total Environment*, 653, 578–588.
123 <https://doi.org/10.1016/j.scitotenv.2018.10.344>
- 124 Jones, A. M., Harrison, R. M., Barratt, B., & Fuller, G. (2012). A large reduction in
125 airborne particle number concentrations at the time of the introduction of “sulphur
126 free” diesel and the london low emission zone. *Atmospheric Environment*, 50, 129–138.
127 <https://doi.org/10.1016/j.atmosenv.2011.12.050>
- 128 Kelly, F., Anderson, H. R., Armstrong, B., Atkinson, R., Barratt, B., Beevers, S., Derwent, D.,
129 Green, D., Mudway, I., Wilkinson, P., & others. (2011). The impact of the congestion
130 charging scheme on air quality in london. Part 1. Emissions modeling and analysis of air
131 pollution measurements. *Research Report (Health Effects Institute)*, 155, 5–71.
- 132 Killick, R., Haynes, K., & Eckley, I. (2016). *Changepoint: An r package for changepoint*
133 *analysis. R package version 2.2*. 2. Comprehensive R Archive Network. [https://CRAN.R-](https://CRAN.R-project.org/)
134 [project.org/. https://CRAN.R-project.org/package=changepoint](https://CRAN.R-project.org/package=changepoint)
- 135 Muggeo, V. M. (2003). Estimating regression models with unknown break-points. *Statistics in*
136 *Medicine*, 22(19), 3055–3071.
- 137 Muggeo, V. M. (2017). Interval estimation for the breakpoint in segmented regression: A
138 smoothed score-based approach. *Australian & New Zealand Journal of Statistics*, 59(3),
139 311–322. <https://doi.org/10.1111/anzs.12200>
- 140 Muggeo, V. M., & others. (2008). Segmented: An r package to fit regression models with
141 broken-line relationships. *R News*, 8(1), 20–25. <https://cran.r-project.org/doc/Rnews/>
- 142 Pearce, J. L., Beringer, J., Nicholls, N., Hyndman, R. J., & Tapper, N. J. (2011). Quantifying
143 the influence of local meteorology on air quality using generalized additive models. *Atmos-*
144 *spheric Environment*, 45(6), 1328–1336. <https://doi.org/10.1016/j.atmosenv.2010.11.051>
- 145 R Core Team, R., & others. (2025). *R: A language and environment for statistical computing.*
146 <https://www.R-project.org/>
- 147 Reeves, J., Chen, J., Wang, X. L., Lund, R., & Lu, Q. Q. (2007). A review and comparison of
148 changepoint detection techniques for climate data. *Journal of Applied Meteorology and*
149 *Climatology*, 46(6), 900–915. <https://doi.org/10.32614/RJ-2012-003>
- 150 Ropkins, K., & Carslaw, D. C. (2012). Openair: Data analysis tools for the air quality
151 community. *The R Journal*, 4(1), 20–29. <https://doi.org/10.32614/RJ-2012-003>
- 152 Truong, C., Oudre, L., & Vayatis, N. (2020). Selective review of offline change point detection
153 methods. *Signal Processing*, 167, 107299. <https://doi.org/10.1016/j.sigpro.2019.107299>
- 154 Wood, S. N. (2017). *Generalized additive models: An introduction with r*. chapman; hall/CRC.
- 155 Wood, S. N. (2025). *Mgcv: Mixed GAM computation vehicle with automatic smoothness*
156 *estimation. R package version 1.9-3*. Comprehensive R Archive Network. [https://CRAN.R-](https://CRAN.R-project.org/)
157 [project.org/.https://CRAN.R-project.org/package=mgcv](https://CRAN.R-project.org/package=mgcv)
- 158 Zeileis, A., Kleiber, C., Krämer, W., & Hornik, K. (2003). Testing and dating of structural
159 changes in practice. *Computational Statistics & Data Analysis*, 44(1-2), 109–123. [https://doi.org/10.1016/S0167-9473\(03\)00030-6](https://doi.org/10.1016/S0167-9473(03)00030-6)
160
- 161 Zeileis, A., Leisch, F., Hornik, K., & Kleiber, C. (2002). Strucchange: An r package for testing
162 for structural change in linear regression models. *Journal of Statistical Software*, 7, 1–38.
163 <http://www.jstatsoft.org/v07/i02/>