

# PICNIC: an open-source Python library for preprocessing of dynamic Positron Emission Tomography (PET) brain imaging data

Francesca Zanderigo<sup>1,2</sup>, Eric Hauser<sup>1,2</sup>, Mike Schmidt<sup>1,2</sup>, Elizabeth A Bartlett<sup>1,2</sup>, and Jeffrey M Miller<sup>1,2</sup>

<sup>1</sup> Department of Psychiatry, Columbia University Medical Center, New York, NY, United States  
<sup>2</sup> Molecular Imaging and Neuropathology Area, New York State Psychiatric Institute, New York, NY, United States

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Marcel Stimberg](#)

## Reviewers:

- [@ghisvail](#)
- [@pjmark](#)

Submitted: 24 July 2025

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

## Summary

PICNIC (Pipeline Initializing Container for Neuro-Imaging Computations) is an open-source Python-based coding library that includes modular wrappers for most standard preprocessing steps for quantification of Positron Emission Tomography (PET) brain imaging data. PICNIC uniquely allows full, transparent modular control over the selection of strategy/software package and associated parameters for each preprocessing step throughout the pipeline. Therefore, the user is provided with the flexibility to select different methods depending on the target of interest and/or radiotracer properties, study population, or other preferences. These settings can be saved as a text-based “input deck” file to support consistent preprocessing across scans, participants, and projects. Furthermore, PICNIC supports freezing software versions within Docker containers, ensuring robust reproducibility.

PICNIC does not require Brain Imaging Data Structure (BIDS) source-level data, but instead takes reconstructed PET static or dynamic data in industry standard imaging filetypes (e.g., DICOM, Nifti, ANALYZE), and converts them to be BIDS-compliant ([Knudsen et al., 2020](#)). PICNIC employs the most commonly used brain imaging software packages, including Advanced Normalization Tools ([Avants et al., 2008](#)), Analysis of Function NeuroImages ([Cox, 1996](#)), FSL ([Jenkinson et al., 2012](#)), FreeSurfer ([Fischl, 2012](#)), dcm2niix ([Li et al., 2016](#)), and SPM ([Laboratory, 2020](#)). PICNIC was designed with PET in mind, but includes steps involving data from multiple imaging modalities, most prominently structural magnetic resonance imaging (sMRI). PICNIC preprocessing capabilities include image reorientation (to force all images into a standard orientation regardless of scanner defaults), brain extraction (to perform skull-stripping of associated sMRI images), tissue segmentation (to classify sMRI voxels within the brain boundary as grey matter vs. white matter vs. cerebrospinal fluid), automatic anatomical parcellation (to identify regions of interest using any available atlas/parcellation), post-reconstruction rigid-body motion-correction (to correct inter-frame participant motion over the duration of the scan), PET-sMRI co-registration (with 20 different PET-sMRI rigid-body transformations that are automatically ranked based on mutual information), and extraction of time-activity-curves (curves that represent the PET signal over the duration of the scan in delineated regions of interest or in each brain voxel).

PICNIC can be run from the command line or through the provided graphical user interface. An html summary output with interactive quality control features allows the user to inspect, comment, and approve/disapprove each module for each participant upon completion. Pre-loaded templates have been provided for commonly used workflows. The user can add, remove or edit pre-processing steps to fully customize their pipeline with regards to their dataset.

PICNIC currently supports over 100,000 module/parameter combinations. PICNIC is designed for in vivo brain investigations by expert PET researchers, and has already been applied to preprocess data for recent scientific publications from our Lab (Bartlett, Patil, et al., 2023, 2024; Bartlett, Zanderigo, Stanley, et al., 2023; Bartlett, Zanderigo, Ogden, et al., 2023; Bartlett, Herzog, et al., 2024; Bartlett, Sublette, et al., 2024; Graves et al., 2024; Herzog et al., 2024, 2025; Mann et al., 2022; Matheson et al., 2024; Miller et al., 2022) (<https://www.columbiapsychiatry.org/research-labs/brain-imaging-lab>). PICNIC's modular nature enables extensive customization of preprocessing for a variety of brain PET studies, and its design may allow future extension to preprocessing of data from imaging modalities other than PET. The source code for PICNIC is available at the following link: <https://github.com/bil-mind-nyspi/PICNIC>.

## Statement of need

PET is a valuable tool used in research and clinical settings to noninvasively image the human brain in vivo. PET can quantify up to nanomolar levels of specific components of brain metabolic and neurochemical processes, thus providing information on the distribution of specific biological targets, such as receptors and enzymes, or on the uptake of specific compounds into the brain, like glucose and polyunsaturated fatty acids. Rigorous preprocessing of the raw data captured by the PET scanner (Norgaard et al., 2019) is key to obtaining the most accurate estimates of the distribution of a biological target or uptake of a substance using PET. Although a few software packages already exist (Funck et al., 2018; Karjalainen et al., 2020; Routier et al., 2021) that perform most of the required preprocessing steps, a fully open-source library that can modularly and flexibly combine appropriate preprocessing strategies depending on the study design at hand is still missing.

## References

- Avants, B. B., Epstein, C. L., Grossman, M., & Gee, J. C. (2008). Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. *Medical Image Analysis*, 12(1), 26–41. <https://doi.org/10.1016/j.media.2007.06.004>
- Bartlett, E. A., Herzog, S., Zanderigo, F., & others. (2024). *Translocator protein (TSPO) binding assessed with 11C-ER176 PET in major depression and healthy volunteers*.
- Bartlett, E. A., Patil, S., Matheson, G., & others. (2024). *Dual input function modeling of nonbrain 11C-ER176 signal for comparison of translocator protein (TSPO) central and peripheral binding*.
- Bartlett, E. A., Patil, S., Schmidt, M., & others. (2023). *In Vivo [11C]ER176 Brain and Non-Brain Translocator Protein (TSPO) PET Binding in Major Depressive Disorder and Healthy Volunteers*.
- Bartlett, E. A., Sublette, M. E., Melhem, N., Mann, J. J., & Zanderigo, F. (2024). *Quantification of 11C-ER176 TSPO distribution volume with one blood sample: optimization and validation of simultaneous estimation (SIME) of input function*.
- Bartlett, E. A., Zanderigo, F., Ogden, R. T., Sublette, M. E., Mann, J. J., & Miller, J. M. (2023). *In vivo investigation of brain translocator protein (TSPO) and monoamine oxidase A (MAO-A) relationship in healthy volunteers*.
- Bartlett, E. A., Zanderigo, F., Stanley, B., & others. (2023). *Relationships of Childhood and Recent Stressors with in Vivo Brain Translocator Protein (TSPO) in Major Depressive Disorder and Healthy Volunteers*.

- 88 Cox, R. W. (1996). AFNI: software for analysis and visualization of functional magnetic  
89 resonance neuroimages. *Computers and Biomedical Research*, 29(3), 162–173. <https://doi.org/10.1006/cbmr.1996.0014>  
90
- 91 Fischl, B. (2012). FreeSurfer. *Neuroimage*, 62(2), 774–781. [https://doi.org/10.1016/j.  
92 neuroimage.2012.01.021](https://doi.org/10.1016/j.neuroimage.2012.01.021)
- 93 Funck, T., Larcher, K., Toussaint, P. J., Evans, A. C., & Thiel, A. (2018). APPIAN:  
94 Automated Pipeline for PET Image Analysis. *Frontiers in Neuroinformatics*, 12, 64.  
95 <https://doi.org/10.3389/fninf.2018.00064>
- 96 Graves, P., Miller, J. M., Zanderigo, F., & others. (2024). *Relationships Between the Gut and  
97 Oral Microbiome and Neuroinflammation in Major Depressive Disorder*.
- 98 Herzog, S., Bartlett, E. A., Galfalvy, H. C., & others. (2024). *11C-ER176 PET Imaging of  
99 Translocator Protein (TSPO) Binding in Major Depression: Relationship to Severity of  
100 Real-Time Suicidal Ideation and Negative Affect in the Context of Naturalistic Stressors*.
- 101 Herzog, S., Bartlett, E. A., Zanderigo, F., & others. (2025). Neuroinflammation, Stress-  
102 Related Suicidal Ideation, and Negative Mood in Depression. *JAMA Psychiatry*, 82(1),  
103 85–93. <https://doi.org/10.1001/jamapsychiatry.2024.3543>
- 104 Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M. (2012).  
105 Fsl. *Neuroimage*, 62(2), 782–790. <https://doi.org/10.1016/j.neuroimage.2011.09.015>
- 106 Karjalainen, T., Tuisku, J., Santavirta, S., & others. (2020). Magia: Robust Automated  
107 Image Processing and Kinetic Modeling Toolbox for PET Neuroinformatics. *Frontiers in  
108 Neuroinformatics*, 14, 3. <https://doi.org/10.3389/fninf.2020.00003>
- 109 Knudsen, G. M., Ganz, M., Appelhoff, S., & others. (2020). Guidelines for the content  
110 and format of PET brain data in publications and archives: A consensus paper. *Journal  
111 of Cerebral Blood Flow & Metabolism*, 40(8), 1576–1585. [https://doi.org/10.1177/  
112 0271678X20905433](https://doi.org/10.1177/0271678X20905433)
- 113 Laboratory, F. I. (2020). *Statistical Parametric Mapping, version 12 (SPM12)*. [https://www.  
114 fil.ion.ucl.ac.uk/spm/software/spm12/](https://www.fil.ion.ucl.ac.uk/spm/software/spm12/)
- 115 Li, X., Morgan, P. S., Ashburner, J., Smith, J., & Rorden, C. (2016). The first step for  
116 neuroimaging data analysis: DICOM to NIfTI conversion. *Journal of Neuroscience Methods*,  
117 264, 47–56. <https://doi.org/10.1016/j.jneumeth.2016.03.001>
- 118 Mann, J. J., Galfalvy, H., Haghighi, F., & others. (2022). *Neuroinflammation and Acute  
119 Psychopathology*.
- 120 Matheson, G. J., Zanderigo, F., Bartlett, E. A., & others. (2024). *Biologically-Informed  
121 Hierarchical Approach for Genotype Correction in Translocator Protein PET and its  
122 Application to [11C]ER176 Data*.
- 123 Miller, J. M., Zanderigo, F., Ogden, R. T., Sublette, M. E., & Mann, J. J. (2022). *Relationships  
124 between Depression, Suicidal Ideation, and Neuroinflammation as Measured Using Positron  
125 Emission Tomography with [11C]ER-176*.
- 126 Norgaard, M., Ganz, M., Svarer, C., & others. (2019). Optimization of preprocessing  
127 strategies in Positron Emission Tomography (PET) neuroimaging: A [(11)C]DASB PET  
128 study. *Neuroimage*, 199, 466–479. <https://doi.org/10.1016/j.neuroimage.2019.05.055>
- 129 Routier, A., Burgos, N., Diaz, M., & others. (2021). Clinica: An Open-Source Software  
130 Platform for Reproducible Clinical Neuroscience Studies. *Frontiers in Neuroinformatics*, 15,  
131 689675. <https://doi.org/10.3389/fninf.2021.689675>