The Journal of Open Source Software

AutoHist.jl: A Julia package for fast and automatic histogram construction

Oskar Høgberg Simensen [1]

1 Scientific Computing Center, Karlsruhe Institute of Technology

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Mehmet Hakan Satman

Reviewers:

- @PieterjanRobbe
- @ajinkya-k

Submitted: 17 July 2025
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

`AutoHist.jl` is a Julia (Bezanson et al., 2017) package for fitting histograms to univariate data, with an automatic and data-based selection of the histogram partition. It currently supports 7 irregular and 12 regular automatic histogram procedures from the statistical literature. Additionally, `AutoHist.jl` provides extensions for `Plots.jl` (Christ et al., 2023) and `Makie.jl` (Danisch & Krumbiegel, 2021), allowing for simple visualization of the resulting histogram fits.

Statement of need

To this day, histograms remain one of the most widely used nonparametric density estimators. Their popularity is in no doubt due to their simplicity and interpretability, and they are routinely used by practitioners with limited mathematical backgrounds as a means of visualizing the distribution of data sets. Although the construction of a histogram for a given interval partition is a simple task, the quality of the resulting density estimate is very sensitive to the choice of bins. As a result, the task of designing automatic histogram procedures, where the number of bins and their location are chosen in a data-driven fashion, has recieved considerable interest in the statistics community. Despite advances in our understanding of different bin selection rules, many popular software libraries only include a limited number of simple rules for selecting a regular histogram partition, and typically do not provide any support for automatic irregular histogram construction. `AutoHist.jl` fills this gap by providing a fast implementation of state-of-the-art regular and irregular bin selection algorithms from the statistics literature. A complete overview of the bin selection procedures that have been implemented so far is given in **Table 1**.

Table 1: Implemented bin selection procedures so far. For methods with type=both, a regular and an irregular variant of the criterion is supported.

Rule	Type (regular/irregular)	Reference
Sturges	regular	Sturges (1926)
Scott	regular	Scott (1979)
Freedman-Diaconis	regular	Freedman & Diaconis (1981)
AIC	regular	Hall (1990)
BIC	regular	Davies et al. (2009)
Birgé-Rozenholc	regular	Birgé & Rozenholc (2006)
MDL	regular	Hall & Hannan (1988)
Wand	regular	Wand (1997)
RRH, Knuth	regular	Knuth (2019)
RMG penA	irregular	Rozenholc et al. (2010)
RMG penB	irregular	Rozenholc et al. (2010)
RMG penR	irregular	Rozenholc et al. (2010)

Rule	Type (regular/irregular)	Reference
RIH	irregular	Simensen et al. (2025)
NML	both	Kontkanen & Myllymäki (2007)
L2CV	both	Rudemo (1982)
KLCV	both	Hall (1990), Simensen et al. (2025)

We note that some automatic histogram selection rules have been implemented in Julia, typically as part of plotting libraries. The `Plots.jl` package includes support for certain plug-in rules used in standard histograms, whereas the `StatsPlots.jl` library (Christ et al., 2023) provides functionality for plotting equal-area histograms. The `R` package `histogram` (Mildenberger et al., 2019) supports some regular and irregular histogram methods, but their implementation covers fewer criteria than ours.

Installation and usage

The `AutoHist` package is part of the Julia general registry, and can as such be installed via the built-in package manager,

```
using Pkg
Pkg.add("AutoHist")
```

To illustrate the basic use of the software, we fit a histogram based on the irregular Bayesian criterion and a regular histogram based on AIC to a standard normal random sample of size $n = 10^6$.

```
using AutoHist, Random, Distributions

n = 10^6
x = rand(Xoshiro(1812), Normal(), n)          # synthetic data

h_irr = fit(AutomaticHistogram, x, RIH())     # fit an irregular histogram
h_reg = fit(AutomaticHistogram, x, AIC())     # fit a regular histogram
```

The call to the `fit` method returns an object of type `AutomaticHistogram`, with fields recording the chosen histogram partition, estimated density and bin counts. `AutoHist.jl` provides plotting recipes for `Plots.jl` and `Makie.jl`, which allows the user to easily visualize the fit. Here, we show how to plot the irregular and the regular histogram fitted in the aabove code snippet using `Makie`.

```
import CairoMakie, Makie # using the CairoMakie backend

fig = Makie.Figure(size=(670, 320))
ax1 = Makie.Axis(fig[1, 1], title="Irregular histogram", xlabel="x",
                 ylabel="Density")
ax2 = Makie.Axis(fig[1, 2], title="Regular histogram", xlabel="x")
p_irr = Makie.plot!(ax1, h_irr, alpha=0.4, color="black")
p_reg = Makie.plot!(ax2, h_reg, alpha=0.4, color="red")
fig
```

DRAFT



Figure 1: Plot of the irregular and regular histogram fit to the standard normal sample.

References

Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. (2017). Julia: A fresh approach to numerical computing. *SIAM Review*, *59*(1), 65–98. https://doi.org/10.1137/141000671

Birgé, L., & Rozenholc, Y. (2006). How many bins should be put in a regular histogram. *ESAIM: Probability and Statistics*, *10*, 24–45. https://doi.org/10.1051/ps:2006001

Christ, S., Schwabeneder, D., Rackauckas, C., Borregaard, M. K., & Breloff, T. (2023). *Plots.jl - a user extendable plotting API for the julia programming language*. https://doi.org/10.5334/jors.431

Danisch, S., & Krumbiegel, J. (2021). Makie.jl: Flexible high-performance data visualization for Julia. *Journal of Open Source Software*, *6*, 3349. https://doi.org/10.21105/joss.03349

Davies, P. L., Gather, U., Nordman, D., & Weinert, H. (2009). A comparison of automatic histogram constructions. *ESAIM: Probability and Statistics*, *13*, 181–196. https://doi.org/10.1051/ps:2008005

Freedman, D., & Diaconis, P. (1981). On the histogram as a density estimator: L2 theory. *Zeitschrift Für Wahrscheinlichkeitstheorie Und Verwandte Gebiete*, *57*, 453–476. https://doi.org/10.1007/BF01025868

Hall, P. (1990). Akaike's information criterion and Kullback–Leibler loss for histogram density estimation. *Probability Theory and Related Fields*, *85*, 449–467. https://doi.org/10.1007/BF01203164

Hall, P., & Hannan, E. J. (1988). On stochastic complexity and nonparametric density estimation. *Biometrika*, *75*, 705–714. https://doi.org/10.1093/biomet/75.4.705

Knuth, K. H. (2019). Optimal data-based binning for histograms and histogram-based probability density models. *Digital Signal Processing*, *95*. https://doi.org/10.1016/j.dsp.2019.102581

Kontkanen, P., & Myllymäki, P. (2007). MDL histogram density estimation. *Proceedings of Machine Learning Research*, *2*, 219–226. https://proceedings.mlr.press/v2/kontkanen07a.html

Mildenberger, T., Rozenholc, Y., & Zasada, D. (2019). *Histogram: Construction of regular and irregular histograms with different options for automatic choice of bins*. https://doi.org/10.32614/cran.package.histogram

Rozenholc, Y., Mildenberger, T., & Gather, U. (2010). Combining regular and irregular histograms by penalized likelihood. *Computational Statistics & Data Analysis*, *54*, 3313–3323. https://doi.org/10.1016/j.csda.2010.04.021

Rudemo, M. (1982). Empirical choice of histograms and kernel density estimators. *Scandinavian Journal of Statistics*, *9*, 65–78. https://www.jstor.org/stable/4615859

Scott, D. W. (1979). On optimal and data-based histograms. *Biometrika*, *66*, 605–610. https://doi.org/10.1093/biomet/66.3.605

Simensen, O. H., Christensen, D., & Hjort, N. L. (2025). Random irregular histograms. *arXiv Preprint*. https://doi.org/10.48550/arXiv.2505.22034

Sturges, H. A. (1926). The choice of a class interval. *Journal of the American Statistical Association*, *21*, 65–66. https://doi.org/10.1080/01621459.1926.10502161

Wand, M. P. (1997). Data-based choice of histogram bin width. *The American Statistician*, *51*, 59–64. https://doi.org/10.2307/2684697

DRAFT