

SpotiPy: A Modular Python Pipeline for Solar Spot Analysis and Tracking

Saqib Sumra^{1,2}, Emily Joe Löbnitz^{1,2}, Alexander G. M. Pietrow¹,
and Carsten Denker¹

¹ Leibniz Institute for Astrophysics Potsdam (AIP), Germany ² University of Potsdam, Germany ¶
Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: ↗

Submitted: 07 July 2026

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#)).

Summary

Solar activity manifests itself in many different ways, from the near-omnipresent network to the more confined active regions, which appear as groups of faculae, spots, and pores in the photosphere, and as filaments and prominences in the chromosphere. Our ever-improving physical understanding and astronomical instrumentation has made these features the subject of study for several centuries ([Cretignier et al., 2024](#); [Palumbo et al., 2024](#); [Solanki, 2003](#)).

In recent years, more focus has been placed on the consistent tracking and observations of active regions with space-based facilities such as the Hinode (Solar-B) Mission ([Kosugi et al., 2007](#)), the Solar Dynamics Observatory (SDO, [Pesnell et al., 2012](#)), the Interface Region Imaging Spectrograph (IRIS, [De Pontieu et al., 2014](#)) and the Chinese H α Solar Explorer (CHASE, [Li et al., 2022](#)), but also in high resolution from the ground with facilities such as the Swedish Solar Telescope (SST, [Scharmer et al., 2003](#)) and the GREGOR solar telescope ([Schmidt et al., 2012](#)), resulting in strong progression of our understanding of these features.

SpotiPy is a Python package developed for feature identification and tracking in full-disk solar observations. The framework has been primarily designed for use with data from the Helioseismic and Magnetic Imager (HMI, [Scherrer et al., 2012](#)) and the Atmospheric Imaging Assembly (AIA, [Lemen et al., 2012](#)) aboard SDO, although the methods are not instrument-specific. The pipeline enables the identification and tracking of arbitrary fields of view over time. While initial applications have focused on photospheric features such as sunspots and plages, the implemented procedures are equally valid for other types of solar features.

Corresponding author: Alexander G.M. Pietrow (apietrow@aip.de)

Statement of Need

While general-purpose solar physics libraries such as SunPy ([SunPy Community et al., 2020](#)) provide functionality for data access, coordinate system handling, and file I/O, they do not aim to offer an integrated pipeline for the tracking of solar active regions. In particular, tasks such as time-series download, cross-instrument alignment between observations, and consistent image preprocessing are typically handled through custom scripts by individual researchers, which can be difficult to reproduce and maintain across projects. While foundational libraries such as astropy ([Astropy Collaboration et al., 2022](#)) handle the underlying coordinate transformations and instrument-specific packages such as aiapy ([Barnes et al., 2020](#)) provide excellent calibration routines for SDO/AIA data, they lack the specific, multi-instrument feature-tracking integration that SpotiPy provides.

SpotiPy addresses this practical gap by providing a modular framework that supports coordinated retrieval of large datasets from HMI and AIA, sub-pixel alignment of multi-

instrument observations, and preprocessing steps such as limb darkening correction. While the package was designed with feature tracking and active region analysis in mind, its modular components can be applied to other full-disk studies or as standalone functions within existing pipelines, enabling reuse and sustained development. SpotiPy was developed to facilitate high-throughput, reproducible analysis of solar features and has already been successfully employed to investigate differential rotation of sunspots (Löbnitz et al., 2025) and center-to-limb variations (CLV) of sunspots, faculae, and the solar network (Pietrow et al., 2026).

Functionality

SpotiPy is organized into five independent modules — (1) Data downloading, (2) Data Alignment, (3) Limb Darkening Correction, (4) Segmentation, and (5) Tracking — that can be used together as a complete pipeline or individually within existing workflows.

1. Data downloading

SpotiPy provides a wrapper around Sunpy's Fido framework that handles JSOC email authentication and batch retrieval of HMI and AIA time series. Because the module queries JSOC directly, it can retrieve any available data series. The most common SDO data products used for feature tracking are listed in Table 1.

Table 1: HMI and AIA full-disk data products and passbands accessible via the downloading module.

Instrument	Observable	Supported JSOC Series
HMI	Continuum Intensity	hmi.Ic_45s, hmi.Ic_720s
HMI	Continuum Intensity (Flattened)	hmi.Ic_nolimbDark_720s
HMI	Line width	hmi.Lw_45s, hmi.Lw_720s
HMI	Line depth	hmi.Ld_45s, hmi.Ld_720s
HMI	LOS Magnetograms	hmi.M_45s, hmi.M_720s
HMI	Dopplergrams	hmi.V_45s, hmi.V_720s
AIA	Continuum Intensity (4500 Å)	aia.lev1_vis_1h
AIA	Ultraviolet (1600 Å, 1700 Å)	aia.lev1_uv_24s
AIA	Extreme UV (94 Å, 131 Å, 193 Å, 211 Å, 304 Å, 335 Å)	aia.lev1_euv_12s

To retrieve a time series, users must configure the following core parameters:

- **series:** the HMI or AIA data product identifier (e.g., "hmi.Ic_720s")
- **start_date:** start of the time series in ISO 8601 format (e.g., "2019-04-07T18:00:00")
- **days:** duration of the time series in days
- **cadence_h:** time step between downloaded frames in hours
- **email:** a JSOC-registered email address required for data export

While `download_series` is a general-purpose wrapper capable of retrieving any JSOC data series, the following example demonstrates fetching the specific HMI continuum and AIA 1700 Å data required for the active region pipeline:

```
# Download HMI continuum intensity and AIA 1700 Å time series
from spotipy.downloading import download_series

hmi_files = download_series(
    series="hmi.Ic_720s",
    start_date="2019-04-07T18:00:00",
```

```

days=11,
cadence_h=6,
email="user@institute.edu",
out_dir="data/HMI"
)

aia_files = download_series(
    series="aia.lev1_uv_24s",
    start_date="2019-04-07T18:00:00",
    days=11,
    cadence_h=6,
    email="user@institute.edu",
    out_dir="data/AIA",
    wavelength_angstrom=1700 # Required to avoid downloading all 10 AIA channels
)

```

69 2. Data Alignment

70 SpotiPy automates the co-alignment of multi-instrument images using World Coordinate
 71 Systems (WCS) and fast interpolation via the reproject package (Robitaille et al., 2020). The
 72 alignment module automatically handles the different FITS file structures (e.g., data stored in
 73 primary versus secondary header data units (HDUs)) across different SDO data products. This
 74 ensures that HMI and AIA observations share a common pixel scale of 0.6 arcseconds per pixel,
 75 as shown in Figure 1.

```

# Reproject AIA onto the HMI pixel grid and orientation
from spotipy.aligning import align_images
align_images(aia_path, hmi_reference_path, output_path)

```




Figure 1: Reprojection of AIA 1700 Å intensity onto the HMI pixel grid. Left: original AIA grid. Center: HMI reference. Right: AIA reprojected to HMI grid.

76 3. Limb Darkening Correction

77 This function implements radial-profile subtraction to remove the center-to-limb intensity
 78 profile for HMI continuum and AIA 1600 Å and 1700 Å observations, which is essential for
 79 reliable intensity thresholding across the full solar disk. Limb darkening is a radiative transfer
 80 effect induced by viewing geometry that allows the observer to look deeper into the atmosphere
 81 at the center of the solar disk than at the limb, where the line of sight passes through the
 82 atmosphere at an angle. The decreasing temperature with height in the solar photosphere
 83 results in a darker limb and a brighter center (Foukal, 2004; Gray, 2005). Removing this
 84 gradient is therefore necessary before applying uniform intensity thresholds across the disk.

85 The correction is illustrated in Figure 2 and Figure 3, where a radial profile is generated and
 86 the limb radius is extracted directly from the FITS header parameters. After removal of this

87 profile a mostly flat disk is left, occasionally with artifacts due to large active regions (see
88 [Figure 3](#)). However, these have not been shown to affect thresholding methods.

```
# Remove limb darkening using radial profile subtraction
from spotipy.limbdarkening_removal import remove_limb_darkening, get_header_geometry
(cx, cy), r_pix = get_header_geometry(header)
aia_corrected = remove_limb_darkening(aia_raw, center=(cx, cy), radius_pix=r_pix)
```




Figure 2: AIA 1700 Å intensity before (left) and after (right) limb darkening correction.




Figure 3: Radial intensity profile of the solar disk (left) and its derivative (right). The left panel displays the raw intensity profile overlaid with a 5th-order polynomial fit describing the CLV. The dashed vertical line marks the limb radius extracted from the FITS header parameters (1564 pixels). The derivative profile on the right illustrates the sharp intensity drop at the limb boundary.

89 4. Segmentation

90 SpotiPy uses intensity thresholding combined with morphological operations to generate binary
91 masks for umbra, penumbra, and extended spot regions from HMI continuum intensity, and
92 for plage, network, and quiet Sun regions from AIA 1700 Å ([Shen et al., 2018](#); [Verma et](#)
93 [al., 2018](#)). The segmentation is performed using intensity thresholding combined with size-
94 based classification. These thresholds were empirically calibrated to 8–43% of the normalized
95 quiet-Sun intensity for the umbra, 59–94% for the penumbra, > 15% for the network, and
96 > 20% for plage. These values are broadly consistent with those reported in previous studies

(Chapman et al., 1989; Györi, 1998; Hoeksema et al., 2014; Kiess et al., 2014; Preminger et al., 2001; Solanki, 2003).

To distinguish between network and plage, contiguous regions exceeding the network intensity threshold are further classified based on their area: namely structures larger than 450 pixels are identified as plage, while smaller structures are classified as network. We note that this introduces a center-to-limb inconsistency, as a fixed pixel threshold corresponds to a larger physical area at smaller μ . However, the chosen value has been found to provide a robust performance across the disk.

The segmentation masks can be applied across all five HMI observables simultaneously — continuum intensity (hmi.Ic_720s), Doppler velocity (hmi.V_720s), magnetic flux density (hmi.M_720s), line depth (hmi.Ld_720s), and line width (hmi.Lw_720s) — enabling studies of the CLV within individual solar features.

Figure 4 shows the full-disk segmentation map, while Figure 5 shows a ROI of an individual sunspot with umbral and penumbral masks.

```
# Generate segmentation masks using intensity thresholding
from spotipy.segmentation import get_masks, get_aia_masks

hmi_masks = get_masks(
    hmi_norm,                    # normalized HMI continuum intensity map
    disk_mask=ondisk,           # boolean mask selecting the solar disk
    umbra_range=(10, 55),       # umbra: 8-bit scale (~8-43% of quiet Sun)
    penumbra_range=(75, 120),   # penumbra: 8-bit scale (~59-94% of quiet Sun)
    cleanup=True                # applies connected-component filter
)

aia_masks = get_aia_masks(
    aia_flat,                   # limb-darkening-corrected AIA image
    spot_mask=hmi_masks["spot"], # main sunspot mask to exclude
    all_dark_mask=hmi_masks["all_dark"], # includes smaller pores for exclusion
    plage_excess_pct=20.0,
    qs_tol_pct=15.0,
    solar_center=(cx, cy),      # required for polar-slice QS reference
    r_pix=r_pix,
    full_disk_aia=aia_flat
)
```




Figure 4: Full-disk segmentation map derived from HMI continuum intensity and AIA 1700 Å intensity. Left: HMI continuum intensity. Center: AIA 1700 Å. Right: segmentation map showing quiet Sun (gray), network (blue), plage (gold), and sunspot (red). Excluded regions such as pores and regions beyond the limb are shown in black.




Figure 5: Sunspot segmentation of active region NOAA 12738 at central meridian passage. Left: HMI continuum ROI. Center: AIA 1700 Å intensity ROI. Right: Combined segmentation map showing umbra (red), penumbra (orange), plage (gold), network (blue), and quiet Sun (gray). Small, isolated dark pores excluded (black) to ensure the analysis remains focused on the primary active region.

5. Tracking

The tracking module allows for consistent feature tracking at an arbitrary cadence, allowing for the study of a region and its parameters as it makes its way across the solar disk, and allows for the calculation of differential rotation rates.

This module is based on the processes laid out in (Löbnitz et al., 2025). A core functionality is based on a (SunPy Community et al., 2020) model to compute the expected position of a solar feature over a time series for a given start position, which can be selected via an interactive clicker or by passing starting coordinates. The resulting coordinates are returned and allow for the isolation of an ROI by cropping out a window around it for each time step. To account for local motions and deviations from the rotation model, the pipeline provides an optional recentering function that refines the feature position centering the window around the darkest feature in the cropped image. This is done by smoothening the ROI with a Gaussian and then finding the center of mass of the largest feature. The extracted ROIs can be visualized in time-summed strips, as shown in Figure 6. This series of crops can then be used for segmentation maps (see Segmentation) or further analysis.

```
# Track a solar feature and generate a time-summed visual strip
from spotipy.tracking import track_spots, refine_centering, strip
# 1. Calculate physical tracks from an initial position
tracks = track_spots(hmi_files, hmi_dir, x_start, y_start)
# 2. Refine tracks frame-by-frame to center on the umbra
refined_tracks = []
for i, frame in enumerate(hmi_files):
    # (Internal logic: Load frame, convert arcsec to pixels)
    # new_cx, new_cy = refine_centering(data_crop, rx0, ry0, frame_size)
    # refined_tracks.append(convert_to_arcsec(new_cx, new_cy))
    pass
# 3. Generate a time-summed strip using the refined coordinates
strip_img, _ = strip(
    series=hmi_files,
    directory=hmi_dir,
    tracks=refined_tracks,
    strip_height_arcsec=400.0,
    frame_size=400,
    overlay=True,
    animate=False
)
```




Figure 6: Time-summed strip of active region NOAA 12738 travelling across the solar disk, with tracked bounding boxes overlaid at each time step. The color of the bounding boxes, from purple to yellow, indicates the progression of time.

The determination of feature rotation rates can be performed using either the masks produced by the [Segmentation](#) module or an internal masking process specifically designed for sunspots. Both approaches identify features such as sunspots at each time step. The centers of the sunspot masks are then determined either from image moments or, for approximately circular features, by fitting a minimal-area ellipse. The resulting positions are tracked over time and finally transformed into the heliographic coordinate system using observational parameters such as the solar radius, tilt, and distance obtained from the FITS header. The heliographic coordinate system expresses points on the solar surface by latitude and longitude ([Thompson, 2006](#)). This transformation is necessary to correct for projection effects caused by the spherical geometry, as illustrated in [Figure 7](#). After deprojection, the tracked positions are used to calculate rotation rates, which are expressed in the Carrington reference frame ([Carrington, 1863](#)). This conversion follows the method described in ([Balthasar, 1979](#)). The Carrington System rotates as a rigid sphere along with the surface of the Sun, ensuring comparability between rotation studies based on SpotiPy and others.




Figure 7: Rotation rate of a single feature (a sunspot in active region NOAA 12738) during its passage across the solar disk. The left panel shows the calculated rotation rate without correcting for projection effects, while the right panel shows the deprojected rotation rate in the Carrington reference frame using the two available methods (image moments and ellipse fitting). Near the solar limb, the apparent motion of the feature corresponds to only a few pixels between consecutive frames. After deprojection, these small positional uncertainties translate into larger uncertainties in the calculated rotation rate. The breaks in the plotted curves occur where the Carrington longitude wraps from 360° to 0°.

Acknowledgements

We acknowledge the use of data from the Solar Dynamics Observatory, provided by NASA. HMI and AIA data were accessed via the JSOC data export service. AP was supported by grant PI 2102/1-1 from the Deutsche Forschungsgemeinschaft (DFG). DeepL Write was used for assistance with proofreading and copy-editing.

References

- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., & others. (2022). The Astropy project: Sustaining and growing a community-oriented open-source project and the latest major release (v5.0) of the core package. *The Astrophysical Journal*, 935(2), 167. <https://doi.org/10.3847/1538-4357/ac7c74>
- Balthasar, H. (1979). *Differentielle Rotation und meridionale Bewegungen der Sonnenflecken in den Jahren 1940–1968* [Diploma Thesis]. Georg-August-Universität Göttingen.
- Barnes, W. T., Cheung, M. C. M., Bobra, M. G., & others. (2020). aiapy: A Python package for analyzing Solar Dynamics Observatory Atmospheric Imaging Assembly data. *Journal of Open Source Software*, 5(55), 2801. <https://doi.org/10.21105/joss.02801>
- Carrington, R. C. (1863). *Observations of the spots on the sun: From November 9, 1853, to March 24, 1861, made at Redhill*. Williams; Norgate.
- Chapman, G. A., Herzog, A. D., Laico, D. E., & others. (1989). Photometric measurements of solar irradiance variations due to sunspots. *The Astrophysical Journal*, 343, 547–556. <https://doi.org/10.1086/167728>
- Cretignier, M., Pietrow, A. G. M., & Aigrain, S. (2024). Stellar surface information from the Ca II H&K lines – i. Intensity profiles of the solar activity components. *Monthly Notices of the Royal Astronomical Society*, 527(2), 2940–2962. <https://doi.org/10.1093/mnras/stad3292>
- De Pontieu, B., Title, A. M., Lemen, J. R., & others. (2014). The Interface Region Imaging Spectrograph (IRIS). *Solar Physics*, 289, 2733–2779. <https://doi.org/10.1007/s11207-014-0485-y>
- Foukal, P. V. (2004). *Solar astrophysics* (2nd ed.). Wiley-VCH.
- Gray, D. F. (2005). *The observation and analysis of stellar photospheres* (3rd ed.). Cambridge University Press. <https://doi.org/10.1017/CBO9781316036570>
- Győri, L. (1998). Automation of area measurement of sunspots. *Solar Physics*, 180(1–2), 109–130. <https://doi.org/10.1023/A:1005081621268>
- Hoeksema, J. T., Liu, Y., Hayashi, K., & others. (2014). The Helioseismic and Magnetic Imager (HMI) vector magnetic field pipeline: Overview and performance. *Solar Physics*, 289(9), 3483–3530. <https://doi.org/10.1007/s11207-014-0516-8>
- Kiess, C., Rezaei, R., & Schmidt, W. (2014). Properties of sunspot umbrae observed in cycle 24. *Astronomy & Astrophysics*, 565, A52. <https://doi.org/10.1051/0004-6361/201321119>
- Kosugi, T., Matsuzaki, K., Sakao, T., & others. (2007). The Hinode (Solar-B) mission: An overview. *Solar Physics*, 243, 3–17. <https://doi.org/10.1007/s11207-007-9014-6>
- Lemen, J. R., Title, A. M., Akin, D. J., & others. (2012). The Atmospheric Imaging Assembly (AIA) on the Solar Dynamics Observatory (SDO). *Solar Physics*, 275, 17–40. <https://doi.org/10.1007/s11207-011-9776-8>
- Li, C., Fang, C., Li, Z., & others. (2022). The Chinese H α Solar Explorer (CHASE) mission: An overview. *Science China Physics, Mechanics & Astronomy*, 65(8), 289602. <https://doi.org/10.1007/s11433-022-1893-3>

- 184 Löbnitz, E. J., Pietrow, A. G. M., Chakraborty, H., & others. (2025). Differential rotation of
185 solar α -sunspots and implications for stellar light curves. *Astronomy & Astrophysics*, 703,
186 A187. <https://doi.org/10.1051/0004-6361/202555654>
- 187 Palumbo, M. L., Saar, S. H., & Haywood, R. D. (2024). Characterizing solar center-to-
188 limb radial-velocity variability with SDO. *The Astrophysical Journal*, 973(1), 11. <https://doi.org/10.3847/1538-4357/ad445a>
- 189
- 190 Pesnell, W. D., Thompson, B. J., & Chamberlin, P. C. (2012). The Solar Dynamics Observatory
191 (SDO). *Solar Physics*, 275, 3–15. <https://doi.org/10.1007/s11207-011-9841-3>
- 192 Pietrow, A. G. M., Sumra, S., Petit dit de la Roche, D. J. M., & others. (2026). Center-to-limb
193 variations of solar active regions: Observations of spots, faculae, and network in the 6173
194 Å continuum. *Astronomy & Astrophysics*. <https://arxiv.org/abs/2606.28887>
- 195 Preminger, D. G., Walton, S. R., & Chapman, G. A. (2001). Solar feature identification
196 using contrasts and contiguity. *Solar Physics*, 202, 53–62. [https://doi.org/10.1023/A:](https://doi.org/10.1023/A:1011896413891)
197 [1011896413891](https://doi.org/10.1023/A:1011896413891)
- 198 Robitaille, T., Deil, C., & Ginsburg, A. (2020). *Reproject: Python-based astronomical image*
199 *reprojection*. Zenodo. <https://doi.org/10.5281/zenodo.1162674>
- 200 Scharmer, G. B., Bjelksjö, K., Korhonen, T. K., & others. (2003). The 1-meter Swedish solar
201 telescope. *Proc. SPIE*, 4853, 341–350. <https://doi.org/10.1117/12.460377>
- 202 Scherrer, P. H., Schou, J., Bush, R. I., & others. (2012). The Helioseismic and Magnetic
203 Imager (HMI) investigation for the Solar Dynamics Observatory (SDO). *Solar Physics*, 275,
204 207–227. <https://doi.org/10.1007/s11207-011-9834-2>
- 205 Schmidt, W., Lühe, O. von der, Volkmer, R., & others. (2012). The GREGOR solar telescope
206 on Tenerife. *Astronomische Nachrichten*, 333, 796–809. [https://doi.org/10.1002/asna.](https://doi.org/10.1002/asna.201211725)
207 [201211725](https://doi.org/10.1002/asna.201211725)
- 208 Shen, Z., Diercke, A., & Denker, C. (2018). Calibration of full-disk He I 10830 Å filtergrams
209 of the Chromospheric Telescope. *arXiv*. <https://doi.org/10.1002/asna.201813536>
- 210 Solanki, S. K. (2003). Sunspots: An overview. *The Astronomy and Astrophysics Review*, 11,
211 153–286. <https://doi.org/10.1007/s00159-003-0018-4>
- 212 SunPy Community, Barnes, W. T., Bobra, M. G., & others. (2020). The SunPy project: Open
213 source development and status of the version 1.0 core package. *The Astrophysical Journal*,
214 890, 68. <https://doi.org/10.3847/1538-4357/ab4f7a>
- 215 Thompson, W. T. (2006). Coordinate systems for solar image data. *Astronomy & Astrophysics*,
216 449, 791–803. <https://doi.org/10.1051/0004-6361:20054262>
- 217 Verma, M., Denker, C., Balthasar, H., & others. (2018). High-resolution imaging and
218 near-infrared spectroscopy of penumbral decay. *Astronomy & Astrophysics*, 614, A2.
219 <https://doi.org/10.1051/0004-6361/201731801>